

SOME ASPECTS OF PARTICLE PHYSICS IN THERMAL BACKGROUND

**THESIS SUBMITTED TO
JADAVPUR UNIVERSITY
FOR THE DEGREE OF
DOCTOR OF PHILOSOPHY (SCIENCE)**

By

**INDRAJIT MITRA
THEORY GROUP
SAHA INSTITUTE OF NUCLEAR PHYSICS
KOLKATA**

2002

ACKNOWLEDGMENTS

I am indebted to Prof. Palash Baran Pal for guidance, encouragement, collaboration and numerous stimulating discussions. I am also grateful to Prof. José F. Nieves for collaboration and many valuable discussions via e-mail.

Interactions with many other members, both past and present, of the Theory Group of SINP, including Dr. Ananda DasGupta, Prof. Asit K. De, Prof. Binayak Dutta-Roy, Dr. Abhee K. Dutt-Mazumder, Prof. A. Harindranath and Prof. Samir Mallik, have helped me learn quantum field theory and thermal field theory. I thank them all.

I express my gratitude to all members of the Theory Group for their help and cooperation. The Research Fellows, in particular, have helped me in every possible way during my stay here.

Last, but not the least, I thank my mother for her support and understanding, without which this work could not have been done.

TABLE OF CONTENTS

	Page
Acknowledgments	ii
List of Figures	vi
Chapters:	
1. Introduction	1
1.1 Importance of particle physics in thermal background	1
1.2 The problems considered in this thesis and their motivations	2
1.3 Outline of the thesis	5
1.4 Some notations and conventions	6
2. Thermal field theory	7
2.1 Introduction	7
2.2 Thermodynamic equilibrium	7
2.3 Time-ordered thermal propagators	10
2.3.1 Scalar field	10
2.3.2 Fermion field	12
2.3.3 Photon field	14
2.4 Additional vertex and propagators	15
3. Linearized theory of gravity	17
3.1 Introduction	17
3.2 Graviton	18
3.3 Photons in curved space	22
3.4 Spinors in curved space	23
3.5 Interaction vertices	28
3.5.1 Photon-photon-graviton vertex	29
3.5.2 Fermion-fermion-graviton vertex	30

3.5.3	Fermion-fermion-photon-graviton vertex	32
3.5.4	Important features of the vertex factors	33
4.	Charged leptons in thermal QED	35
4.1	Dispersion relation and inertial mass	35
4.2	Spinor and wavefunction renormalization factor	39
4.3	Effect of charge conjugation invariance	43
4.4	Relations when background is not C invariant	47
4.5	Calculations at $O(e^2)$	48
4.5.1	Self-energy	48
4.5.2	Calculation of inertial mass	51
4.5.3	Calculation of wavefunction renormalization factor	53
5.	Gravitational couplings of charged leptons in a medium	55
5.1	Introduction	55
5.2	Gravitational mass	55
5.3	Operational definition at $O(e^2)$	59
5.4	Gravitational vertex	62
5.4.1	Irreducible diagrams	62
5.4.2	Bubble diagrams given in Fig. 5.1	64
5.4.3	Tadpole diagrams given in Fig. 5.2	66
5.4.4	Additional remarks	69
5.5	Calculation of the gravitational mass	70
5.5.1	Terms with the photon distribution from Fig. 5.1	70
5.5.2	Terms with the electron distribution from Fig. 5.1	72
5.5.3	Terms from Fig. 5.2	74
5.5.4	Summary	75
5.6	Results for particular cases	76
5.6.1	Classical electron gas and classical proton gas	77
5.6.2	Degenerate electron gas and classical proton gas	80
6.	Some aspects of high-temperature QED	83
6.1	Introduction	83
6.2	One-loop electron self-energy in a general linear covariant gauge	84
6.3	Equation governing gauge dependence of dispersion relation at one loop, and general considerations	88
6.4	Gauge independence of one-loop dispersion relation at momenta much larger than eT	89
6.5	Gauge independence of effective mass at one-loop	91
6.5.1	Subtleties at zero momentum	91
6.5.2	Formula for the effective mass	91

6.5.3	Investigation of gauge independence	93
7.	Conclusions	96
Appendices:		
A.	Field quantization using discrete momenta	99
B.	Full propagator and wavefunction renormalization factor in vacuum	101
C.	Charge conjugation and the 1PI gravitational vertex	104
D.	Transversality of the gravitational vertex	107
E.	Apparently ill-defined contributions to the gravitational vertex	111
E.1	The B1 contribution	111
E.2	The A2 contribution	114
E.3	The X contribution	117
E.4	Alternative evaluation of the B1 and A2 contributions	119
F.	Thermal contribution to the equation for the effective mass at high temperature	123
Bibliography		126
List of Publications		130

LIST OF FIGURES

Figure		Page
3.1	Feynman rule for the coupling of the photon with the graviton. $C_{\mu\nu\lambda\rho}(k, k')$ is given in Eq. (3.79).	29
3.2	Feynman rule for the tree-level gravitational vertex of a fermion. $V_{\lambda\rho}(p, p')$ is given in Eq. (3.95).	32
3.3	Diagrams of $O(e\kappa)$ with two fermions, one photon and one graviton in the external lines. These are the tree-level diagrams for the photoproduction of gravitons off electrons.	33
4.1	One-loop diagrams for the self-energy of a charged lepton ℓ in a medium. . .	48
5.1	Bubble diagrams for the one-loop gravitational vertex of charged leptons. These diagrams contribute in a background of photons and electrons. The braided line represents the graviton.	63
5.2	Tadpole diagrams for the one-loop gravitational vertex of charged leptons. These diagrams contribute to the vertex of any charged lepton in a background of electrons and nucleons.	63
5.3	The one-loop diagrams that contribute to order κ to the photon tadpole in a medium, in the presence of a static and homogeneous ($q = 0$) gravitational potential. The fermion loop involves a sum over all the species of fermions present in the medium.	67

CHAPTER 1

Introduction

1.1 Importance of particle physics in thermal background

The study of elementary particles in vacuum is based on quantum field theory. However, there are situations in which it is necessary to consider these particles in a non-trivial background consisting of a large number of particles at finite temperature and/or density. Such temperatures and densities of significance occur in stellar interiors and were also present in the early universe. Another relevant situation occurs in the laboratory during heavy-ion collisions. The formalism needed for the study of these situations is provided by thermal field theory, also known as finite temperature field theory [1, 2, 3].

From a theoretical point of view (as we will see in the course of the present work), two major features are induced by the background medium. These are the occurrence of an additional four-vector in the form of the four-velocity of the medium, and the breaking of discrete symmetries like C , P and T even when the underlying Lagrangian respects these symmetries. As a result, self-energies, vertex functions and other Green's functions involve new tensor structures and associated form factors, together with a relaxation of some of the constraints restricting the form factors. A third important feature induced by the medium (also to be encountered in the present work) is the breaking of flavour symmetry. All these features give rise to rich physics which was absent in the vacuum. A modified self-energy, for example, gives rise to modification of the dispersion relation (energy-momentum relation). This, in turn, among other things, can open up the phase space, and allow processes which were kinematically forbidden in the vacuum to take place in the medium.

Let us now turn to some specific illustrations of changes in the properties of elementary particles due to a thermal background. Gauge symmetry spontaneously broken in vacuum can be restored by finite temperature corrections. The masses of the fermions and gauge

bosons of the Standard Model vanish when this happens [4]. In the phase of restored chiral invariance, one finds the generation of thermal effective masses of the particles, different effective masses of left-handed and right-handed leptons, and different dispersion relations of transverse and longitudinal modes of gauge bosons [5, 6]. Modification of photon dispersion relations leads to Cerenkov radiation (emission of a photon) by even a single neutrino in a medium [7]. The dispersion relation of neutrinos become flavour dependent in a medium containing electrons but not the other charged leptons [8, 9]. In such a flavor asymmetric background, the electromagnetic [10] and gravitational [11] interactions of the neutrinos in response to external fields are also flavor dependent, while the radiative decay of a massive neutrino is strongly enhanced compared to the decay in vacuum (as the leptonic GIM mechanism becomes inoperative) [12]. When neutrinos travel in a background of electrons with a static external magnetic field also present, the latter causes an anisotropy of the neutrino dispersion relation [10]. An entirely new property of the Standard Model neutrino acquired in a background of electrons is an effective electric charge [13]. A *CPT*-asymmetric background medium causes rotation of the plane of polarization of light passing through it [14], a phenomenon known as optical activity.

There are a number of extremely important applications of background-induced particle physics, some of which we now mention. The temperature dependence of the effective potential for the Higgs field gives rise to the inflationary scenario in the early universe [15]. The flavour dependence of the neutrino dispersion relation in matter leads to the MSW effect, with consequences for the solar neutrino problem [16]. The decay of a photon propagating through a medium, known as plasmon, into neutrino-antineutrino pair is the major cooling mechanism in white dwarfs and red giants [17, 18]. The anisotropy of neutrino dispersion relation in matter with background magnetic field has been suggested as a possible explanation for the large birth velocities of pulsars [19].

1.2 The problems considered in this thesis and their motivations

The major problem considered in this thesis is the gravitational interaction of charged leptons in a medium. Gravitational interactions (in the vacuum) are universal in the sense that the ratio of the inertial and the gravitational masses of any particle is a constant. This fact, expressed in the form of the equivalence principle, is one of the basic axioms of the

general theory of relativity. Although this is a feature of the theory at the classical level, it has been shown by Donoghue, Holstein and Robinett (DHR) [20] that the corresponding linearized quantum theory of gravity also respects this ratio, at least to $O(\alpha)$.

However, it was also shown by DHR [20, 21] that this property is lost when the particles are in the presence of a thermal background rather than the vacuum. To arrive at this idea, the inertial and the gravitational masses must be defined in the context of quantum field theory. We consider in Chapters 4 and 5 their precise definitions in terms of the particle propagator and the gravitational vertex, which we will need in the subsequent work. For the moment, let us denote the mass in vacuum by m , the inertial mass by M and the gravitational mass by M' , and summarize the results of DHR. These authors considered the case of the electron, and calculated M_e and M'_e to $O(\alpha)$, in a background with a temperature $T \ll m_e$ and zero chemical potential. Thus, the background contained only photons, but not electrons or any other matter particles. Although in those calculations only the case of the electron was considered explicitly, the results are equally applicable to other charged leptons, namely, the muon and the tau lepton. Thus for any charged lepton ℓ , the DHR results imply

$$M_\ell = m_\ell + \frac{\alpha\pi T^2}{3m_\ell}, \quad (1.1)$$

$$M'_\ell = m_\ell - \frac{\alpha\pi T^2}{3m_\ell}, \quad (1.2)$$

$$\frac{M'_\ell}{M_\ell} = 1 - \frac{2\alpha\pi T^2}{3m_\ell^2} \quad (1.3)$$

to $O(\alpha)$. Therefore, not only the inertial and gravitational masses of a given charged lepton cease to be equal when the background effects are taken into account, but, in addition, the ratio of these two quantities is no longer the same for all the particles; i.e., universality is lost as well.

Now, the DHR results, as mentioned before, were obtained by considering only photons in the thermal background. But in a matter background with a non-zero chemical potential (such as the Sun), one can no longer neglect the matter particles in medium, even when $T \ll m_e$. It is then possible that the matter contribution to the charged lepton masses will dominate over the photon background contribution (the latter being, as the above results indicate, quite small for $T \ll m_e$). Moreover, since a medium with non-zero chemical potential is asymmetric with respect to charge conjugation, the matter-induced corrections will not be the same for the corresponding antileptons.

Motivated by these considerations, in this work we calculate the leading matter-induced QED corrections to the inertial and gravitational masses of charged fermions in a medium that consists of a photon background and a matter background of electrons and nucleons [22]. These represent the dominant corrections for charged leptons and antileptons (weak interaction corrections being negligible in comparison). For strongly interacting particles such as the quarks, gluon exchange corrections are expected to be even stronger and our results will not apply. After finding the general expressions for the matter-induced corrections in a generic matter background, we give explicit formulas for the corrections in terms of the macroscopic parameters of the background medium for a few special cases of the background gases.

While these calculations are performed for low temperatures and densities, in the last part of the thesis, we turn to a background at high temperature. We consider high-temperature QED with exact chiral invariance and at zero chemical potential, and investigate gauge independence of the one-loop electron dispersion relation in this situation.

It is well-known that the one-loop electron dispersion relation is gauge independent to leading order in the temperature T . This was demonstrated by Weldon [5], by showing that the leading part of the one-loop electron self-energy (the self-energy being the field-theoretic input to the dispersion relation) goes like T^2 and is gauge independent, while the gauge dependent part goes like T . Now, a somewhat similar situation occurs for the one-loop self-energy of neutrinos in material medium, which is gauge independent at $O(1/M_W^2)$ but gauge dependent when the $O(1/M_W^4)$ terms are included. However, it has been shown [9] that the one-loop dispersion relation following from the latter is still independent of the gauge parameter. This raises the possibility that the gauge dependence of the part of the one-loop electron self-energy subleading in T may also not show up in the one-loop dispersion relation. This possibility is explored in the present work in two different limiting cases [23]. One of these cases is the limit of momenta much larger than eT . The other is the zero momentum limit, when we investigate the gauge independence of the effective mass.

It may be mentioned in this context that the demonstration of gauge independence of the properties of elementary particles in material medium has turned out to be of extreme importance in thermal field theory. In particular, the gauge dependence of the gluon damping rate at one loop was a long-standing problem. In Ref. [24], the Ward identities determining the gauge dependence of the gluon dispersion relations were deduced and

used to prove gauge independence in a self-consistent perturbative expansion. In Refs. [25] and [26], it was shown that such a consistent expansion requires resummation of the hard thermal loops. The use of this idea finally led to a gauge-independent gluon damping rate to leading order in the coupling constant [27]. It will be seen that the gauge dependence identities of Ref. [24], although dealing with gluons, are still relevant for the case studied by us.

1.3 Outline of the thesis

In Chapter 2, we introduce the real-time formulation of thermal field theory. After working out the thermal averages of bilinears of ladder operators, we deduce the time-ordered thermal propagators for various fields. Additional vertex and propagators in the real-time formulation are then briefly mentioned.

In Chapter 3, we discuss the linearized theory of gravity. The linearization of the metric and how it gives rise to the graviton are explained. Photons and spinors in curved space are discussed, leading to the QED Lagrangian of curved space. This Lagrangian is then linearized, and the Feynman rules for the lowest order interaction of the graviton with photons and fermions are deduced.

In Chapter 4, we consider charged leptons in thermal QED. General expressions for the dispersion relation, the inertial mass, the spinor and the wavefunction renormalization factor are obtained at $O(e^2)$ for a fermion as well as an antifermion. Relations between the fermion and antifermion properties are then deduced for both C -symmetric and C -asymmetric backgrounds. Explicit $O(e^2)$ calculation of the inertial masses and wavefunction renormalization factors of charged leptons and antileptons are performed.

In Chapter 5, we turn to gravitational couplings in a medium. We show in detail how the gravitational mass is determined from the gravitational vertex function, and obtain an operational formula for the gravitational mass at $O(e^2)$. We then calculate the gravitational vertex of the leptons, and use it to find the gravitational masses of charged leptons and antileptons in a medium. Since the terms involving the fermion distribution function cannot be evaluated exactly, we evaluate the corrections in two different limits, viz., the classical and the degenerate limits for the electron gas.

In Chapter 6, we consider some aspects of high-temperature QED. The expression for the one-loop self-energy of the electron in a general linear covariant gauge is deduced.

The gauge dependence relation for the one-loop electron dispersion relation is obtained, and gauge independence is proved for momenta much larger than eT . After obtaining a formula for the effective mass of the electron, gauge independence of the one-loop effective mass is investigated.

In Chapter 7, we present our conclusions.

1.4 Some notations and conventions

We now summarize some notations and conventions which will be frequently used in this thesis. Although these will be mentioned again in the appropriate places, it was felt that a separate summary of them may be useful as a ready reference in going through the present work.

Throughout this work, we will denote four-momenta by lower case letters, and the corresponding three-momenta by upper case letters. For example, we write $p^\mu = (p^0, \vec{P})$. The magnitude of $\vec{P}$ is denoted by P . Also, we use the notation $E_P \equiv \sqrt{P^2 + m^2}$, where m is the mass of the particle concerned.

The mass in the vacuum, the inertial mass in a medium and the gravitational mass in a medium are denoted by m , M and M' respectively. They come with various subscripts: f denotes a generic fermion, p a proton, ℓ a charged lepton and $\bar{\ell}$ a charged antilepton. Further, ℓ can be e or μ or τ .

The notations for the thermal corrections to m , as defined in Eqs. (4.104) and (5.84), contain the additional subscripts 1 and 2. Here 1 stands for the contribution from the photon background, and 2 stands for the contribution from the electron background. The symbols 1 and 2 are used in this way for various other medium effects also. Thus, it is used to denote the contributions from a diagram: the B1 contribution, for example, denotes the photon background contribution to the diagram B, and the A2 contribution denotes the electron background contribution to the diagram A.

Finally, in our work on high-temperature QED, the effective mass of the electron is denoted by M , while M_0 denotes the effective mass to leading order in temperature.

CHAPTER 2

Thermal field theory

2.1 Introduction

Thermal field theory is the theoretical framework for studying a large number of quantum mechanical and relativistic particles in thermodynamic equilibrium. The two well-known versions of this subject are the imaginary-time formalism [1, 2, 3, 28, 29] and the real-time formalism [2, 3, 28, 29, 30], of which we are going to discuss only the latter. The real-time formalism has the following advantages. The calculations are manifestly covariant, the effects of the medium are separated out at the very beginning from the part already present in the vacuum, and it can be easily understood that the medium does not introduce any new ultraviolet divergence. The particular approach to the real-time formalism to be discussed by us is the canonical approach [30, 31]. This approach involves a generalization of the canonical quantization formalism of ordinary quantum field theory, by defining the Green's functions as thermal averages instead of vacuum expectation values.

2.2 Thermodynamic equilibrium

Thermal field theory makes use of quantum field theory as well as statistical mechanics. As mentioned in Sec. 2.1, we are going to use the canonical quantization formalism of quantum field theory. For doing statistical mechanics, we have to choose a particular ensemble according to our convenience. In the present case, each of the systems making up the ensemble is in some state of the Fock space, and the calculations are the simplest (see below) if the ensemble contains all the states of the Fock space, without any restriction on the total number of particles in a system. Consequently, we consider the grand canonical

ensemble, which is characterized by the temperature

$$T \equiv \frac{1}{\beta} \quad (2.1)$$

and the chemical potential μ . (We have chosen our units such that the Boltzmann constant is equal to unity.) The thermal expectation value (ensemble average) of any operator $\mathcal{O}$ is

$$\langle \mathcal{O} \rangle = \text{Tr}(\rho_G \mathcal{O}), \quad (2.2)$$

with the density operator ρ_G in the grand canonical ensemble given by

$$\rho_G = \frac{e^{-\beta J}}{\text{Tr } e^{-\beta J}}, \quad (2.3)$$

where

$$J = H - \mu Q. \quad (2.4)$$

Here H is the normal-ordered hamiltonian, and Q the normal-ordered charge operator, i.e., the total number operator for the particle minus that for the antiparticle.

We shall now consider the case of free fields. Let us begin with a real scalar field, for which (and, in general, for any self-conjugate field) $\mu = 0$. For the sake of convenience, we shall consider quantization with discrete momenta in this section. (From Sec. 2.3 onward, we shall consider quantization with continuous momenta; Appendix A relates the two approaches.) We therefore have

$$H = \sum_{\vec{K}} E_K a_K^\dagger a_{\vec{K}}. \quad (2.5)$$

Here

$$E_K \equiv \sqrt{K^2 + m^2}, \quad (2.6)$$

m being the mass of the quantum of the scalar field, and $a_K^\dagger a_{\vec{K}}$ is the number operator for quanta of momentum $\vec{K}$.

We are now going to find out the thermal average of the bilinears of the ladder operators. First consider

$$\langle a_{\vec{P}}^\dagger a_{\vec{P}} \rangle = \frac{\text{Tr}[a_{\vec{P}}^\dagger a_{\vec{P}} \exp(-\beta \sum_{\vec{K}} E_K a_K^\dagger a_{\vec{K}})]}{\text{Tr}[\exp(-\beta \sum_{\vec{K}} E_K a_K^\dagger a_{\vec{K}})]}, \quad (2.7)$$

where Eqs. (2.2)-(2.5) (with $\mu = 0$) were used. The trace is in the Fock space, which is spanned by the direct product of the eigenstates of the number operators $a_{\vec{K}}^\dagger a_{\vec{K}}$ for the various modes $\vec{K}$. Therefore for any operator $\mathcal{O}'$,

$$\text{Tr } \mathcal{O}' = \sum_{n_{\vec{K}_1}, n_{\vec{K}_2}, \dots} \langle n_{\vec{K}_1} | \langle n_{\vec{K}_2} | \dots \mathcal{O}' \dots | n_{\vec{K}_2} \rangle | n_{\vec{K}_1} \rangle. \quad (2.8)$$

Here $\langle n_{\vec{K}_1} | \langle n_{\vec{K}_2} | \dots$ denotes the direct product of the states $\langle n_{\vec{K}_1} |$, $\langle n_{\vec{K}_2} |$, $\dots$. Also, since we are considering a bosonic field, the sum over each eigenvalue $n_{\vec{K}}$ runs from zero to ∞ .

Use of Eq. (2.8) in both the numerator and denominator of Eq. (2.7) leads to

$$\begin{aligned} \langle a_{\vec{P}}^\dagger a_{\vec{P}} \rangle &= \frac{\sum_{n_{\vec{P}}} n_{\vec{P}} e^{-\beta E_P n_{\vec{P}}} \prod_{\vec{K} \neq \vec{P}} \sum_{n_{\vec{K}}} e^{-\beta E_K n_{\vec{K}}}}{\prod_{\vec{K}} \sum_{n_{\vec{K}}} e^{-\beta E_K n_{\vec{K}}}} \\ &= \frac{\sum_{n_{\vec{P}}} n_{\vec{P}} e^{-\beta E_P n_{\vec{P}}}}{\sum_{n_{\vec{P}}} e^{-\beta E_P n_{\vec{P}}}} \end{aligned} \quad (2.9)$$

$$= -\frac{\partial}{\partial(\beta E_P)} \ln \sum_{n_{\vec{P}}} e^{-\beta E_P n_{\vec{P}}} \quad (2.10)$$

The sum over $n_{\vec{P}}$ is an infinite geometric series, and equals $1/(1 - e^{-\beta E_P})$. So

$$\langle a_{\vec{P}}^\dagger a_{\vec{P}} \rangle = \frac{1}{e^{\beta E_P} - 1}, \quad (2.11)$$

which is the Bose distribution function. Using $[a_{\vec{P}}, a_{\vec{P}}^\dagger]_- = 1$, this leads to

$$\langle a_{\vec{P}} a_{\vec{P}}^\dagger \rangle = 1 + \frac{1}{e^{\beta E_P} - 1}. \quad (2.12)$$

Also using Eqs. (2.2)-(2.5) and Eq. (2.8), it follows at once that

$$\langle a_{\vec{P}}^\dagger a_{\vec{P}'} \rangle = \langle a_{\vec{P}} a_{\vec{P}'}^\dagger \rangle = 0 \quad \text{for } \vec{P} \neq \vec{P}' \quad (2.13)$$

$$\langle a_{\vec{P}} a_{\vec{P}'} \rangle = \langle a_{\vec{P}}^\dagger a_{\vec{P}'}^\dagger \rangle = 0. \quad (2.14)$$

For a complex scalar field with the number operators $a_{\vec{K}}^\dagger a_{\vec{K}}$ for the particle and $b_{\vec{K}}^\dagger b_{\vec{K}}$ for the antiparticle of momentum $\vec{K}$, Eq. (2.4) gives

$$J = \sum_{\vec{K}} [(E_K - \mu) a_{\vec{K}}^\dagger a_{\vec{K}} + (E_K + \mu) b_{\vec{K}}^\dagger b_{\vec{K}}], \quad (2.15)$$

The Fock space states are now the direct product of the eigenstates of $a_{\vec{K}}^\dagger a_{\vec{K}}$ and $b_{\vec{K}}^\dagger b_{\vec{K}}$ for various $\vec{K}$. The only change in Eqs. (2.11) – (2.14) is then found to be $E_P \rightarrow E_P - \mu$.

For the bilinears of the antiparticle ladder operators, relations similar to (2.11) – (2.14) hold with $E_P \rightarrow E_P + \mu$. Also, averages of mixed operators, like $\langle a_{\vec{P}} b_{\vec{P}'} \rangle$, are all zero.

We next turn to the case of the Majorana fermion field, which is analogous to the real scalar field, and has $\mu = 0$ necessarily. The important difference from the case of the real scalar field relevant for our purpose is that the number operator $c_{\vec{K},s}^\dagger c_{\vec{K},s}$ for any given mode can only have the eigenvalues 0 and 1. Consequently, the geometric series represented by the sum in Eq. (2.10) now contains only two terms and equals $1 + e^{-\beta E_P}$. So

$$\langle c_{\vec{P},s}^\dagger c_{\vec{P},s} \rangle = \frac{1}{e^{\beta E_P} + 1}, \quad (2.16)$$

which is the Fermi distribution function. Using $[c_{\vec{P},s}, c_{\vec{P},s}^\dagger]_+ = 1$, this leads to

$$\langle c_{\vec{P},s} c_{\vec{P},s}^\dagger \rangle = 1 - \frac{1}{e^{\beta E_P} + 1}. \quad (2.17)$$

The other thermal averages are zero, as for the real scalar field.

Finally, for a fermion field which is not self-conjugate,

$$J = \sum_{\vec{K},s} [(E_K - \mu) c_{\vec{K},s}^\dagger c_{\vec{K},s} + (E_K + \mu) d_{\vec{K},s}^\dagger d_{\vec{K},s}] \quad (2.18)$$

(analogous to Eq. (2.15)), so that the replacements $E_P \rightarrow E_P \mp \mu$ result, just as in the case of the complex scalar field.

2.3 Time-ordered thermal propagators

2.3.1 Scalar field

Time-ordered thermal propagators are defined by replacing the vacuum expectation value in the definitions of the vacuum propagators by a thermal average. Beginning with scalar fields, we shall first work out the more complicated case of the complex scalar field in detail, and then state the result for the case of the real scalar field. We shall use the Fourier expansion

$$\phi(x) = \int \frac{d^3 P}{\sqrt{(2\pi)^3 2E_P}} (a(\vec{P}) e^{-ip \cdot x} + b^\dagger(\vec{P}) e^{ip \cdot x}), \quad (2.19)$$

for the complex scalar field, where $p^\mu = (p^0, \vec{P})$ and $p^0 = E_P$. Since we are now considering continuous momenta, the thermal averages read

$$\langle a^\dagger(\vec{P}) a(\vec{P}') \rangle = \delta^{(3)}(\vec{P} - \vec{P}') f_B(E_P, \mu), \quad (2.20)$$

$$\langle a(\vec{P})a^\dagger(\vec{P}') \rangle = \delta^{(3)}(\vec{P} - \vec{P}') [1 + f_B(E_P, \mu)], \quad (2.21)$$

$$\langle b^\dagger(\vec{P})b(\vec{P}') \rangle = \delta^{(3)}(\vec{P} - \vec{P}') f_B(E_P, -\mu), \quad (2.22)$$

$$\langle b(\vec{P})b^\dagger(\vec{P}') \rangle = \delta^{(3)}(\vec{P} - \vec{P}') [1 + f_B(E_P, -\mu)], \quad (2.23)$$

with the other thermal averages equalling zero. Here the Bose distribution function for the particle is given by

$$f_B(E_P, \mu) = \frac{1}{e^{\beta(E_P - \mu)} - 1}. \quad (2.24)$$

Eqs. (2.20) and (2.21), consistent with Eq. (A.1), correspond to Eqs. (2.11)–(2.13).

The time-ordered thermal propagator for the complex scalar field is defined by

$$iD(x - y) \equiv \langle T[\phi(x)\phi^\dagger(y)] \rangle \quad (2.25)$$

$$= \theta(x_0 - y_0) \langle \phi(x)\phi^\dagger(y) \rangle + \theta(y_0 - x_0) \langle \phi^\dagger(y)\phi(x) \rangle. \quad (2.26)$$

Using Eqs. (2.19) – (2.23), we can obtain $\langle \phi(x)\phi^\dagger(y) \rangle$ and $\langle \phi^\dagger(y)\phi(x) \rangle$. After substituting these in Eq. (2.26), let us split up the resultant expression as

$$iD(x - y) = iD_F(x - y) + D_T(x - y), \quad (2.27)$$

where $iD_F(x - y)$ is the part independent of f_B , and $D_T(x - y)$ is the part arising out of the medium. Now, if f_B were set equal to zero in Eqs. (2.20) – (2.23), the thermal averages would reduce to the vacuum expectation values, and the thermal propagator to the propagator in the vacuum. So, we can immediately write down

$$\begin{aligned} D_F(x - y) &= \int \frac{d^4p}{(2\pi)^4} e^{-ip \cdot (x - y)} D_F(p), \\ D_F(p) &= \frac{1}{p^2 - m^2 + i\epsilon}. \end{aligned} \quad (2.28)$$

On the other hand,

$$D_T(x - y) = \int \frac{d^3P}{(2\pi)^3 2E_P} [f_B(E_P, \mu) e^{-ip \cdot (x - y)} + f_B(E_P, -\mu) e^{ip \cdot (x - y)}]. \quad (2.29)$$

(One can also get Eq. (2.29) by using Eq. (A.9) and the discretized versions of Eqs. (2.20) – (2.23), and finally using Eq. (A.4) in reverse.) Now, Eq. (2.29) can be rewritten as

$$\begin{aligned} D_T(x - y) = \int \frac{d^4p}{(2\pi)^3 2E_P} & [\delta(p_0 - E_P) f_B(p_0, \mu) e^{-ip_0(x_0 - y_0) + i\vec{P} \cdot (\vec{x} - \vec{y})} \\ & + \delta(p_0 + E_P) f_B(-p_0, -\mu) e^{-ip_0(x_0 - y_0) - i\vec{P} \cdot (\vec{x} - \vec{y})}]. \end{aligned} \quad (2.30)$$

Changing $\vec{P}$ to $-\vec{P}$ in the second term, and using

$$\theta(\pm p_0)\delta(p^2 - m^2) = \frac{1}{2E_P}\delta(p_0 \mp E_P), \quad (2.31)$$

we obtain

$$\begin{aligned} D_T(x-y) &= \int \frac{d^4p}{(2\pi)^4} e^{-ip \cdot (x-y)} D_T(p), \\ D_T(p) &= 2\pi\delta(p^2 - m^2)\eta_B(p), \\ \eta_B(p) &= \theta(p_0)f_B(p_0, \mu) + \theta(-p_0)f_B(-p_0, -\mu). \end{aligned} \quad (2.32)$$

Thus, the thermal propagator is given by Eqs. (2.27), (2.28) and (2.32). However, this form is not manifestly covariant since the distribution function (2.24) was written in the rest frame of the medium. Therefore we introduce the four-velocity v^μ of the centre-of-mass of the medium. In the rest frame of the medium,

$$v^\mu = (1, \vec{0}). \quad (2.33)$$

Then carrying out the replacement

$$p_0 \rightarrow p \cdot v \quad (2.34)$$

in Eq. (2.32), we arrive at the manifestly covariant form of the thermal propagator.

For the real scalar field, we have to use the Fourier expansion

$$\phi(x) = \int \frac{d^3P}{\sqrt{(2\pi)^3 2E_P}} \left(a(\vec{P})e^{-ip \cdot x} + a^\dagger(\vec{P})e^{ip \cdot x} \right), \quad (2.35)$$

and the thermal averages (2.20) and (2.21) with $\mu = 0$. This leads to the thermal propagator

$$\langle T[\phi(x)\phi(y)] \rangle = \int \frac{d^4p}{(2\pi)^4} e^{-ip \cdot (x-y)} \left[\frac{i}{p^2 - m^2 + i\epsilon} + 2\pi\delta(p^2 - m^2) \frac{1}{e^{\beta|p \cdot v|} - 1} \right], \quad (2.36)$$

which is seen to be the thermal propagator of the complex scalar field at $\mu = 0$.

2.3.2 Fermion field

For the fermion field, we shall use the Fourier expansion

$$\psi(x) = \int \frac{d^3P}{(2\pi)^{3/2}} \sum_s \left(c_s(\vec{P})u_s(\vec{P})e^{-ip \cdot x} + d_s^\dagger(\vec{P})v_s(\vec{P})e^{ip \cdot x} \right) \quad (2.37)$$

and normalize the spinors as

$$u_s^\dagger(\vec{P})u_{s'}(\vec{P}) = \delta_{ss'}, \quad v_s^\dagger(\vec{P})v_{s'}(\vec{P}) = \delta_{ss'}. \quad (2.38)$$

(This normalization will be later extended to the spinors in a medium; see Eqs. (4.27) and (4.46).) The thermal averages then read

$$\langle c_s^\dagger(\vec{P})c_{s'}(\vec{P}') \rangle = \delta_{ss'}\delta^{(3)}(\vec{P} - \vec{P}')f_f(E_P), \quad (2.39)$$

$$\langle c_s(\vec{P})c_{s'}^\dagger(\vec{P}') \rangle = \delta_{ss'}\delta^{(3)}(\vec{P} - \vec{P}')[1 - f_f(E_P)], \quad (2.40)$$

$$\langle d_s^\dagger(\vec{P})d_{s'}(\vec{P}') \rangle = \delta_{ss'}\delta^{(3)}(\vec{P} - \vec{P}')f_{\bar{f}}(E_P), \quad (2.41)$$

$$\langle d_s(\vec{P})d_{s'}^\dagger(\vec{P}') \rangle = \delta_{ss'}\delta^{(3)}(\vec{P} - \vec{P}')[1 - f_{\bar{f}}(E_P)], \quad (2.42)$$

with the other thermal averages equalling zero. Here the distribution functions for a fermion and an antifermion are given by

$$f_{f,\bar{f}}(E_P) = \frac{1}{e^{\beta(E_P \mp \mu_f)} + 1}, \quad (2.43)$$

respectively. (We have denoted the fermion chemical potential by μ_f , since we will later deal with several species of the fermion f .) Eqs. (2.39) and (2.40), consistent with Eq. (A.10), correspond to Eqs. (2.16) and (2.17).

The thermal propagator for the fermion is defined by

$$iS_{fab}(x) \equiv \langle T[\psi_a(x)\bar{\psi}_b(y)] \rangle \quad (2.44)$$

$$= \theta(x_0 - y_0)\langle \psi_a(x)\bar{\psi}_b(y) \rangle - \theta(y_0 - x_0)\langle \bar{\psi}_b(y)\psi_a(x) \rangle. \quad (2.45)$$

Using Eq. (2.37), Eqs. (2.39) – (2.42), and the spin-sum relations

$$\begin{aligned} \sum_s u_s(\vec{P})\bar{u}_s(\vec{P}) &= \frac{E_P\gamma_0 - \vec{P} \cdot \vec{\gamma} + m_f}{2E_P}, \\ \sum_s v_s(\vec{P})\bar{v}_s(\vec{P}) &= \frac{E_P\gamma_0 - \vec{P} \cdot \vec{\gamma} - m_f}{2E_P} \end{aligned} \quad (2.46)$$

following from Eq. (2.38), we can find the R.H.S. of Eq. (2.45). The expression is then split up into the vacuum part and the thermal part:

$$S_f(x - y) = \int \frac{d^4p}{(2\pi)^4} e^{-ip \cdot (x - y)} S_f(p), \quad (2.47)$$

$$iS_f(p) = iS_{Ff}(p) + S_{Tf}(p). \quad (2.48)$$

The vacuum part is

$$S_{Ff} = \frac{\not{p} + m_f}{p^2 - m_f^2 + i\epsilon}, \quad (2.49)$$

while, following the derivation in the case of the complex scalar field, the thermal part is found to be

$$S_{Tf}(p) = -2\pi(\not{p} + m_f)\delta(p^2 - m_f^2)\eta_f(p), \quad (2.50)$$

$$\eta_f(p) = \frac{\theta(p \cdot v)}{e^{\beta(p \cdot v - \mu_f)} + 1} + \frac{\theta(-p \cdot v)}{e^{-\beta(p \cdot v - \mu_f)} + 1}. \quad (2.51)$$

2.3.3 Photon field

In this case, the thermal propagator in the Feynman gauge is the same as that for the real scalar field, except for the facts that the photon is massless, and that there is an extra factor of $-\eta_{\mu\nu}$ (as in the vacuum). Thus, we have the propagator

$$iD_{\mu\nu}(k) = -\eta_{\mu\nu} [i\Delta_F(k) + \Delta_T(k)], \quad (2.52)$$

where

$$\Delta_F(k) = \frac{1}{k^2 + i\epsilon}, \quad (2.53)$$

$$\Delta_T(k) = 2\pi\delta(k^2)\eta_\gamma(k), \quad (2.54)$$

$$\eta_\gamma(k) = \frac{1}{e^{\beta|k \cdot v|} - 1}. \quad (2.55)$$

The factor $\eta_\gamma(k)$ arises out of the photon distribution function

$$f_\gamma(K) = \frac{1}{e^{\beta K} - 1}, \quad (2.56)$$

where K is the magnitude of the three-momentum of the photon.

We end this section with the following observation. At any finite temperature, the distribution function contained in the medium-dependent part of the thermal propagator becomes an exponential damping factor for momenta much larger than T . Consequently, the corresponding loop integration will have an effective cut-off of $O(T)$. If $T = 0$, we can still have a Fermi gas; in that case, the Fermi momentum provides a cut-off. Thus, the medium does not introduce any new ultraviolet divergence, and renormalization in the vacuum suffices to make the theory finite in the presence of thermal background.

2.4 Additional vertex and propagators

The full structure of the real-time formulation of thermal field theory is more complicated than the replacement of vacuum propagators with time-ordered thermal propagators. However, as will be seen later, the additional features of the full structure do not play a role in the present work. So, in this section, we shall just outline the main results involving those additional features [32].

Ultimately, one has to calculate an n -point Green's function like $\langle T[\phi(x_1)\phi^\dagger(x_2)\cdots\phi(x_n)] \rangle$, where $\phi(x_i)$ are the Heisenberg picture fields. (We shall consider only complex scalar fields in this section. Similar results hold for other fields.) In a medium, as in the vacuum, one can prove Wick's theorem and reduce the n -point function to a sum of fully contracted terms. But the fact that the Green's function in the present case involves a thermal average over all the Fock space states, rather than a vacuum expectation value, leads to four types of contractions or “propagators.” They are defined as

$$iD_{11}(x-y) = \langle T[\phi(x)\phi^\dagger(y)] \rangle, \quad (2.57)$$

$$iD_{22}(x-y) = \langle \bar{T}[\phi(x)\phi^\dagger(y)] \rangle, \quad (2.58)$$

$$iD_{12}(x-y) = \langle \phi^\dagger(y)\phi(x) \rangle, \quad (2.59)$$

$$iD_{21}(x-y) = \langle \phi(x)\phi^\dagger(y) \rangle. \quad (2.60)$$

Here $\bar{T}$ stands for anti-time ordering. In Eqs. (2.57) – (2.60), the fields are the interaction picture fields, which can be expressed in terms of the ladder operators. Thus, in addition to the time-ordered propagator of Sec. 2.3, we also have anti-time ordered propagator, as well as propagators without any ordering in time. As given by Eqs. (2.27), (2.28) and (2.32)

$$iD_{11}(p) = \frac{i}{p^2 - m^2 + i\epsilon} + 2\pi\delta(p^2 - m^2)\eta_B(p). \quad (2.61)$$

The others are found to be

$$iD_{22}(p) = -\frac{i}{p^2 - m^2 - i\epsilon} + 2\pi\delta(p^2 - m^2)\eta_B(p), \quad (2.62)$$

$$iD_{12}(p) = 2\pi\delta(p^2 - m^2)[\eta_B(p) + \theta(-p \cdot v)], \quad (2.63)$$

$$iD_{21}(p) = 2\pi\delta(p^2 - m^2)[\eta_B(p) + \theta(p \cdot v)]. \quad (2.64)$$

Here it is to be understood that the replacement (2.34) has been carried out in $\eta_B(p)$.

The Feynman diagrams now contain two types of vertices: the 1-type vertex is the usual one, while the 2-type vertex has a vertex factor which differs from that for the 1-type vertex by a sign. Green's functions can now be evaluated with these two types of vertices and four types of propagators. However, the external legs representing physical particles can only be connected to 1-type vertices.

The importance of this doubling of degrees of freedom manifests itself in diagrammatic calculations in the following way. If one uses only the 1-type vertex and “11” propagator, one sometimes ends up with singular expressions involving products of delta functions with the same argument. However the use of the additional vertex and propagators leads to the cancellation of such singularities [33].

CHAPTER 3

Linearized theory of gravity

3.1 Introduction

Gravity is governed by the general theory of relativity which tells us how the geometry of space-time, as given by the metric tensor $g_{\mu\nu}(x)$, is affected by the presence of matter. The Einstein field equations for $g_{\mu\nu}$ in the absence of external source, derived from the action for pure gravity, are

$$R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} = 0. \quad (3.1)$$

Here $R_{\mu\nu}$ is the Ricci tensor and R is the curvature scalar. One important point is that these equations are non-linear in $g_{\mu\nu}$. This is in contrast with the linear free-field equations for the scalar, fermion and photon fields, where we use the principle of linear superposition to write down the general solution before quantizing the field. The reason for this non-linearity in the case of gravity is that the gravitational field possesses energy and hence effective mass, and therefore *produces* gravitational field. The gravitational field thus contributes to its own source. To ignore this feedback effect, we have to consider a weak gravitational field. One thus considers only small deviations from the flat space metric, and writes

$$g_{\mu\nu}(x) = \eta_{\mu\nu} + 2\kappa h_{\mu\nu}(x) \quad (3.2)$$

with $|\kappa h_{\mu\nu}| \ll 1$. Here κ is related to the Newton constant G . The approximation stated in Eq. (3.2) is the starting point of the linearized theory of gravity [34, 35]. Now $\eta_{\mu\nu}$, and hence from Eq. (3.2), $h_{\mu\nu}$, are symmetric flat space tensors, but not coordinate tensors (in contrast with $g_{\mu\nu}$, which is a symmetric coordinate tensor, and so, as a special case, also a tensor in flat space). Consequently, *all* tensors in an expression should be regarded only

as flat space tensors *after* we use the approximation of Eq. (3.2) to perform an expansion of the expression in powers of κ , and we will adopt the convention of raising and lowering the indices of such tensors with the flat space metric.

The precise value of κ , to be deduced in the next section by identifying $h_{\mu\nu}$ with the graviton field, is given by

$$\kappa = \sqrt{8\pi G}. \quad (3.3)$$

Since $\sqrt{G} = 1/m_{\text{Planck}}$ in natural units, it is clear that the weak field approximation given by Eq. (3.2) is valid when all mass-scales in the problem are much smaller than the Planck mass. These mass-scales include not only particle masses, but also temperature and chemical potential, as we shall later perform calculations in the presence of a thermal background. Since m_{Planck} is $O(10^{19})$ GeV, the weak field approximation holds in all physical situations except in the very early universe.

At this point, it is interesting to consider the case of the Yang-Mills theory for comparison, because the equation of motion for the non-Abelian gauge field derived from the pure Yang-Mills Lagrangian is also non-linear. However, there the equation of motion *is* linear when one considers the free gauge fields, setting the coupling constant equal to zero, and so the need to linearize does not arise. In contrast, the flat space metric tensor $\eta_{\mu\nu}$, being a constant, does not have any dynamics (and the LHS of Eq. (3.1) reduces to zero in this case). So, to do quantum field theory with gravity it is essential to impose the condition (3.2), which considers linear deviations from the flat space metric case.

3.2 Graviton

The field equations (3.1) follow from the Einstein-Hilbert action given by

$$\mathcal{A} = \int d^4x \mathcal{L}, \quad (3.4)$$

$$\mathcal{L} = \frac{1}{16\pi G} \sqrt{-g} R. \quad (3.5)$$

The curvature scalar, the Ricci tensor and the affine connection are respectively given by

$$R = g^{\mu\nu} R_{\mu\nu}, \quad (3.6)$$

$$R_{\mu\nu} = \partial_\nu \Gamma_{\mu\lambda}^\lambda - \partial_\lambda \Gamma_{\mu\nu}^\lambda + \Gamma_{\mu\lambda}^\eta \Gamma_{\nu\eta}^\lambda - \Gamma_{\mu\nu}^\eta \Gamma_{\lambda\eta}^\lambda, \quad (3.7)$$

$$\Gamma_{\mu\nu}^\lambda = \frac{1}{2} g^{\lambda\sigma} (\partial_\mu g_{\nu\sigma} + \partial_\nu g_{\mu\sigma} - \partial_\sigma g_{\mu\nu}), \quad (3.8)$$

while

$$g \equiv \det g_{\mu\nu}. \quad (3.9)$$

In anticipation of the transition to quantum field theory, here and elsewhere we always write the action in the form (3.4) (although $\mathcal{L}$ is not a coordinate scalar).

From the weak field condition (3.2), we obtain

$$g^{\mu\nu} = \eta^{\mu\nu} - 2\kappa h^{\mu\nu} + O(\kappa^2) \quad (3.10)$$

(since $g^{\mu\nu}g_{\nu\lambda} = \delta_\lambda^\mu$), and

$$\sqrt{-g} = 1 + \kappa h + O(\kappa^2) \quad (3.11)$$

where

$$h \equiv h^\mu{}_\mu = \eta^{\mu\nu} h_{\mu\nu}. \quad (3.12)$$

Note that we are raising and lowering the indices of the graviton field with the flat space metric [36]; this is in accord with the convention stated in Sec. 3.1.

Use of Eqs. (3.2) and (3.10) in Eq. (3.8) leads to

$$\Gamma_{\mu\nu}^\lambda = \kappa(\eta^{\lambda\sigma} - 2\kappa h^{\lambda\sigma})(\partial_\mu h_{\nu\sigma} + \partial_\nu h_{\mu\sigma} - \partial_\sigma h_{\nu\mu}) + O(\kappa^3). \quad (3.13)$$

Putting this in Eq. (3.7), one can determine $R_{\mu\nu}$ to $O(\kappa^2)$. Let us write

$$R_{\mu\nu} = R_{\mu\nu}^{(1)} + R_{\mu\nu}^{(2)} + O(\kappa^3) \quad (3.14)$$

with $R_{\mu\nu}^{(1)}$ and $R_{\mu\nu}^{(2)}$ denoting the parts linear and quadratic in κ respectively.

Following again the convention stated in Sec. 3.1, we shall now raise and lower the indices of $R_{\mu\nu}^{(1)}$, $R_{\mu\nu}^{(2)}$ and $\partial/\partial x^\mu$ with the flat space metric [36]. Then

$$\begin{aligned} 16\pi G \mathcal{L} &= \sqrt{-g} g^{\mu\nu} R_{\mu\nu} \\ &= R^\mu{}_\mu^{(1)} + \kappa h R^\mu{}_\mu^{(1)} - 2\kappa h^{\mu\nu} R_{\mu\nu}^{(1)} + R^\mu{}_\mu^{(2)} + O(\kappa^3). \end{aligned} \quad (3.15)$$

The first term on the RHS is the only term of $O(\kappa)$. In fact,

$$R^\mu{}_\mu^{(1)} = 2\kappa \partial_\mu (\partial^\mu h - \partial_\lambda h^{\lambda\mu}), \quad (3.16)$$

and this, being a surface term, does not contribute to the Einstein-Hilbert action. So we now turn to the three $O(\kappa^2)$ terms on the RHS of Eq. (3.15). First consider $R^\mu{}_\mu{}^{(2)}$. After leaving out surface terms, we obtain

$$R^\mu{}_\mu{}^{(2)} = \kappa^2[(\partial_\mu h_{\eta\lambda} + \partial_\lambda h_{\mu\eta} - \partial_\eta h_{\lambda\mu})(\partial^\mu h^{\eta\lambda} - \partial^\lambda h^{\mu\eta} + \partial^\eta h^{\lambda\mu}) + (\partial_\mu h)(\partial^\mu h - 2\partial_\lambda h^{\lambda\mu})]. \quad (3.17)$$

Carrying out of several partial integrations with discarding of surface terms [for example, $(\partial_\lambda h_{\mu\eta})(\partial^\eta h^{\lambda\mu}) \rightarrow -h_{\mu\eta}(\partial_\lambda \partial^\eta h^{\lambda\mu}) \rightarrow (\partial^\eta h_{\mu\eta})(\partial_\lambda h^{\lambda\mu})$] leads to

$$R^\mu{}_\mu{}^{(2)} = -\kappa^2[(\partial_\mu h_{\eta\lambda})(\partial^\mu h^{\eta\lambda}) - 2(\partial_\lambda h^{\mu\lambda})(\partial^\eta h_{\mu\eta}) + 2(\partial_\mu h)(\partial_\lambda h^{\mu\lambda}) - (\partial_\mu h)(\partial^\mu h)]. \quad (3.18)$$

Similarly one can write $\kappa h R^\mu{}_\mu{}^{(1)}$ and $2\kappa h^{\mu\nu} R_{\mu\nu}{}^{(1)}$ in terms of the combinations contained on the R.H.S. of Eq. (3.18). Putting everything back into Eq. (3.15), we get

$$16\pi G\mathcal{L} = \kappa^2[(\partial_\mu h_{\eta\lambda})(\partial^\mu h^{\eta\lambda}) - 2(\partial_\lambda h^{\mu\lambda})(\partial^\eta h_{\mu\eta}) + 2(\partial_\mu h)(\partial_\lambda h^{\mu\lambda}) - (\partial_\mu h)(\partial^\mu h)]. \quad (3.19)$$

Here and henceforth it is to be understood that the expression for $16\pi G\mathcal{L}$ has been written neglecting the $O(\kappa^3)$ terms.

With the aim of arriving at the propagator, we carry out further partial integrations to obtain

$$16\pi G\mathcal{L} = \kappa^2 h_{\alpha\beta} \mathcal{P}^{\alpha\beta\mu\nu} h_{\mu\nu}, \quad (3.20)$$

$$\begin{aligned} \mathcal{P}^{\alpha\beta\mu\nu} = & -\frac{1}{2}[(\eta^{\alpha\mu}\eta^{\beta\nu} + \eta^{\alpha\nu}\eta^{\beta\mu} - 2\eta^{\alpha\beta}\eta^{\mu\nu})\square - (\eta^{\beta\nu}\partial^\alpha\partial^\mu + \eta^{\alpha\nu}\partial^\beta\partial^\mu + \eta^{\beta\mu}\partial^\alpha\partial^\nu \\ & + \eta^{\alpha\mu}\partial^\beta\partial^\nu) + 4\eta^{\alpha\beta}\partial^\mu\partial^\nu]. \end{aligned} \quad (3.21)$$

We have explicitly symmetrized $\mathcal{P}^{\alpha\beta\mu\nu}$ with respect to $\alpha \leftrightarrow \beta$ and with respect to $\mu \leftrightarrow \nu$ (such symmetrization helps one to guess the form of the inverse; see later). But this operator does not possess an inverse! This is easily seen from the fact that

$$\mathcal{P}^{\alpha\beta\mu\nu} \partial_\alpha \partial_\beta \partial_\mu \partial_\nu \Lambda(x) = 0 \quad (3.22)$$

where $\Lambda(x)$ is any function of x , i.e., the operator has a zero eigenvalue.

The problem is similar to the one encountered in quantizing the photon field, and the solution also is the same, viz., gauge-fixing. To this end, one chooses the harmonic or de Donder gauge

$$F^\mu \equiv \partial_\nu h^{\mu\nu} - \frac{1}{2}\partial^\mu h = 0 \quad (3.23)$$

and adds to $16\pi G\mathcal{L}$ the gauge-fixing term [37]

$$16\pi G\mathcal{L}_{\text{GF}} = 2\kappa^2 F^\mu F_\mu. \quad (3.24)$$

Before proceeding further, let us dwell a little on gauge invariance.

While the Einstein-Hilbert action is invariant under general coordinate transformations, the allowed coordinate transformations in the present case are only those which sustain the weak field approximation. Thus, we consider

$$x^\mu \rightarrow x'^\mu = x^\mu + 2\kappa\epsilon^\mu(x) \quad (3.25)$$

and demand that

$$g'^{\mu\nu} = \eta^{\mu\nu} - 2\kappa h'^{\mu\nu} + O(\kappa^2). \quad (3.26)$$

Using these relations in

$$g'^{\mu\nu} = \frac{\partial x'^\mu}{\partial x^\lambda} \frac{\partial x'^\nu}{\partial x^\rho} g^{\lambda\rho}, \quad (3.27)$$

we arrive at the following gauge transformation of $h_{\mu\nu}$ which keeps the action invariant:

$$h'_{\mu\nu} = h_{\mu\nu} - (\partial_\mu \epsilon_\nu + \partial_\nu \epsilon_\mu), \quad (3.28)$$

where $\epsilon_\mu \equiv \eta_{\mu\nu}\epsilon^\nu$. It can easily be checked that if $h_{\mu\nu}$ does not satisfy the gauge condition stated in Eq. (3.23), $h'_{\mu\nu}$ does, provided

$$\square\epsilon^\mu = F^\mu. \quad (3.29)$$

So it is always possible to implement the harmonic gauge condition. Another point to be confirmed is that the gauge-fixing term given in Eq. (3.24) does break the gauge invariance of the action. This readily follows from

$$g^{\mu\nu}\Gamma_{\mu\nu}^\lambda = -4\kappa F^\lambda + O(\kappa^2). \quad (3.30)$$

Since the affine connection is not a coordinate tensor, the square of the L.H.S. is not a coordinate scalar. (Note that $\mathcal{A}_{\text{GF}}$ involves d^4x which also is not a scalar. But the transformation of d^4x cannot compensate the breaking of gauge invariance, as $16\pi G\mathcal{L}_{\text{GF}}$ is already of $O(\kappa^2)$.)

We now explore the consequences of fixing the gauge. Addition of the gauge non-invariant term given in Eq. (3.24) immediately yields the correctly normalized kinetic terms for the massless spin-2 field, to be identified as the graviton field:

$$\mathcal{L} + \mathcal{L}_{\text{GF}} = \frac{1}{2}(\partial_\mu h_{\nu\lambda})(\partial^\mu h^{\nu\lambda}) - \frac{1}{4}(\partial_\mu h)(\partial^\mu h), \quad (3.31)$$

provided we set κ equal to the value stated in Eq. (3.3).

We conclude this section by obtaining the graviton propagator. Partial integration of the last expression leads to

$$\mathcal{A} + \mathcal{A}_{\text{GF}} = \int d^4x \frac{1}{2} h_{\alpha\beta} \mathcal{Q}^{\alpha\beta\mu\nu} h_{\mu\nu}, \quad (3.32)$$

$$\mathcal{Q}^{\alpha\beta\mu\nu} = -\frac{1}{2}(\eta^{\alpha\mu}\eta^{\beta\nu} + \eta^{\alpha\nu}\eta^{\beta\mu} - \eta^{\alpha\beta}\eta^{\mu\nu})\Box. \quad (3.33)$$

So the graviton propagator $iD_{\alpha\beta,\mu\nu}$ is i/k^2 times the inverse of

$$M^{\alpha\beta,\mu\nu} \equiv \frac{1}{2}(\eta^{\alpha\mu}\eta^{\beta\nu} + \eta^{\alpha\nu}\eta^{\beta\mu} - \eta^{\alpha\beta}\eta^{\mu\nu}). \quad (3.34)$$

The inverse must be determined from

$$M^{\alpha\beta,\mu\nu}(M^{-1})_{\alpha\beta,\lambda\rho} = \frac{1}{2}(\delta_\lambda^\mu\delta_\rho^\nu + \delta_\rho^\mu\delta_\lambda^\nu). \quad (3.35)$$

Guided by the fact that M^{-1} must be symmetric under $\alpha \leftrightarrow \beta$ and (separately) under $\lambda \leftrightarrow \rho$, one can write down

$$(M^{-1})_{\alpha\beta,\lambda\rho} = A(\eta_{\alpha\lambda}\eta_{\beta\rho} + \eta_{\alpha\rho}\eta_{\beta\lambda}) + B\eta_{\alpha\beta}\eta_{\lambda\rho}. \quad (3.36)$$

We then obtain $A = 1/2 = -B$. Therefore the graviton propagator is given by

$$iD_{\alpha\beta,\lambda\rho} = i\frac{\eta_{\alpha\lambda}\eta_{\beta\rho} + \eta_{\alpha\rho}\eta_{\beta\lambda} - \eta_{\alpha\beta}\eta_{\lambda\rho}}{2k^2}. \quad (3.37)$$

One should now compare the photon propagator in the Feynman gauge, given by $iD_{\mu\nu} = -i\eta_{\mu\nu}/k^2$. Clearly D_{00} is negative, while $D_{00,00}$ is positive. This has the consequence that the fermion-fermion interaction in the non-relativistic limit is repulsive when mediated by the photon and attractive when mediated by the graviton [38].

3.3 Photons in curved space

For the electromagnetic field, the Lagrangian in curved space is given by

$$\mathcal{L}_G^{(\gamma)} = -\frac{1}{4}\sqrt{-g}F_{\mu\nu}F_{\alpha\beta}g^{\mu\alpha}g^{\nu\beta} \quad (3.38)$$

where

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu. \quad (3.39)$$

This has been written down following the standard prescription of passing from flat space to curved space, i.e., by replacing all flat space tensors with coordinate tensors, all ordinary derivatives with covariant derivatives (by which we shall mean gravity-covariant derivatives) and the flat space metric tensor with $g_{\mu\nu}$. The additional $\sqrt{-g}$ in Eq. (3.38) comes because d^4x in the action gets replaced with the coordinate scalar $d^4x\sqrt{-g}$. It may be noted that the use of covariant derivative and ordinary derivative in Eq. (3.39) are equivalent due to the symmetry of the Christoffel symbol in the two lower indices.

3.4 Spinors in curved space

The procedure of replacing flat space tensors with coordinate tensors cannot be extended to the case of spinors. The procedure works with tensors because the tensor representations of $GL(4, \mathbb{R})$, the group of 4×4 real matrices, behave like tensors under the subgroup $SO(3,1)$. Thus, considering the vector representation as an example, $V'^\mu(x') = (\partial x'^\mu / \partial x^\nu) V^\nu(x)$ becomes $V'^\mu(x') = \Lambda^\mu{}_\nu V^\nu(x)$ if we restrict ourselves to $x'^\mu = \Lambda^\mu{}_\nu x^\nu$. But there are no representations of $GL(4, \mathbb{R})$ which behave like spinors under $SO(3,1)$ (there is no function of x and x' which reduces to $D(\Lambda)$ for $x'^\mu = \Lambda^\mu{}_\nu x^\nu$).

Consequently, spinors have to be considered as scalars under general coordinate transformations. But the problem of incorporating the flat space spinorial property into curved space remains. This is tackled by making use of the tangent space formalism, to be described now [39, 40, 41].

At any given point of the curved space, we can choose a locally inertial coordinate system. Let the locally inertial coordinates at the point x^μ be $\xi^a(x)$ with $a = 0, 1, 2, 3$. Since the locally inertial coordinate system at each point may be redefined by an arbitrary Lorentz transformation (LT), the action must be invariant under the local LT

$$\xi^a(x) \rightarrow \xi'^a(x) = \Lambda^a{}_b(x) \xi^b(x), \quad (3.40)$$

with $\Lambda^a{}_b(x)$ a real matrix satisfying

$$\eta_{ab} \Lambda^a{}_c(x) \Lambda^b{}_d(x) = \eta_{cd}. \quad (3.41)$$

The action, of course, has to be also invariant under the general coordinate transformation $x^\mu \rightarrow x'^\mu$, which is independent of the local LT. The requirement of invariance under local LT does not affect the actions discussed earlier, since coordinate tensors like $g_{\mu\nu}$ and A_μ are taken to be local Lorentz scalars (i.e. scalars under local LT).

Let us now consider the vierbein or tetrad, defined by

$$e^a{}_\mu(X) \equiv \left. \frac{\partial \xi^a(x)}{\partial x^\mu} \right|_{x=X}. \quad (3.42)$$

These constitute a set of four coordinate vectors, forming a basis for the (flat) tangent space to the curved space at the point $x = X$. Under $x^\mu \rightarrow x'^\mu$, they transform as

$$e'^a{}_\mu(x') = \frac{\partial x^\nu}{\partial x'^\mu} e^a{}_\nu(x), \quad (3.43)$$

since $\xi'^a(x') = \xi^a(x)$. Under local LT, the vierbein transforms just like $\xi^a(x)$:

$$e'^a{}_\mu(x) = \Lambda^a{}_b(x) e^b{}_\nu(x), \quad (3.44)$$

since x^μ does not transform. From the definition given by Eq. (3.42), it follows that raising or lowering of the local Lorentz index a (in general, a latin index) and the general coordinate index μ (in general, a greek index) of the vierbein must be done with η_{ab} and $g_{\mu\nu}$ respectively.

The metric $g_{\mu\nu}$ is given by

$$g_{\mu\nu}(x) = \eta_{ab} e^a{}_\mu(x) e^b{}_\nu(x). \quad (3.45)$$

Therefore $\delta^\mu_\nu = e_b{}^\mu e^b{}_\nu$, i.e., $e_b{}^\mu$ is the inverse of $e^b{}_\mu$. We can rewrite this as $\mathbf{1}_{\mu\nu} = (\mathbf{E}^{-1})_{\mu b} \mathbf{E}_{b\nu}$, where the indices now denote matrix indices. It then also follows that $\mathbf{1}_{ab} = \mathbf{E}_{a\mu} (\mathbf{E}^{-1})_{\mu b}$. This means that $\delta^a_b = e^a{}_\mu e_b{}^\mu$, or,

$$\eta^{ab} = g^{\mu\nu}(x) e^a{}_\mu(x) e^b{}_\nu(x). \quad (3.46)$$

Eq. (3.45), or equivalently, Eq. (3.46), expresses the orthonormality of the vierbeins.

The vierbeins turn out to be necessary for constructing the curved space Lagrangian for the Dirac fermion. This will involve modifying the flat space Dirac Lagrangian

$$\mathcal{L}_0^{(f)} = \left[\frac{i}{2} \bar{\psi} \gamma^\mu \partial_\mu \psi + \text{h.c.} \right] - m_f \bar{\psi} \psi \quad (3.47)$$

such that the action is both a coordinate scalar and a local Lorentz scalar. With this aim in mind, we turn to the transformation properties of the spinor field.

Spinor fields are coordinate scalars which transform under local LT as

$$\psi_\alpha(x) \rightarrow \psi'_\alpha(x) = D_{\alpha\beta}(\Lambda(x))\psi_\beta(x). \quad (3.48)$$

Here $D_{\alpha\beta}(\Lambda)$ is the spinor representation of the Lorentz group. Note that we use α and β to denote the components of ψ , $D(\Lambda)$ and other quantities associated with the spinor representation; they must not be confused with general coordinate indices. Eq. (3.48) is the relation which incorporates the flat space spinorial property into general relativity.

Since Λ is a function of x^μ , $\partial_\mu\psi_\alpha$ does not transform like ψ_α under local LT. Hence one defines the covariant derivative

$$\mathcal{D}_\mu\psi_\alpha \equiv \partial_\mu\psi_\alpha + [\Omega_\mu]_{\alpha\beta}\psi_\beta \quad (3.49)$$

with the connection matrix Ω_μ transforming as

$$\Omega'_\mu = D(\Lambda)\Omega_\mu D^{-1}(\Lambda) - (\partial_\mu D(\Lambda))D^{-1}(\Lambda), \quad (3.50)$$

so that

$$\mathcal{D}_\mu\psi_\alpha(x) \rightarrow D_{\alpha\beta}(\Lambda(x))\mathcal{D}_\mu\psi_\beta(x). \quad (3.51)$$

The connection matrix can be written as

$$[\Omega_\mu]_{\alpha\beta}(x) = \frac{i}{2}[S_{ab}]_{\alpha\beta}\omega_\mu^{ab}(x) \quad (3.52)$$

where S_{ab} are the generators of the Lorentz group in the spinor representation:

$$S_{ab} = \frac{1}{2}\sigma_{ab} = \frac{i}{4}[\gamma_a, \gamma_b] \quad (3.53)$$

(with γ_a denoting the ordinary gamma matrices), and ω_μ^{ab} is a coordinate vector field antisymmetric in a and b , called the spin connection.

From Eq. (3.51), it follows that $\bar{\psi}\gamma^a\mathcal{D}_\mu\psi$ is a local Lorentz vector (just like $\bar{\psi}\gamma^a\psi$). It is also a coordinate vector (as ψ is a coordinate scalar). So when we contract it with e_a^μ , we get a quantity which is a local Lorentz scalar and a coordinate scalar. We can now modify Eq. (3.47) to write down the curved space Lagrangian

$$\mathcal{L}_G^{(f)} = \sqrt{-g}\left[\left(\frac{i}{2}\bar{\psi}\gamma^a e_a^\mu \mathcal{D}_\mu\psi + \text{h.c.}\right) - m_f\bar{\psi}\psi\right] \quad (3.54)$$

$$= \sqrt{-g}\left[\left(\frac{i}{2}\bar{\psi}\gamma^a e_a^\mu \left(\partial_\mu + \frac{i}{4}\sigma^{bc}\omega_{\mu bc}\right)\psi + \text{h.c.}\right) - m_f\bar{\psi}\psi\right] \quad (3.55)$$

The latin indices of the spin connection are by definition raised and lowered with the flat space metric.

We now explain why we started with the explicitly hermitian form $\mathcal{L}_0^{(f)}$ of the flat space Dirac Lagrangian. As long as we stay in flat space, the form $\mathcal{L}_0^{(f)'} = \bar{\psi}(i\gamma^\mu\partial_\mu - m_f)\psi$ can also be used, since it differs from $\mathcal{L}_0^{(f)}$ by $\frac{i}{2}\partial_\mu(\bar{\psi}\gamma^\mu\psi)$ which contributes a surface term to the action. But this difference is modified to $\frac{i}{2}\sqrt{-g}e_a^\mu\partial_\mu(\bar{\psi}\gamma^a\psi)$ in curved space, which is no longer a surface term. Thus we would have missed some contribution if we naively modified $\mathcal{L}_0^{(f)}'$ to make the transition to curved space.

Our next aim is to express the spin connections in terms of the vierbeins. The procedure to be adopted by us will involve the covariant derivative of the vierbein. So we first discuss the covariant derivative of a local Lorentz vector.

We have already discussed the covariant derivative of a local Lorentz spinor, and similar considerations apply for other representations of the Lorentz group. The analogues of Eqs. (3.48)-(3.52) for a local Lorentz vector field V^a are

$$V^a(x) \rightarrow V'^a(x) = \Lambda^a_b(x)V^b(x), \quad (3.56)$$

$$\mathcal{D}_\mu V^a \equiv \partial_\mu V^a + [\Omega_\mu^V]^a_b V^b, \quad (3.57)$$

$$[\Omega_\mu^V]^a_b = \Lambda^a_c [\Omega_\mu^V]^c_d (\Lambda^{-1})^d_b - (\partial_\mu \Lambda^a_c) (\Lambda^{-1})^c_b \quad (3.58)$$

$$\mathcal{D}_\mu V^a(x) \rightarrow \Lambda^a_b(x) \mathcal{D}_\mu V^b(x), \quad (3.59)$$

$$[\Omega_\mu^V]^a_b(x) = \frac{i}{2} [J_{cd}]^a_b \omega_\mu^{cd}(x). \quad (3.60)$$

Here Ω_μ^V is the connection matrix and J_{cd} the Lorentz group generators in the vector representation:

$$[J_{cd}]^{ab} = i(\delta_c^a \delta_d^b - \delta_d^a \delta_c^b). \quad (3.61)$$

The spin connection is representation-independent and the same as that in Eq. (3.52). Eqs. (3.60) and (3.61) yield

$$[\Omega_\mu^V]^{ab} = -\omega_\mu^{ab} \quad (3.62)$$

and so Eq. (3.57) reduces to

$$\mathcal{D}_\mu V^a = \partial_\mu V^a - \omega_\mu{}^a{}_b V^b. \quad (3.63)$$

For completeness we now give the transformation of $\omega_{\mu ab}$ (which are the analogues of the gauge fields of Yang-Mills theory) under an infinitesimal local LT, parametrized by

$$\Lambda^a{}_b(x) = \delta^a_b + \theta^a{}_b(x). \quad (3.64)$$

Eq. (3.41) then gives the condition

$$\theta_{ba} = -\theta_{ab}. \quad (3.65)$$

where $\theta_{ab} \equiv \eta_{ac}\theta^c{}_b$. Using Eq. (3.62), Eq. (3.64) and $(\Lambda^{-1})^a{}_b = \delta^a_b - \theta^a{}_b$ in Eq. (3.58), we obtain

$$\omega'_{\mu ab} = \omega_{\mu ab} + \partial_\mu \theta_{ab} - \omega_{\mu ac}\theta^c{}_b + \omega_{\mu bc}\theta^c{}_a. \quad (3.66)$$

One can check that the same transformation is obtained on using Eq. (3.52),

$$D(\Lambda(x)) = 1 - \frac{i}{2} S_{ab} \theta^{ab}(x), \quad (3.67)$$

and the fact that S_{ab} satisfy the Lorentz algebra

$$[S_{ab}, S_{cd}] = i(\eta_{bc}S_{ad} - \eta_{ac}S_{bd} - \eta_{bd}S_{ac} + \eta_{ad}S_{bc}) \quad (3.68)$$

in Eq. (3.50). (It is necessary to make $\omega'_{\mu ab}$ manifestly antisymmetric in a and b for having a complete agreement with Eq. (3.66).) Note that Eq. (3.67) is consistent with Eq. (3.64), since the analogue of the former, viz., $\Lambda^a{}_b(x) = \delta^a_b - \frac{i}{2}[J_{cd}]^a{}_b \theta^{cd}(x)$ leads to the latter on use of Eq. (3.61).

Let us now return to Eq. (3.63). From this equation, it is straightforward to write down the covariant derivative of the vierbein:

$$\mathcal{D}_\mu e^a{}_\nu = \partial_\mu e^a{}_\nu - \Gamma_{\mu\nu}^\lambda e^a{}_\lambda - \omega_\mu{}^a{}_b e^b{}_\nu \quad (3.69)$$

Here we have added the usual affine connection term for the covariant derivative of a coordinate vector. Since the vierbein is like the square-root of the curved space metric, it is natural to demand that its covariant derivative be zero. Put in another way,

$$\mathcal{D}_\mu e^a{}_\nu = 0 \quad (3.70)$$

ensures that the covariant derivative of the metric is zero, since Eq. (3.45) holds and the covariant derivative obeys the Leibniz rule. The spin connections can now be determined in terms of the vierbeins by making use of the above condition.

Actually, instead of using the condition (3.70), it is more convenient to use the weaker condition

$$\mathcal{D}_\mu e_{a\nu} - \mathcal{D}_\nu e_{a\mu} = 0. \quad (3.71)$$

The antisymmetrization on the L.H.S. of Eq. (3.71) ensures that the Christoffel symbol does not appear when we use Eq. (3.69) in it:

$$\partial_\mu e_{a\nu} - \partial_\nu e_{a\mu} - \omega_{\mu ab} e^b{}_\nu + \omega_{\nu ab} e^b{}_\mu = 0. \quad (3.72)$$

Now we multiply with $e_c{}^\mu e_d{}^\nu e^a{}_\lambda$ to get

$$e_c{}^\mu e_d{}^\nu e^a{}_\lambda (\partial_\mu e_{a\nu} - \partial_\nu e_{a\mu}) + \omega_{\mu da} e_c{}^\mu e^a{}_\lambda - \omega_{\nu ca} e_d{}^\nu e^a{}_\lambda = 0. \quad (3.73)$$

Here we used the vierbein orthonormality relation Eq. (3.46). Next we use Eq. (3.72) to replace $\omega_{\mu da} e^a{}_\lambda$ and $\omega_{\nu ca} e^a{}_\lambda$ in Eq. (3.73). After using Eq. (3.46) again, this gives

$$\omega_{\mu ab} = \frac{1}{2} \left[e_a{}^\nu (\partial_\nu e_{b\mu} - \partial_\mu e_{b\nu}) - e_b{}^\nu (\partial_\nu e_{a\mu} - \partial_\mu e_{a\nu}) + e_a{}^\nu e_b{}^\lambda e^c{}_\mu (\partial_\nu e_{c\lambda} - \partial_\lambda e_{c\nu}) \right]. \quad (3.74)$$

Finally we note that the Lagrangian given in Eq. (3.55) accounts only for the gravitational interaction of the Dirac fermion. Electromagnetic interaction is introduced in the usual way by further demanding invariance under $U(1)_{\text{EM}}$ gauge transformation:

$$\mathcal{L}_{\text{G,EM}}^{(f)} = \sqrt{-g} \left[\left(\frac{i}{2} \bar{\psi} \gamma^a e_a{}^\mu (\partial_\mu + ie \mathcal{Q}_f A_\mu + \frac{i}{4} \sigma^{bc} \omega_{\mu bc}) \psi + \text{h.c.} \right) - m_f \bar{\psi} \psi \right] \quad (3.75)$$

where $\mathcal{Q}_f$ is the charge of the fermion. Since A_μ is a local Lorentz scalar and a coordinate vector, the new term maintains the symmetries we already had.

3.5 Interaction vertices

QED in curved space is governed by the Lagrangians $\mathcal{L}_G^{(\gamma)}$ and $\mathcal{L}_{\text{G,EM}}^{(f)}$ given by Eqs. (3.38) and (3.75) respectively. In this section, we expand these Lagrangians to $O(\kappa)$ in the approximation of linearized gravity, and obtain the Feynman rules for the lowest order interaction of the graviton with photons and fermions.




Figure 3.1: Feynman rule for the coupling of the photon with the graviton. $C_{\mu\nu\lambda\rho}(k, k')$ is given in Eq. (3.79).

3.5.1 Photon-photon-graviton vertex

On use of Eqs. (3.11) and (3.10) in Eq. (3.38), the $O(\kappa)$ term in $\mathcal{L}_G^{(\gamma)}$ turns out to be

$$\mathcal{L}_{AAh} = -\kappa h^{\lambda\rho}(x) \hat{T}_{\lambda\rho}^{(A)}(x) \quad (3.76)$$

with the energy-momentum tensor operator for the photon given by

$$\hat{T}_{\lambda\rho}^{(A)} = \frac{1}{4} \eta_{\lambda\rho} F_{\alpha\beta} F^{\alpha\beta} + F_{\lambda\alpha} F^{\alpha}_{\rho}. \quad (3.77)$$

Writing in terms of the photon field,

$$\begin{aligned} \hat{T}_{\lambda\rho}^{(A)} = & \frac{1}{2} \eta_{\lambda\rho} [(\partial_\alpha A_\beta)(\partial^\alpha A^\beta) - (\partial_\alpha A_\beta)(\partial^\beta A^\alpha)] + (\partial_\lambda A_\alpha)(\partial^\alpha A_\rho) + (\partial_\rho A_\alpha)(\partial^\alpha A_\lambda) \\ & - (\partial_\alpha A_\lambda)(\partial^\alpha A_\rho) - (\partial_\lambda A_\alpha)(\partial_\rho A^\alpha) \end{aligned} \quad (3.78)$$

Let us now consider the vertex given in Fig. 3.1. Each term of $\mathcal{L}_{AAh}$ gives two contributions to this vertex, depending on which of the two photon field operators in the term destroys the photon and which creates it. Let us take, as an example, the $(\partial_\alpha A_\beta)(\partial^\alpha A^\beta)$ part in Eq. (3.78). When A_β destroys the photon of momentum k and A^β creates the photon of momentum k' , the contribution is $(-ik_\alpha)\eta_{\beta\mu}(ik'^\alpha)\delta_\nu^\beta$; while for the reverse case, it is $(ik'_\alpha)\eta_{\beta\nu}(-ik^\alpha)\delta_\mu^\beta$. Adding up, we find that the total contribution to the vertex from the $-\kappa h^{\lambda\rho} \frac{1}{2} \eta_{\lambda\rho} (\partial_\alpha A_\beta)(\partial^\alpha A^\beta)$ term of $\mathcal{L}_{AAh}$ is $-i\kappa \eta_{\lambda\rho} \eta_{\mu\nu} k \cdot k'$. Carrying out a similar exercise for the other terms of $\mathcal{L}_{AAh}$, we finally obtain [37, 42]

$$\begin{aligned} C_{\mu\nu\lambda\rho}(k, k') = & \eta_{\lambda\rho}(\eta_{\mu\nu} k \cdot k' - k'_\mu k_\nu) - \eta_{\mu\nu}(k_\lambda k'_\rho + k'_\lambda k_\rho) + k_\nu(\eta_{\lambda\mu} k'_\rho + \eta_{\rho\mu} k'_\lambda) \\ & + k'_\mu(\eta_{\lambda\nu} k_\rho + \eta_{\rho\nu} k_\lambda) - k \cdot k'(\eta_{\lambda\mu} \eta_{\rho\nu} + \eta_{\lambda\nu} \eta_{\rho\mu}). \end{aligned} \quad (3.79)$$

3.5.2 Fermion-fermion-graviton vertex

This vertex emerges from the Lagrangian $\mathcal{L}_{\text{G,EM}}^{(f)}$ given by Eq. (3.75), which involves the vierbein. In the weak field approximation, the vierbein can be written as

$$e^a{}_\mu(x) = \delta^a_\mu + \kappa \phi^a{}_\mu(x). \quad (3.80)$$

Here and below, we drop terms of higher order than what is necessary to determine the vierbein to $O(\kappa)$. Note that δ^a_μ , and hence $\phi^a{}_\mu$, are neither local Lorentz vectors nor coordinate vectors, but only flat space tensors of rank two. Consequently, *all* tensors in an expression should be regarded only as flat space tensors *after* we use the approximation of Eq. (3.80) to perform an expansion of the expression in powers of κ . The distinction between greek and latin indices would then be dropped, and they would be raised and lowered with the flat space metric.

The vierbein is determined from the orthonormality relation given by Eq. (3.45) or Eq. (3.46). Thus, use of Eqs. (3.80) and (3.2) in Eq. (3.45) leads to

$$\phi_{\mu\nu} + \phi_{\nu\mu} = 2h_{\mu\nu}. \quad (3.81)$$

Eq. (3.81) does not determine the antisymmetric part of $\phi_{\mu\nu}$. This freedom corresponds to the fact that the curved space action for the Dirac field is invariant under local LT: a new vierbein, given by Eq. (3.44), is as good as the old one. The ambiguity is removed by imposing the condition

$$\phi_{\mu\nu} = \phi_{\nu\mu}. \quad (3.82)$$

This is called the Lorentz symmetric gauge [43]. Let us now dwell a little on this condition.

The allowed local LT's in the weak field approximation are only those which sustain the closeness of the vierbein to the unit matrix. So we consider the local LT

$$\Lambda^a{}_b(x) = \delta^a_b + \kappa \theta^a{}_b(x). \quad (3.83)$$

Using this and Eq. (3.80) in Eq. (3.44), we arrive at the transformation

$$\phi'_{a\mu} = \phi_{a\mu} + \theta_{a\mu}. \quad (3.84)$$

Because $\theta_{a\mu}$ is antisymmetric (see Eq. (3.65)), we can always choose it to cancel out the antisymmetric part of $\phi_{a\mu}$ and implement the gauge condition given by Eq. (3.82). Another point to note is that the gauge condition does not allow any further transformation

like (3.84), and so it does break the local LT invariance. Finally we remark that the harmonic gauge condition, which was implemented earlier using general coordinate transformation invariance, is not disturbed by the implementation of the Lorentz symmetric gauge condition, since $g_{\mu\nu}$ is a local Lorentz scalar.

Let us now return to Eq. (3.80). Since Eqs. (3.81) and (3.82) give $\phi_{\mu\nu} = h_{\mu\nu}$, we have

$$e^a{}_\mu(x) = \delta^a_\mu + \kappa h^a{}_\mu(x). \quad (3.85)$$

This implies that $e_{a\mu} = \eta_{a\mu} + \kappa h_{a\mu}$. And use of Eq. (3.10) in $e_a{}^\mu = g^{\mu\nu} e_{a\nu}$ gives

$$e_a{}^\mu = \delta_a^\mu - \kappa h_a{}^\mu. \quad (3.86)$$

Now we are in a position to expand the Lagrangian $\mathcal{L}_{\text{G,EM}}^{(f)}$ to $O(\kappa)$. First consider the terms in Eq. (3.75) involving the spin connection:

$$-\frac{1}{8}\sqrt{-g}e_a{}^\mu\omega_{\mu bc}(\bar{\psi}\gamma^a\sigma^{bc}\psi + \text{h.c.}). \quad (3.87)$$

On using the weak field expressions for the vierbeins in Eq. (3.74), we obtain

$$\sqrt{-g}e_a{}^\mu\omega_{\mu bc} = \kappa(\partial_c h_{ba} - \partial_b h_{ca}). \quad (3.88)$$

Also,

$$\begin{aligned} \bar{\psi}\gamma^a\sigma^{bc}\psi + \text{h.c.} &= \bar{\psi}[\gamma^a, \sigma^{bc}]_+\psi \\ &= \frac{i}{2}\bar{\psi}\left((\gamma^a\gamma^b\gamma^c - \gamma^c\gamma^b\gamma^a) + (\gamma^b\gamma^c\gamma^a - \gamma^a\gamma^c\gamma^b)\right)\psi \\ &= 2\epsilon^{abcd}\bar{\psi}\gamma_d\gamma_5\psi, \end{aligned} \quad (3.89)$$

using

$$\gamma^a\gamma^b\gamma^c = \eta^{ab}\gamma^c + \eta^{bc}\gamma^a - \eta^{ca}\gamma^b - i\epsilon^{abcd}\gamma_d\gamma_5. \quad (3.90)$$


The product of the expressions given by Eq. (3.88) and (3.89) vanish because the graviton field is symmetric. Thus the expression (3.87) vanishes at $O(\kappa)$.

The other terms in Eq. (3.75) not involving A_μ are

$$\sqrt{-g}\left[\left(\frac{i}{2}\bar{\psi}\gamma^a e_a{}^\mu\partial_\mu\psi + \text{h.c.}\right) - m_f\bar{\psi}\psi\right]. \quad (3.91)$$

Using Eqs. (3.11) and (3.86), the $O(\kappa)$ term in the above expression turns out to be

$$\mathcal{L}_{fh} = -\kappa h^{\lambda\rho}(x)\hat{T}_{\lambda\rho}^{(f)}(x) \quad (3.92)$$



$$\text{Feynman diagram: } \frac{\text{fermion line } f(p) \rightarrow f(p')}{\text{graviton line } h_{\lambda\rho}} = -i\kappa V_{\lambda\rho}(p, p')$$

Figure 3.2: Feynman rule for the tree-level gravitational vertex of a fermion. $V_{\lambda\rho}(p, p')$ is given in Eq. (3.95).

with the energy-momentum tensor operator for the fermion given by

$$\hat{T}_{\lambda\rho}^{(f)}(x) = \left(\frac{i}{4} \bar{\psi}(x) [\gamma_\lambda \partial_\rho + \gamma_\rho \partial_\lambda] \psi(x) + \text{h.c.} \right) - \eta_{\lambda\rho} \mathcal{L}_0^{(f)}(x). \quad (3.93)$$

(see Eq. (3.47) for $\mathcal{L}_0^{(f)}$). Here we have explicitly symmetrized the term within square brackets in Eq. (3.93), making use of the fact that it comes multiplied with $h^{\lambda\rho}$. Writing out in full,

$$\begin{aligned} \hat{T}_{\lambda\rho}^{(f)} &= \frac{i}{4} \left(\bar{\psi} \gamma_\lambda \partial_\rho \psi + \bar{\psi} \gamma_\rho \partial_\lambda \psi - (\partial_\rho \bar{\psi}) \gamma_\lambda \psi - (\partial_\lambda \bar{\psi}) \gamma_\rho \psi \right) \\ &\quad - \eta_{\lambda\rho} \left(\frac{i}{2} (\bar{\psi} \gamma^\mu \partial_\mu \psi - (\partial_\mu \bar{\psi}) \gamma^\mu \psi) - m_f \bar{\psi} \psi \right). \end{aligned} \quad (3.94)$$

Let us now consider the vertex given in Fig. 3.2. Each term of $\mathcal{L}_{fgh}$ gives a contribution to this vertex, with ψ destroying the fermion and $\bar{\psi}$ creating it. We thus obtain [43, 44, 11]

$$V_{\lambda\rho}(p, p') = \frac{1}{4} [\gamma_\lambda (p + p')_\rho + \gamma_\rho (p + p')_\lambda] - \frac{1}{2} \eta_{\lambda\rho} [(\not{p} - m) + (\not{p}' - m)]. \quad (3.95)$$

3.5.3 Fermion-fermion-photon-graviton vertex

The term involving A_μ in Eq. (3.75) is

$$-e \mathcal{Q}_f \sqrt{-g} e_a{}^\mu \bar{\psi} \gamma^a \psi A_\mu. \quad (3.96)$$

Using Eqs. (3.11) and (3.86), the $O(\kappa)$ term in this expression turns out to be [42, 44]

$$\mathcal{L}_{f f A h} = -e \kappa \mathcal{Q}_f a_{\mu\nu\lambda\rho} h^{\lambda\rho} \bar{\psi} \gamma^\nu \psi A^\mu, \quad (3.97)$$

with

$$a_{\mu\nu\lambda\rho} = \eta_{\mu\nu} \eta_{\lambda\rho} - \frac{1}{2} (\eta_{\mu\lambda} \eta_{\nu\rho} + \eta_{\mu\rho} \eta_{\nu\lambda}). \quad (3.98)$$

Here we have explicitly symmetrized $a_{\mu\nu\lambda\rho}$ in λ and ρ . The Lagrangian of Eq. (3.97) gives rise to the fermion-fermion-photon-graviton vertex with the vertex factor $-ie\kappa \mathcal{Q}_f a_{\mu\nu\lambda\rho} \gamma^\nu$.




Figure 3.3: Diagrams of $O(e\kappa)$ with two fermions, one photon and one graviton in the external lines. These are the tree-level diagrams for the photoproduction of gravitons off electrons.

3.5.4 Important features of the vertex factors

All the three vertex factors deduced by us are seen to be symmetric under the exchange of λ and ρ . This is because they represent the coupling to $h_{\lambda\rho}$, which is a symmetric tensor. We now proceed to discuss the other important features of these vertex factors.

The photon-photon-graviton vertex factor $-i\kappa C_{\mu\nu\lambda\rho}(k, k')$, with $C_{\mu\nu\lambda\rho}(k, k')$ given by Eq. (3.79), is also symmetric under the simultaneous exchange of μ and ν , and, k and k' . This reflects the Bose symmetry under the interchange of the two photons. The vertex factor also satisfies electromagnetic gauge invariance:

$$k^\mu C_{\mu\nu\lambda\rho}(k, k') = 0 = k'^\nu C_{\mu\nu\lambda\rho}(k, k'). \quad (3.99)$$

It also satisfies gravitational gauge invariance:

$$(k - k')^\lambda \epsilon^\mu(k) \epsilon^\nu(k') C_{\mu\nu\lambda\rho}(k, k') = 0 = (k - k')^\rho \epsilon^\mu(k) \epsilon^\nu(k') C_{\mu\nu\lambda\rho}(k, k'), \quad (3.100)$$

$k - k'$ being the four-momentum of the graviton. In writing Eq. (3.100), we have taken the polarization vectors of the photons to be real. To prove Eq. (3.100), one has to use the on-shell conditions $k^2 = 0 = k'^2$ and $k \cdot \epsilon(k) = 0 = k' \cdot \epsilon(k')$ for the photons.

It may be noted that to prove a given kind of gauge invariance, the particles carrying the corresponding charge must be put on-shell. Since neither the photon nor the graviton carries electric charge, it was not necessary to put them on-shell in Eq. (3.99). On the other hand, the photon has gravitational coupling, i.e. carries “gravitational charge” (this is in fact true for any particle, as all particles possess energy). So the photons were put on-shell in Eq. (3.100).

We now turn to the fermion-fermion-graviton vertex factor $-i\kappa V_{\lambda\rho}(p, p')$, with $V_{\lambda\rho}(p, p')$ given by Eq. (3.95). Here again one can check gravitational gauge invariance:

$$(p - p')^\lambda \bar{u}(\vec{P}') V_{\lambda\rho}(p, p') u(\vec{P}) = 0 = (p - p')^\rho \bar{u}(\vec{P}') V_{\lambda\rho}(p, p') u(\vec{P}) \quad (3.101)$$

with the electrons put on-shell for reasons stated just now.

Finally we consider the fermion-fermion-photon-graviton vertex factor $ie\kappa a_{\mu\nu\lambda\rho}\gamma^\nu$ with $a_{\mu\nu\lambda\rho}$ given by Eq. (3.98). The checking of gauge invariance is now less trivial. This is because there are three other diagrams of $O(e\kappa)$ with the same external lines. These four diagrams, given in Fig. 3.3, represent the photoproduction of gravitons off electrons at tree-level (note that the third diagram from the left is just the vertex under consideration). When all the four diagrams are taken together, it is possible to prove electromagnetic and gravitational gauge invariance [42].

CHAPTER 4

Charged leptons in thermal QED

4.1 Dispersion relation and inertial mass

We saw in Chapter 2 that in the real-time formulation of thermal field theory, there are four free-field propagators. Similarly there are four full or exact propagators with the Heisenberg picture fields replacing the interaction picture fields in the definitions of the free-field propagators. There are also four self-energies Σ_{ab} with $-i\Sigma_{ab}$ ($a, b = 1, 2$) denoting the 1PI self-energy diagram in which the momentum enters at a b -type vertex and leaves at an a -type vertex. (We are now dealing with the specific case of fermions.) Thus the free-field propagator, the full or exact propagator and the self-energy are all 2×2 matrices [28, 30]. The poles of the full propagator for a fermion of momentum p_μ occur at the poles of the function

$$S'_f(p) = [\not{p} - m_f - \Sigma_f(p)]^{-1} \quad (4.1)$$

where $\Sigma_f(p)$ is related to the components of the self-energy matrix. Let us define

$$\text{Re } \Sigma_f \equiv \frac{1}{2}(\Sigma_f + \gamma^0 \Sigma_f^\dagger \gamma^0) \quad (4.2)$$

and

$$\text{Im } \Sigma_f \equiv \frac{1}{2i}(\Sigma_f - \gamma^0 \Sigma_f^\dagger \gamma^0), \quad (4.3)$$

and likewise for Σ_{ab} . Then the aforementioned relations are [28, 30]

$$\text{Re } \Sigma_f(p) = \text{Re } \Sigma_{11}(p), \quad (4.4)$$

$$\text{Im } \Sigma_f(p) = \frac{\text{Im } \Sigma_{11}(p)}{1 - 2\eta_f(p)}, \quad (4.5)$$

with $\eta_f(p)$ given by Eq. (2.51).

Considering the analogy of Eq. (4.1) to the vacuum case, we shall call $\Sigma_f(p)$ the *self-energy of the fermion*. It has the general form

$$\Sigma_f(p) = a\not{p} + b\not{v} + c \quad (4.6)$$

at one loop in an isotropic medium with four-velocity v_μ . Here a , b and c are Lorentz-invariant functions of the two Lorentz scalars

$$\omega \equiv p \cdot v, \quad (4.7)$$

$$P \equiv [(p \cdot v)^2 - p^2]^{1/2}. \quad (4.8)$$

ω and P are the Lorentz-invariant energy and three-momentum respectively, which satisfy

$$p^2 = \omega^2 - P^2. \quad (4.9)$$

Eq. (4.6) can contain an additional term proportional to $\sigma^{\mu\nu}p_\mu v_\nu$ in the more general case. However, such a term does not appear at the level of the one-loop calculations [5] that we are considering in this work, and therefore we omit it.

We shall be interested only in the dispersive part of the self-energy. This means that the corresponding term $\bar{\psi}\Sigma_f\psi$ of the Lagrangian in momentum space is hermitian, which translates to the condition

$$\Sigma_f(p) = \gamma^0 \Sigma_f^\dagger(p) \gamma^0. \quad (4.10)$$

Thus, the dispersive part of the self-energy is given by $\text{Re } \Sigma_f$. On the other hand, when one is also interested in the absorption of the fermion in the medium, the Lagrangian must no longer be hermitian, and so one has to consider the entire Σ_f . For the relation between $\text{Im } \Sigma_f$ and the damping rate of the fermion, see, for example, Refs. [45] and [23]. We shall deal only with $\text{Re } \Sigma_f$ and *by Σ_f , we shall henceforth refer only to $\text{Re } \Sigma_f$* . Clearly, the Lorentz scalars a , b and c of Eq. (4.6) are real in our case.

Now $S'_f(p)$ of Eq. (4.1) can be written in the form

$$S'_f(p) = N(p)[D(p)]^{-1} \quad (4.11)$$

with

$$N(p) = (1 - a)\not{p} - b\not{v} + (m_f + c), \quad (4.12)$$

$$D(p) = [(1-a)p - bv]^2 - (m_f + c)^2. \quad (4.13)$$

The poles of $S'_f(p)$ are given by $D(p) = 0$. At $O(e^2)$ (i.e., on keeping only the terms independent of or linear in a , b and c), this leads to

$$\omega^2 = P^2 + m_f^2 + 2[a(\omega^2 - P^2) + b\omega + m_f c] \quad (4.14)$$

$$= P^2 + m_f^2 + \frac{1}{2} \text{Tr}[(\not{p} + m_f)\Sigma_f]. \quad (4.15)$$

Taking the square root, we obtain the pair of solutions

$$\omega_{\pm} = \pm \sqrt{P^2 + m_f^2} \pm \frac{1}{4\sqrt{P^2 + m_f^2}} \text{Tr}[(\not{p} + m_f)\Sigma_f]. \quad (4.16)$$

These are implicit equations for ω as function of P , since the R.H.S. contains a , b and c which are again functions of ω . We now write

$$\omega_{\pm}(P) = \pm E_{f,\bar{f}}(P) \quad (4.17)$$

Eq. (4.16) can then be used to solve for $E_f(P)$ and $E_{\bar{f}}(P)$. These are the dispersion relations corresponding to the fermion and the antifermion respectively. In perturbation theory, the justification for Eq. (4.17) is that it reduces to the usual dispersion relations at the tree-level. (See also Secs. 4.3 and 4.4 on charge conjugation. For justification at a non-perturbative level, see the last paragraph of this section.) The inertial masses are defined as

$$M_{f,\bar{f}} \equiv E_{f,\bar{f}}(0) = \pm \omega_{\pm}(0). \quad (4.18)$$

Eq. (4.16) then gives

$$M_{f,\bar{f}} = m_f + \frac{1}{4m_f} \left\{ \text{Tr}[(\not{p} + m_f)\Sigma_f] \right\}_{\substack{\omega=\pm m_f \\ P=0}}, \quad (4.19)$$

using the tree-level values of ω in the $O(e^2)$ part.

We shall perform all calculations in the rest frame of the medium, in which v^μ has the components given by Eq. (2.33), and in that frame, we define the components of p^μ by writing

$$p^\mu = (p^0, \vec{P}). \quad (4.20)$$

(Eq. (4.8) shows that the magnitude of the rest-frame three-momentum $\vec{P}$ is indeed equal to the Lorentz-invariant three-momentum P . So our notations are consistent.) Then

a, b, c are functions of the variables p^0 and P , which we will indicate by writing them as $a(p^0, P)$, and similarly for the other ones, when we need to show it explicitly. In the same way, $N(p)$ and $D(p)$ of Eqs. (4.12) and (4.13) will be written as $N(p^0, \vec{P})$ and $D(p^0, P)$. Since the condition $\omega = \pm m_f$ now reduces to $p^0 = \pm m_f$, we have

$$M_{f,\bar{f}} = m_f + \frac{1}{4} \left\{ \text{Tr}[(1 \pm \gamma^0) \Sigma_f] \right\}_{p^\mu = (\pm m_f, \vec{0})} \quad (4.21)$$

Defining

$$\mathcal{E}_{f,\bar{f}}(p^0, \vec{P}) = \frac{1}{4} \text{Tr}[(\gamma^0 \pm 1) \Sigma_f] \quad (4.22)$$

$$= ap^0 + b \pm c, \quad (4.23)$$

the inertial masses to $O(e^2)$ are given by

$$\begin{aligned} M_f &= m_f + \mathcal{E}_f(m_f, \vec{0}), \\ M_{\bar{f}} &= m_f - \mathcal{E}_{\bar{f}}(-m_f, \vec{0}). \end{aligned} \quad (4.24)$$

Eq. (4.22) is a useful formula that allows us to extract the matter-induced corrections to the inertial mass directly from the one-loop expression for Σ_f . As we will see later, the wavefunction renormalization factor is determined in terms of the same quantities $\mathcal{E}_f$ and $\mathcal{E}_{\bar{f}}$.

It may be noted CPT symmetry, if present, ensures that $E_f(P) = E_{\bar{f}}(P)$, and consequently, $M_f = M_{\bar{f}}$. However, normal matter is CPT -asymmetric, and, as we shall see from explicit calculations, M_f no longer equals $M_{\bar{f}}$.

The antifermion dispersion relation of Eq. (4.17) and the hole dispersion relation of Ref. [46] (see also Ref. [47]) are based on the same concept, namely, the removal of a particle from the thermal background. However, Ref. [46] considers a different situation: high-temperature massless QED (or QCD) at zero chemical potential. In this situation, while one can decompose the propagator as $D(p) = (\omega - \omega_+)(\omega - \omega_-)$ such that $\omega_\pm = \pm P$ at the tree-level, there are some important differences from our case. Firstly, one gets two dispersion relation from ω_+ itself. One relation, of course, is obtained by writing $\omega_+ = E_f(P)$. The other is obtained by writing $\omega_+ = -E_h(P)$. The form of ω_+ is such that there is an $E_h(P)$ which is positive for all P . This is the hole state, also called plasmino [48]. Secondly, ω_- does not give dispersion relations distinct from those obtained from ω_+ , since the medium is CPT symmetric.

4.2 Spinor and wavefunction renormalization factor

Consider the momentum-space Dirac equation

$$(p^0 \gamma^0 - \vec{P} \cdot \vec{\gamma} - m_f - \Sigma_f(p^0, \vec{P})) \xi(\vec{P}) = 0 \quad (4.25)$$

in a medium. Requiring Eq. (4.25) to have non-trivial solutions yields the condition $D(p^0, P) = 0$ (see Eq. (4.13)). This equation, as we have seen, possesses the two solutions $p_0 = E_f(P)$ and $p_0 = -E_{\bar{f}}(P)$. Now $\xi(\vec{P})$ for $p_0 = E_f(P)$ is the fermion spinor $U_s(\vec{P})$. On the other hand, $\xi(\vec{P})$ for $p_0 = -E_{\bar{f}}(P)$ is the antifermion spinor at three-momentum $-\vec{P}$.

Thus, the fermion spinor $U_s(\vec{P})$ in a medium satisfies the equation

$$(\not{p} - m_f - \Sigma_f(p)) U_s(\vec{P}) = 0 \quad (4.26)$$

with $p^\mu = (E_f(P), \vec{P})$. We choose the normalization

$$U_s^\dagger(\vec{P}) U_s(\vec{P}) = 1. \quad (4.27)$$

The hermitian conjugate of Eq. (4.26) is

$$\overline{U}_s(\vec{P}) (\not{p} - m_f - \Sigma_f(p)) = 0 \quad (4.28)$$

where the condition (4.10) was used. We now use the general form of Eq. (4.6). Multiplying Eq. (4.26) from the left with $\overline{U}_s(\vec{P}) \gamma_\mu$ and Eq. (4.28) from the right with $\gamma_\mu U_s(\vec{P})$, and adding the two resulting equations, one then arrives at the identity

$$\overline{U}_s(\vec{P}) \gamma_\mu U_s(\vec{P}) = \left[\frac{(1-a)p_\mu - bv_\mu}{m_f + c} \right] \overline{U}_s(\vec{P}) U_s(\vec{P}). \quad (4.29)$$

The time-component of Eq. (4.29), together with the normalization adopted in Eq. (4.27), give

$$\overline{U}_s(\vec{P}) U_s(\vec{P}) = \frac{m_f + c}{(1-a)E_f - b}, \quad (4.30)$$

which, when fed back into Eq. (4.29), yields

$$\overline{U}_s(\vec{P}) \gamma_\mu U_s(\vec{P}) = \frac{(1-a)p_\mu - bv_\mu}{(1-a)E_f - b}. \quad (4.31)$$

For future reference, we note that in particular, in the frame specified by Eq. (2.33),

$$[\overline{U}_s(\vec{P}) U_s(\vec{P})]_{\vec{P}=0} = 1 \quad (4.32)$$

(use Eq. (4.23) and the first of Eqs. (4.24), or use Eq. (4.34) below), and

$$\left[\overline{U}_s(\vec{P}) \gamma_\mu U_s(\vec{P}) \right]_{\vec{P}=0} = v_\mu. \quad (4.33)$$

From Eq. (4.26), the explicit form of $U_s(\vec{P})$ can easily be worked out in the rest frame of the medium, using any particular representation of the gamma matrices. The condition for Eq. (4.26) to possess non-trivial solution is given by

$$[(1-a)E_f - b]^2 = (1-a)^2 P^2 + (m_f + c)^2 \quad (4.34)$$

which, of course, is nothing but the equation $D(p_0, P) = 0$ with $p_0 = E_f$. The U -spinors in the Dirac-Pauli representation are then given by

$$U_\pm(\vec{P}) = \sqrt{\frac{(1-a)E_f - b + m_f + c}{2[(1-a)E_f - b]}} \begin{pmatrix} \chi_\pm \\ \frac{(1-a)\vec{\sigma} \cdot \vec{P}}{(1-a)E_f - b + m_f + c} \chi_\pm \end{pmatrix} \quad (4.35)$$

with the prefactor chosen in accord with the normalization condition of Eq. (4.27). Here

$$\chi_+ = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \chi_- = \begin{pmatrix} 0 \\ 1 \end{pmatrix}. \quad (4.36)$$

The solutions in Eq. (4.35) can be used to find that

$$\sum_s U_s(\vec{P}) \overline{U}_s(\vec{P}) = \frac{N(E_f, \vec{P})}{2[(1-a)E_f - b]}. \quad (4.37)$$

This relation is representation-independent.

Eq. (4.37) is useful in finding out the fermion wavefunction renormalization factor $Z_f(P)$. In a medium Z_f is defined through the behaviour of the propagator at the one-particle pole $p_0 = E_f(P)$ [46]:

$$S'_f(p) \Big|_{1\text{-particle}} \approx \frac{Z_f(P) \sum_s U_s(\vec{P}) \overline{U}_s(\vec{P})}{p_0 - E_f}. \quad (4.38)$$

(See App. B for a proof of this relation in vacuum.) Note that we have explicitly indicated the fact that Z_f depends on $\vec{P}$, but only on the magnitude of it, in an isotropic medium. This will be checked below. Near the pole considered in Eq. (4.38), Eq. (4.11) reduces to

$$S'_f(p) \approx \frac{N(E_f, \vec{P})}{(p_0 - E_f) \left(\frac{\partial D}{\partial p_0} \right)_{p^0=E_f}}. \quad (4.39)$$

The requirement that the residues of these two expressions coincide, then yields

$$Z_f(P) = \left\{ 2[(1-a)E_f - b] \left(\frac{\partial D}{\partial p_0} \right)^{-1} \right\}_{p^0=E_f}, \quad (4.40)$$

where we have used Eq. (4.37). Note that since a , b , c and (hence) D depend on $\vec{P}$ only through P , so also does Z_f . Now use

$$D(p_0, P) = (1-2a)(p_0^2 - P^2) - 2bp_0 - m_f^2 - 2m_f c \quad (4.41)$$

to $O(e^2)$. Then, to this order,

$$Z_f(P) = 1 + \left[a + \left(p_0 - \frac{P^2}{p_0} \right) \frac{\partial a}{\partial p_0} + \frac{\partial b}{\partial p_0} + \frac{m_f}{p_0} \frac{\partial c}{\partial p_0} \right]_{p_0=E_f}. \quad (4.42)$$

For the particular case $\vec{P} = 0$ in which we shall be interested, Eq. (4.42) reduces to

$$Z_f = 1 + \zeta_f, \quad (4.43)$$

where

$$\zeta_f = \left. \frac{\partial \mathcal{E}_f}{\partial p^0} \right|_{p^\mu=(m_f, \vec{0})}, \quad (4.44)$$

with $\mathcal{E}_f$ given by Eq. (4.22) or Eq. (4.23). From now on whenever we omit the dependence of Z_f on P in an equation, it is to be understood as the quantity evaluated at $P = 0$.

In this context, we note that in vacuum one can use the result $Z_f^{-1} = 1 - (d\Sigma_f/d\not{p})_{\not{p}=m}$ just as well as Eq. (4.40) in order to determine the wavefunction renormalization factor. But since in a medium a , b and c depend on $p \cdot v$, one cannot define $d\Sigma_f/d\not{p}$ unambiguously, and so Eq. (4.40) is the result to be used in the present case.

We now consider the case of the antiparticles. We have already defined the antifermion spinor in the beginning of this section (see after Eq. (4.25)). Thus, the antifermion spinor $V_s(\vec{P})$ satisfies the equation

$$(\not{p} + m_f + \Sigma_f(-p))V_s(\vec{P}) = 0, \quad (4.45)$$

where $p^\mu = (E_{\vec{P}}(P), \vec{P})$. The normalization is chosen to be

$$V_s^\dagger(\vec{P})V_s(\vec{P}) = 1. \quad (4.46)$$

The analogy of Eq. (4.29) in the present case is

$$\bar{V}_s(\vec{P})\gamma_\mu V_s(\vec{P}) = - \left[\frac{(1 - a(-p))p_\mu + b(-p)v_\mu}{m_f + c(-p)} \right] \bar{V}_s(\vec{P})V_s(\vec{P}). \quad (4.47)$$

From Eqs. (4.47) and (4.46), one can obtain the expressions for $\bar{V}_s(\vec{P})V_s(\vec{P})$ and $\bar{V}_s(\vec{P})\gamma^\mu V_s(\vec{P})$ just as in the case of the U -spinors. It may be noted that since $a(p)$ depends on $\vec{P}$ only through P , the quantity $a(-p)$ essentially translates to $a(p)$ evaluated at $p_0 = -E_{\bar{f}}$. Similar remarks hold for b and c .

The condition for Eq. (4.45) to possess non-trivial solution is given by

$$[(1 - a(-p))E_{\bar{f}} + b(-p)]^2 = (1 - a(-p))^2 P^2 + (m_f + c(-p))^2 \quad (4.48)$$

which is the equation $D(p_0, P) = 0$ with $p_0 = -E_{\bar{f}}$. The normalized V -spinors in the Dirac-Pauli representation are then given by

$$V_\pm(\vec{P}) = \sqrt{\frac{(1 - a(-p))E_{\bar{f}} + b(-p) + m_f + c(-p)}{2[(1 - a(-p))E_{\bar{f}} + b(-p)]}} \times \begin{pmatrix} \frac{(1 - a(-p))\vec{\sigma} \cdot \vec{P}}{(1 - a(-p))E_{\bar{f}} + b(-p) + m_f + c(-p)} \chi'_\pm \\ \chi'_\pm \end{pmatrix}. \quad (4.49)$$

Here

$$\chi'_+ = - \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad \chi'_- = \begin{pmatrix} 1 \\ 0 \end{pmatrix}. \quad (4.50)$$

(The choice of $\chi'_\pm$ is motivated by consideration of charge conjugation; this will be explained at the end of Sec. 4.3.) Eq. (4.49) leads to

$$\sum_s V_s(\vec{P})\bar{V}_s(\vec{P}) = - \frac{N(-E_{\bar{f}}, -\vec{P})}{2[(1 - a(-p))E_{\bar{f}} + b(-p)]}. \quad (4.51)$$

$Z_{\bar{f}}(P)$ is defined through the relation

$$S'_f(p) \Big|_{1\text{-antiparticle}} \approx \frac{Z_{\bar{f}}(P) \sum_s V_s(-\vec{P})\bar{V}_s(-\vec{P})}{p_0 + E_{\bar{f}}}. \quad (4.52)$$

(See App. B for a proof of this relation in vacuum.) Near the pole considered in Eq. (4.52), Eq. (4.11) reduces to

$$S'_f(p) \approx \frac{N(-E_{\bar{f}}, \vec{P})}{(p_0 + E_{\bar{f}}) \left(\frac{\partial D}{\partial p_0} \right)_{p^0 = -E_{\bar{f}}}}. \quad (4.53)$$

Eqs. (4.51), (4.52) and (4.53) give

$$Z_{\bar{f}}(P) = - \left\{ 2 \left[(1 - a(p)) E_{\bar{f}} + b(p) \right] \left(\frac{\partial D}{\partial p_0} \right)^{-1} \right\}_{p^0 = -E_{\bar{f}}} . \quad (4.54)$$

Hence we obtain, to $O(e^2)$,

$$Z_{\bar{f}}(P) = 1 + \left[a + \left(p_0 - \frac{P^2}{p_0} \right) \frac{\partial a}{\partial p_0} + \frac{\partial b}{\partial p_0} + \frac{m_f}{p_0} \frac{\partial c}{\partial p_0} \right]_{p_0 = -E_{\bar{f}}} . \quad (4.55)$$

At $P = 0$, this reduces to

$$Z_{\bar{f}} = 1 + \zeta_{\bar{f}} , \quad (4.56)$$

where

$$\zeta_{\bar{f}} = \left. \frac{\partial \mathcal{E}_{\bar{f}}}{\partial p^0} \right|_{p^\mu = (-m_f, \vec{0})} . \quad (4.57)$$

The tree-level spinors u_s and v_s are the limiting cases of the spinors U_s and V_s when the effects of the medium are neglected. They satisfy the free Dirac equation in the vacuum, as well as the relations

$$\bar{u}_s \gamma_\mu u_s = \frac{p_\mu}{m_f} \bar{u}_s u_s \quad (4.58)$$

$$\bar{u}_s u_s = \frac{m_f}{E_f} \quad (4.59)$$

(where $p_0 = \sqrt{P^2 + m_f^2}$), with similar relations for v_s but with the substitution $p_\mu \rightarrow -p_\mu$ in the above equations.

We conclude this section by pointing out why a medium is expected to impart new features to the wavefunction renormalization factor which are absent in vacuum. Firstly, had the wavefunction renormalization factor been a function of momentum in vacuum, it would have led to new kinds of divergence, spoiling the renormalizability of the theory. But there is no such problem in a medium, since the thermal corrections do not involve ultraviolet divergences. Secondly, CPT invariance in vacuum ensures that $Z_f = Z_{\bar{f}}$. But since a medium can be CPT -asymmetric, this equality may not generally hold.

4.3 Effect of charge conjugation invariance

In this section, we consider the consequences of charge conjugation invariance for the self-energy, the dispersion relations, the wavefunction renormalization factors and the

antifermion spinor. C invariance is present when the background contains only photons, and also when it contains, in addition, charged leptons at zero chemical potential. So these consequences can be checked in such cases.

We begin by writing down the effect of the charge conjugation operator C on a spinor field:

$$\psi_C(x) \equiv C\psi(x)C = \mathbf{C}\gamma_0^T\psi^*(x). \quad (4.60)$$

(Note that $C^2 = 1$, so that $C^{-1} = C$.) C invariance of the free Dirac Lagrangian is ensured by the relation

$$\mathbf{C}^{-1}\gamma^\mu\mathbf{C} = -\gamma_\mu^T \quad (4.61)$$

where the matrix $\mathbf{C}$ is unitary:

$$\mathbf{C}^{-1} = \mathbf{C}^\dagger. \quad (4.62)$$

Using $(\psi_C)_C = \psi$, one can then show that the matrix $\mathbf{C}$ is also antisymmetric:

$$\mathbf{C}^T = -\mathbf{C}. \quad (4.63)$$

It will be useful to first go through the demonstration of the C invariance of the tree-level Dirac action

$$\mathcal{A}_f = \int d^4x \bar{\psi}(x)(i\gamma^\mu\partial_\mu - m_f)\psi(x). \quad (4.64)$$

Let us define the Fourier transform of the spinor field by

$$\psi(x) = \int \frac{d^4p}{(2\pi)^4} e^{-ip \cdot x} \psi(p). \quad (4.65)$$

Putting this in Eq. (4.64), we obtain

$$\mathcal{A}_f = \int \frac{d^4p}{(2\pi)^4} \bar{\psi}(p)(\not{p} - m_f)\psi(p). \quad (4.66)$$

Now from Eqs. (4.60) and (4.65),

$$\psi_C(x) = \int \frac{d^4p}{(2\pi)^4} e^{ip \cdot x} \mathbf{C}\gamma_0^T\psi^*(p), \quad (4.67)$$

and so,

$$\bar{\psi}_C(x) = \psi_C^\dagger(x)\gamma_0 = - \int \frac{d^4p}{(2\pi)^4} e^{-ip \cdot x} \psi^T(p)\mathbf{C}^{-1}, \quad (4.68)$$

where we used Eqs. (4.61) and (4.62). Substitution in Eq. (4.64) with $\psi \rightarrow \psi_C$ gives

$$\mathcal{A}_{fC} = - \int \frac{d^4 p}{(2\pi)^4} \psi^T(p) \mathbf{C}^{-1} (-\not{p} - m_f) \mathbf{C} \gamma_0^T \psi^*(p). \quad (4.69)$$

Note that we have $-\not{p} - m_f$ in the place of $\not{p} - m_f$ of Eq. (4.66), due to the presence of $e^{ip \cdot x}$ in Eq. (4.67) in the place of $e^{-ip \cdot x}$ in Eq. (4.65). A little rearrangement in Eq. (4.69) leads to

$$\mathcal{A}_{fC} = - \int \frac{d^4 p}{(2\pi)^4} \bar{\psi}(p) [\mathbf{C}^{-1} (\not{p} + m_f) \mathbf{C}]^T \psi(p), \quad (4.70)$$

which, on using Eq. (4.61), is found to equal $\mathcal{A}_f$, as it should.

Now consider the full Dirac action

$$\mathcal{A}'_f = \int \frac{d^4 p}{(2\pi)^4} \bar{\psi}(p) (\not{p} - m_f - \Sigma_f(p)) \psi(p), \quad (4.71)$$

Following the derivation of Eq. (4.69), we can write down

$$\mathcal{A}'_{fC} = - \int \frac{d^4 p}{(2\pi)^4} \psi^T(p) \mathbf{C}^{-1} (-\not{p} - m_f - \Sigma_f(-p)) \mathbf{C} \gamma_0^T \psi^*(p). \quad (4.72)$$

Rearranging as in the case A_{fC} and demanding that

$$\mathcal{A}'_f = \mathcal{A}'_{fC}, \quad (4.73)$$

we arrive at the condition for C invariance:

$$\Sigma_f(p) = [\mathbf{C}^{-1} \Sigma_f(-p) \mathbf{C}]^T. \quad (4.74)$$

Using the general form of $\Sigma_f(p)$ given in Eq. (4.6), and also using Eq. (4.61), it follows that the Lorentz scalars a , b and c then satisfy

$$a(-p) = a(p), \quad b(-p) = -b(p), \quad c(-p) = c(p). \quad (4.75)$$

Substitution of these conditions in Eq. (4.48) and comparison of the resulting equation with Eq. (4.34) show that E_f and $E_{\bar{f}}$ satisfy identical equations under C invariance, i.e., the fermion and antifermion dispersion relations are then the same.

Turning to the wavefunction renormalization factor, we rewrite Eq. (4.55) as

$$Z_{\bar{f}}(P) = 1 + \left[a(-p) + \left(p_0 - \frac{P^2}{p_0} \right) \frac{\partial a(-p)}{\partial p_0} - \frac{\partial b(-p)}{\partial p_0} + \frac{m_f}{p_0} \frac{\partial c(-p)}{\partial p_0} \right]_{p_0=E_{\bar{f}}}. \quad (4.76)$$

Substitution of the conditions of Eq. (4.75) and comparison with Eq. (4.42) then shows that $Z_{\bar{f}}(P) = Z_f(P)$ (on using $E_{\bar{f}} = E_f$). So C invariance also leads to the equality of the fermion and antifermion wavefunction renormalization factors.

Such symmetries between the properties of the particle and the antiparticle are, in general, the consequence of CPT invariance. However, the form of the self-energy as given by Eq. (4.6) already implies invariance of the action under parity and time reversal. So the additional imposition of C invariance through Eq. (4.75) is, in fact, equivalent to the requirement of CPT invariance.

Finally we consider the antifermion spinor. At the tree-level, the fermion spinor is related to the antifermion spinor by

$$v_s(\vec{P}) = \mathbf{C}\gamma_0^T u_s^*(\vec{P}). \quad (4.77)$$

To verify this, start with the equation $(\not{p} - m_f)u_s(\vec{P}) = 0$. Then $v_s(\vec{P})$, as defined above, satisfies

$$(p^\mu \mathbf{C}\gamma_0^T \gamma_\mu^* (\mathbf{C}\gamma_0^T)^{-1} - m_f)v_s(\vec{P}) = 0. \quad (4.78)$$

This, on using the hermitian conjugation property of the gamma matrices and then Eq. (4.61), reduces to the familiar equation $(\not{p} + m_f)v_s(\vec{P}) = 0$.

Let us now try to generalize this exercise in the presence of radiative corrections. That is, let us start with Eq. (4.26), and write

$$V_s(\vec{P}) = \mathbf{C}\gamma_0^T U_s^*(\vec{P}). \quad (4.79)$$

Then one can find an equation for $V_s(\vec{P})$ just as we found Eq. (4.78). Demanding that this equation for $V_s(\vec{P})$ be the same as Eq. (4.45) leads, on using $E_f = E_{\bar{f}}$, to the condition

$$\Sigma_f(p) = \mathbf{C}\gamma_0^T \Sigma_f^*(-p)\gamma_0^T \mathbf{C}^{-1}. \quad (4.80)$$

Use Eq. (4.10) and then Eq. (4.63) in Eq. (4.80). This reduces it to Eq. (4.74). We therefore conclude that the tree-level definition (4.77) of the antifermion spinor can be extended to (4.79) in the presence of C invariance. In the Dirac-Pauli representation, one can choose $\mathbf{C} = i\gamma^2\gamma^0$ (thereby fixing the overall phase in $\mathbf{C}$ which is left arbitrary by Eqs. (4.61), (4.62) and (4.63)), while the explicit solutions for the spinors have already been given in Eqs. (4.35) and (4.49). One can then verify Eq. (4.79), making use of the conditions of Eq. (4.75) together with $E_f = E_{\bar{f}}$.

4.4 Relations when background is not C invariant

Let us now consider the case when the background contains charged leptons at non-zero chemical potential. This breaks the invariance under charge conjugation, even though the underlying Lagrangian, being that of QED, is C invariant.

Recall from Sec. 2.2 that the effect of the medium is introduced through the operator J given by Eq. (2.4), which occurs in the density operator. The operator J is made up of H and Q , which transform under charge conjugation as

$$CHC = H, \quad CQC = -Q. \quad (4.81)$$

The relations (4.81) can be explicitly checked at the tree-level, as follows. Considering the contribution from the charged leptons only, the expressions for H and Q are those given in Eq. (A.11). Let us now consider Eq. (4.60) and substitute the expansion (2.37) into it. Using Eq. (4.77), and $u_s(\vec{P}) = C\gamma_0^T v_s^*(\vec{P})$ (following from Eq. (4.77) and the properties of the matrix C), we obtain

$$Cc_s(\vec{P})C = d_s(\vec{P}), \quad Cd_s(\vec{P})C = c_s(\vec{P}), \quad (4.82)$$

so that

$$Cc_s^\dagger(\vec{P})c_s(\vec{P})C = d_s^\dagger(\vec{P})d_s(\vec{P}), \quad Cd_s^\dagger(\vec{P})d_s(\vec{P})C = c_s^\dagger(\vec{P})c_s(\vec{P}). \quad (4.83)$$

(For these steps, it is convenient to use $C^\dagger = C$, which follows from $C^2 = 1$ and the fact that C is a unitary operator.) Use of Eqs. (4.83) in Eqs. (A.11) at once leads to Eqs. (4.81).

From Eqs. (2.4) and (4.81), it is clear that CJC does not equal J when $\mu \neq 0$. Note, however, that J stays the same if charge conjugation is followed by a reversal of the sign of μ . Consequently, while the relation in Eq. (4.73) breaks down, the relation

$$\mathcal{A}'_f = [\mathcal{A}'_{fC}]_{\mu \rightarrow -\mu} \quad (4.84)$$

holds. Eq. (4.75) is then modified to

$$[a(-p)]_{\mu \rightarrow -\mu} = a(p), \quad [b(-p)]_{\mu \rightarrow -\mu} = -b(p), \quad [c(-p)]_{\mu \rightarrow -\mu} = c(p). \quad (4.85)$$

Retracing the subsequent steps of Sec. 4.3, one then arrives at

$$[E_{\bar{f}}(P)]_{\mu \rightarrow -\mu} = E_f(P), \quad (4.86)$$

$$[Z_{\bar{f}}(P)]_{\mu \rightarrow -\mu} = Z_f(P), \quad (4.87)$$

$$[V_s(\vec{P})]_{\mu \rightarrow -\mu} = C\gamma_0^T U_s^*(\vec{P}). \quad (4.88)$$




Figure 4.1: One-loop diagrams for the self-energy of a charged lepton ℓ in a medium.

4.5 Calculations at $O(e^2)$

4.5.1 Self-energy

The self-energy diagrams are shown in Fig. 4.1. In view of Eq. (4.4), we need to concern ourselves only with the 1-type vertex. (Actually, the upper vertex of Fig. 4.1B can still be of 2-type, but we shall see that there is no contribution in this case.) The propagators for the internal lines are therefore the “11” propagators of Chap. 2. For a fermion, the propagator is thus given by Eqs. (2.48), (2.49), (2.50) and (2.51), while for the photon, the propagator in the Feynman gauge is given by Eqs. (2.52), (2.53), (2.54) and (2.55).

The contribution for Fig. 4.1B vanishes because the photon tadpole is zero. There are several reasons for the vanishing of the photon tadpole in vacuum, but, as we shall see, only one of these arguments continue to hold in normal matter. First consider Lorentz invariance. The photon tadpole must equal a four-vector, but there being no four-vector available in vacuum (the photon line of the tadpole carries zero momentum), the tadpole has to vanish. However in a medium, we do have a four-vector, namely, the four-velocity v^μ of the medium, and the tadpole can be proportional to it. Next consider invariance under the action of the charge conjugation operator C . Using $C|0\rangle = 0$ and $CA_\mu C = -A_\mu$, it is clear that $\langle 0|A_\mu|0\rangle$ vanishes. But in a medium, the expectation value is with respect to the thermal state which need not be C -invariant (normal matter, e.g., contains electrons but no positrons) and in that case this argument also is invalidated. Finally one must realize that the photon tadpole, since it has no other external lines, represents the coupling of

the photon to the net current in vacuum or in the medium, as the case may be. This net current is zero not only in vacuum, but also in an electrically neutral medium, thereby ensuring the vanishing of the photon tadpole.

As an illustration of the formal reasoning presented above, let us now explicitly evaluate the photon tadpole contained in Fig. 4.1B. The tadpole is given by

$$T^\mu = \sum_f \int \frac{d^4 k}{(2\pi)^4} \text{Tr} \left[i e \mathcal{Q}_f \gamma^\mu i S_f(k) \right]. \quad (4.89)$$

Here the sum is over all species of fermions in the medium, the charge of each species being denoted by $\mathcal{Q}_f$ with the convention that $\mathcal{Q}_e = -1$. The vacuum part of Eq. (4.89), coming from $i S_{Ff}(p)$ of Eq. (2.48), vanishes separately for each fermion in the loop. On the other hand, the thermal part, coming from $S_{Tf}(p)$ of Eq. (2.48), equals

$$T^\mu = -i 4e \sum_f \mathcal{Q}_f \int \frac{d^4 k}{(2\pi)^3} \delta(k^2 - m_f^2) \eta_f(k) k^\mu. \quad (4.90)$$

Let us write $T^\mu = v^\mu T$ where T is a scalar, so that $T = v_\mu T^\mu$. It is easy to evaluate T by going over to the rest frame given by Eq. (2.33), and performing the k_0 integration. We thus obtain

$$T^\mu = -i e v^\mu \sum_f \mathcal{Q}_f \left[2 \int \frac{d^3 K}{(2\pi)^3} [f_f(E_K) - f_{\bar{f}}(E_K)] \right] \quad (4.91)$$

The distribution functions are given by Eq. (2.43). Here, as elsewhere, we write the components of a four-vector as $k^\mu = (k^0, \vec{K})$, and use the notation $E_K \equiv \sqrt{K^2 + m_f^2}$. Since the number densities are given by

$$n_{f,\bar{f}} = 2 \int \frac{d^3 K}{(2\pi)^3} f_{f,\bar{f}}(E), \quad (4.92)$$

Eq. (4.91) shows that T^μ is proportional to the total charge density of the medium, and indeed vanishes for an electrically neutral medium such as normal matter.

At this point, we consider the case when the upper vertex in Fig. 4.1B is a 2-type vertex. Because the thermal part of the “22” propagator is the same as that of the “11” propagator, and the 2-type vertex differs from the 1-type vertex by a sign, the photon tadpole in this case is just the negative of the value given in Eq. (4.91), and therefore again vanishes in a neutral medium.

However, in anticipation of the calculation of the gravitational interaction of the fermion in Chap. 5, we mention here that in the presence of the gravitational potential, the diagram of Fig. 4.1B is not zero by itself. This is because the condition of

the vanishing of the photon tadpole (which is equivalent to require that the medium be electrically neutral) involves other diagrams. This will be discussed in detail in Sec. 5.4.3.

Let us now return to our present calculation of the self-energy (in the absence of gravitational potential), for which we need to consider only Fig. 4.1A. Thus the charged lepton self-energy is given by

$$-i\Sigma_\ell(p) = (ie)^2 \int \frac{d^4k}{(2\pi)^4} \gamma^\mu iS_\ell(p+k) \gamma^\nu iD_{\mu\nu}(k). \quad (4.93)$$

When Eqs. (2.48) and (2.52) are substituted into Eq. (4.93), four terms are produced. Since we are interested in the background induced contributions only, we disregard the term involving both $S_{F\ell}$ and Δ_F . Among the other three, the one involving both $S_{T\ell}$ and Δ_T contributes only to the absorptive part of the self-energy (see Eq. (4.5)), and does not concern us here. The contributions to the dispersive part of the self-energy (see Eq. (4.4)) arises from the remaining two terms, which can be written in the form

$$\Sigma'_\ell(p) = \Sigma'_{\ell 1}(p) + \Sigma'_{\ell 2}(p), \quad (4.94)$$

where

$$\Sigma'_{\ell 1}(p) = 2e^2 \int \frac{d^4k}{(2\pi)^3} \delta(k^2) \eta_\gamma(k) \frac{\not{p} + \not{k} - 2m_\ell}{p^2 + 2k \cdot p - m_\ell^2}, \quad (4.95)$$

$$\Sigma'_{\ell 2}(p) = -2e^2 \int \frac{d^4k}{(2\pi)^3} \delta(k^2 - m_\ell^2) \eta_\ell(k) \frac{\not{k} - 2m_\ell}{p^2 - 2k \cdot p + m_\ell^2}. \quad (4.96)$$

We shall now write

$$\Sigma'_{\ell 1}(p) = a_1(p) \not{p} + b_1(p) \not{p} + c_1(p) \quad (4.97)$$

and check the conditions of C invariance stated in Eq. (4.75). Eq. (4.97) can be inverted to express $a_1(p)$, $b_1(p)$ and $c_1(p)$ in terms of $\text{Tr}[\not{p}\Sigma'_{\ell 1}(p)]$, $\text{Tr}[\not{p}\Sigma'_{\ell 1}(p)]$ and $\text{Tr}[\Sigma'_{\ell 1}(p)]$. Then using the expression of Eq. (4.95), we obtain

$$\begin{aligned} a_1(p) &= \frac{2e^2}{(p \cdot v)^2 - p^2} \int \frac{d^4k}{(2\pi)^3} \delta(k^2) \eta_\gamma(k) \frac{(p \cdot v)(k \cdot v + p \cdot v) - (k \cdot p + p^2)}{p^2 + 2k \cdot p - m_\ell^2}, \\ b_1(p) &= \frac{2e^2}{(p \cdot v)^2 - p^2} \int \frac{d^4k}{(2\pi)^3} \delta(k^2) \eta_\gamma(k) \frac{(p \cdot v)(k \cdot p) - p^2 k \cdot v}{p^2 + 2k \cdot p - m_\ell^2}, \\ c_1(p) &= -4e^2 m_\ell \int \frac{d^4k}{(2\pi)^3} \delta(k^2) \eta_\gamma(k) \frac{1}{p^2 + 2k \cdot p - m_\ell^2}. \end{aligned} \quad (4.98)$$

Changing p to $-p$, and, simultaneously, k to $-k$, in each of these expressions, one can readily check the properties given by Eq. (4.75).

Let us next write

$$\Sigma'_{\ell 2}(p) = a_2(p)\not{p} + b_2(p)\not{p} + c_2(p). \quad (4.99)$$

We then have

$$\begin{aligned} a_2(p) &= \frac{2e^2}{(p \cdot v)^2 - p^2} \int \frac{d^4 k}{(2\pi)^3} \delta(k^2 - m_\ell^2) \eta_\ell(k) \frac{k \cdot p - (p \cdot v)(k \cdot v)}{p^2 - 2k \cdot p + m_\ell^2}, \\ b_2(p) &= \frac{2e^2}{(p \cdot v)^2 - p^2} \int \frac{d^4 k}{(2\pi)^3} \delta(k^2 - m_\ell^2) \eta_\ell(k) \frac{p^2 k \cdot v - (p \cdot v)(k \cdot p)}{p^2 - 2k \cdot p + m_\ell^2}, \\ c_2(p) &= 4e^2 m_\ell \int \frac{d^4 k}{(2\pi)^3} \delta(k^2 - m_\ell^2) \eta_\ell(k) \frac{1}{p^2 - 2k \cdot p + m_\ell^2}. \end{aligned} \quad (4.100)$$

To check the conditions of Eq. (4.85), we first have to change μ_ℓ to $-\mu_\ell$. Eq. (2.51) shows that this is equivalent to changing $\eta_\ell(k)$ to $\eta_\ell(-k)$. After making this change, one only has to change p to $-p$, and, simultaneously, k to $-k$, and the conditions check out.

4.5.2 Calculation of inertial mass

Using Eq. (4.22), and according to the decomposition given in Eq. (4.94), we write

$$\mathcal{E}_\ell = \mathcal{E}_{\ell 1} + \mathcal{E}_{\ell 2}, \quad (4.101)$$

where

$$\mathcal{E}_{\ell 1} = 2e^2 \int \frac{d^4 k}{(2\pi)^3} \delta(k^2) \eta_\gamma(k) \frac{p_0 + k_0 - 2m_\ell}{p^2 + 2k \cdot p - m_\ell^2}, \quad (4.102)$$

$$\mathcal{E}_{\ell 2} = -2e^2 \int \frac{d^4 k}{(2\pi)^3} \delta(k^2 - m_\ell^2) \eta_\ell(k) \frac{k_0 - 2m_\ell}{p^2 - 2k \cdot p + m_\ell^2}. \quad (4.103)$$

We can make a similar decomposition of $\mathcal{E}_{\bar{\ell}}$. The quantities $\mathcal{E}_{\bar{\ell} 1}$ and $\mathcal{E}_{\bar{\ell} 2}$ are obtained from $\mathcal{E}_{\ell 1}$ and $\mathcal{E}_{\ell 2}$ by replacing m_ℓ by $-m_\ell$.

The inertial mass is determined by applying Eq. (4.24), and, according to the decomposition given in Eq. (4.101), we write it as

$$M_\ell = m_\ell + m_{\ell 1} + m_{\ell 2}, \quad (4.104)$$

and similarly for the anti-leptons. Substituting $p^\mu = (m_\ell, \vec{0})$ in Eq. (4.102), and using the fact that the terms in the integrand that are odd in k yield zero, we obtain

$$\begin{aligned} m_{\ell 1} \equiv \mathcal{E}_{\ell 1}(m_\ell, \vec{0}) &= \frac{e^2}{m_\ell} \int \frac{d^4 k}{(2\pi)^3} \delta(k^2) \eta_\gamma(k) \\ &= \frac{e^2 T^2}{12m_\ell}. \end{aligned} \quad (4.105)$$

This is the contribution to the inertial mass from the photons in the background, in agreement with the result quoted in Eq. (1.1), and it is non-zero for any the charged lepton propagating through the medium. In a similar fashion we find

$$m_{\bar{\ell}1} \equiv -\mathcal{E}_{\bar{\ell}1}(-m_\ell, \vec{0}) = \frac{e^2 T^2}{12m_\ell}, \quad (4.106)$$

and therefore the photon contribution for the anti-particle is the same as for the corresponding particle.

In order to justify the usage of the term “inertial mass,” we now deduce the $O(e^2)$ dispersion relation when only photons are present in the background. For this purpose, Eq. (4.95) has to be used in Eq. (4.15). Since the tree-level dispersion relation $p^2 = \omega^2 - P^2 = m_\ell^2$ can be used in $O(e^2)$ terms, we obtain

$$\text{Tr}[(\not{p} + m_\ell)\Sigma'_{\ell 1}] = \frac{e^2 T^2}{3}, \quad (4.107)$$

after a term odd in the integration variable drops out, leading to the same integral as in Eq. (4.105). Consequently, Eq. (4.15) reads

$$\omega^2 = P^2 + m_\ell^2 + \frac{e^2 T^2}{6}, \quad (4.108)$$

and the dispersion relations are

$$E_\ell(P) = E_{\bar{\ell}}(P) = \sqrt{P^2 + m_{\ell T}^2} \quad (4.109)$$

with

$$m_{\ell T} = m_\ell + \frac{e^2 T^2}{12m_\ell}. \quad (4.110)$$

Thus the dispersion relations in photon background are of the same form as in vacuum, and reduce to $E_\ell = E_{\bar{\ell}} = m_{\ell T} + P^2/2m_{\ell T}$ in the non-relativistic limit. From this, it is clear that $m_{\ell T}$ is the inertial mass. Now, from Eq. (4.109), $m_{\ell T}$ also equals $E_\ell(0)$ or $E_{\bar{\ell}}(0)$, thereby justifying our definition of inertial mass given in Eq. (4.18). Note that Eq. (4.110) is in agreement with the results given in Eqs. (4.105) and (4.106).

When the medium also contains fermions, one finds that the dispersion relations no longer possess the form given in Eq. (4.109). This is because, unlike in Eq. (4.107), $\text{Tr}[(\not{p} + m_\ell)\Sigma'_{\ell 2}]$ is a function of momentum. We have chosen to call $E_\ell(0)$ and $E_{\bar{\ell}}(0)$ the inertial masses even in this case, by a generalization of the usual application of the term. Sometimes these quantities are called the phase-space masses [20].

In passing, we remark that the first equality in Eq. (4.109) is as expected from C invariance, which is present in the case of a photon background.

We now turn to the fermion background term given in Eq. (4.103), and find its effect on the inertial mass. In a background that contains electrons but not the other charged leptons, the distribution functions for the muon and the tau vanish. As a result,

$$m_{\mu 2} = m_{\tau 2} = m_{\bar{\mu} 2} = m_{\bar{\tau} 2} = 0. \quad (4.111)$$

For the electron, we obtain

$$m_{e2} \equiv \mathcal{E}_{e2}(m_e, \vec{0}) = \frac{e^2}{m_e} \int \frac{d^4 k}{(2\pi)^3} \delta(k^2 - m_e^2) \eta_e(k) \left[\frac{k_0 - 2m_e}{k_0 - m_e} \right] \quad (4.112)$$

Performing the integrations over k_0 and the angular variables, we obtain

$$m_{e2} = \frac{e^2}{2\pi^2 m_e} \int_0^\infty dK \frac{K^2}{2E_K} \left[\left(\frac{E_K - 2m_e}{E_K - m_e} \right) f_e(E_K) + \left(\frac{E_K + 2m_e}{E_K + m_e} \right) f_{\bar{e}}(E_K) \right]. \quad (4.113)$$

Here E_K is given by Eq. (2.6) (with m replaced by m_e) and the distribution functions by Eq. (2.43). Similarly,

$$\begin{aligned} m_{\bar{e}2} &\equiv -\mathcal{E}_{\bar{e}2}(-m_e, \vec{0}) \\ &= \frac{e^2}{2\pi^2 m_e} \int_0^\infty dK \frac{K^2}{2E_K} \left[\left(\frac{E_K + 2m_e}{E_K + m_e} \right) f_e(E_K) + \left(\frac{E_K - 2m_e}{E_K - m_e} \right) f_{\bar{e}}(E_K) \right]. \end{aligned} \quad (4.114)$$

The integration over K can be performed only when the distribution functions are specified, and we will consider some examples in Sec. 5.6. Here we only note the general result of Eq. (4.86), obtained by charge conjugation, is borne out at $P = 0$ by Eqs. (4.113) and (4.114): the inertial mass corrections for the particle and the antiparticle are the same only after we implement $\mu_e \rightarrow -\mu_e$, or equivalently,

$$f_e \leftrightarrow f_{\bar{e}}, \quad (4.115)$$

in one of them.

4.5.3 Calculation of wavefunction renormalization factor

We decompose

$$\zeta_\ell = \zeta_{\ell 1} + \zeta_{\ell 2} \quad (4.116)$$

with a similar decomposition for the anti-leptons, where, from Eq. (4.44),

$$\zeta_{\ell i} = \left. \frac{\partial \mathcal{E}_{\ell i}}{\partial p^0} \right|_{p^\mu = (m_\ell, \vec{0})} \quad \text{for } i = 1, 2, \quad (4.117)$$

and, from Eq. (4.57),

$$\zeta_{\bar{\ell} i} = \left. \frac{\partial \mathcal{E}_{\bar{\ell} i}}{\partial p^0} \right|_{p^\mu = (-m_\ell, \vec{0})} \quad \text{for } i = 1, 2. \quad (4.118)$$

From Eq. (4.102), we obtain

$$\left. \frac{\partial \mathcal{E}_{\ell 1}}{\partial p^0} \right|_{p^\mu = (m_\ell, \vec{0})} = \left. \frac{\partial \mathcal{E}_{\bar{\ell} 1}}{\partial p^0} \right|_{p^\mu = (-m_\ell, \vec{0})} = -\frac{e^2}{m_\ell^2} \int \frac{d^3 K}{(2\pi)^3} \frac{f_\gamma(K)}{K} \left(1 - \frac{m_\ell^2}{K^2} \right), \quad (4.119)$$

which implies

$$\zeta_{\ell 1} = \zeta_{\bar{\ell} 1} = -\frac{e^2 T^2}{12 m_\ell^2} + \frac{e^2}{2\pi^2} \int_0^\infty \frac{dK}{K} f_\gamma(K), \quad (4.120)$$

the photon distribution function being given by Eq. (2.56). Notice that the integral in Eq. (4.120) is infrared divergent. This divergence will turn out to be of importance in Chap. 5, since there it will cancel a similarly divergent term in the gravitational vertex contribution to the gravitational mass (see Eq. (5.59)).

Since the electron background terms do not contribute to the self-energy of the muon or the tau, it follows that

$$\zeta_{\mu 2} = \zeta_{\tau 2} = \zeta_{\bar{\mu} 2} = \zeta_{\bar{\tau} 2} = 0. \quad (4.121)$$

For the electron, Eq. (4.103) implies

$$\begin{aligned} \left. \frac{\partial \mathcal{E}_{e2}}{\partial p^0} \right|_{p^\mu = (m_e, \vec{0})} &= -\frac{\mathcal{E}_{e2}(m_e, \vec{0})}{m_e} \\ \left. \frac{\partial \mathcal{E}_{\bar{e}2}}{\partial p^0} \right|_{p^\mu = (-m_e, \vec{0})} &= \frac{\mathcal{E}_{\bar{e}2}(-m_e, \vec{0})}{m_e}, \end{aligned} \quad (4.122)$$

which yield

$$\begin{aligned} \zeta_{e2} &= -\frac{m_{e2}}{m_e} \\ \zeta_{\bar{e}2} &= -\frac{m_{\bar{e}2}}{m_e}, \end{aligned} \quad (4.123)$$

with m_{e2} and $m_{\bar{e}2}$ given in Eqs. (4.113) and (4.114) respectively. The result (4.120) shows that $Z_e = Z_{\bar{e}}$ in a photon background, while the result (4.123) shows that Z_e transforms to $Z_{\bar{e}}$ under $f_e \leftrightarrow f_{\bar{e}}$ (so that m_{e2} becomes $m_{\bar{e}2}$) as expected from considerations of charge conjugation.

CHAPTER 5

Gravitational couplings of charged leptons in a medium

5.1 Introduction

The underlying Lagrangian for the calculations of this chapter is that of the weak gravitational field version of QED in curved space, which was discussed in Chapter 3. Thus we shall use the graviton couplings of Sec. 3.5, in addition to, of course, the flat space QED coupling. Since the coupling constant κ is of mass dimension -1 , the theory we are considering is non-renormalizable. But this will not be a problem, since our calculations will be confined to the lowest order in κ , and the radiative corrections to be considered will involve an expansion in the electromagnetic coupling constant e . All our calculations will be valid at energies low compared to $\kappa^{-1} \approx m_{\text{Planck}}$, as stated in Sec. 3.1. (This is comparable, to some extent, to tree-level calculations using the four-Fermi interaction, which are valid at energies much less than m_W .)

5.2 Gravitational mass

The gravitational mass is a measure of the strength of the coupling of the fermion to the graviton. It can be determined in terms of the fermion's vertex function for the gravitational interaction. For this purpose, we are going to consider the S -matrix element of quantum field theory for the case of scattering of a slowly moving fermion, by a weak and static external gravitational field (these conditions define the Newtonian limit of gravitational interaction). Since our aim is to calculate gravitational mass in a medium, we must consider this scattering to take place in the presence of a thermal background.

So, let us first consider the relevant part

$$\mathcal{L}_{\text{eff}} = -\kappa h^{\lambda\rho}(x) \hat{T}_{\lambda\rho}(x) \tag{5.1}$$

of the interaction Lagrangian. Here $\hat{T}_{\lambda\rho}(x)$ is the total energy-momentum tensor operator, involving quantized fermion and photon fields, such that $\mathcal{L}_{\text{eff}}$ represents the possible interactions of these fields with the classical field $h_{\lambda\rho}(x)$ to all orders in e . The S -matrix, to lowest order in κ , is then

$$S = 1 - i\kappa \int d^4x h^{\lambda\rho}(x) \hat{T}_{\lambda\rho}(x). \quad (5.2)$$

Eq. (5.2) follows from the well-known S -matrix expansion

$$S = T[\exp(i \int d^4x \mathcal{L}_{\text{int}})] \quad (5.3)$$

where T denotes time-ordering. The standard derivation of this formula assumes that the interaction $\mathcal{L}_{\text{int}}$ does not contain derivatives of fields, so that the commutation relations of the interacting theory are the same as those in the free-field theory, and also $\mathcal{H}_{\text{int}} = -\mathcal{L}_{\text{int}}$. In the present case, $\mathcal{L}_{\text{eff}}$ does contain derivatives through $\hat{T}_{\lambda\rho}$ (see, for example, Eqs. (3.78) and (3.93)). However the formula (5.3) can be shown to be generally valid, irrespective of $\mathcal{L}_{\text{int}}$ containing derivatives or not, provided one uses the commutation relations of the free theory in calculating the S -matrix elements [49]. This justifies our use of Eq. (5.3), and also the derivation of the interaction vertices in Secs. 3.5.1 and 3.5.2.

Returning to Eq. (5.2), we now consider the S -matrix element between incoming and outgoing fermion states:

$$S_{ff} \equiv \langle f(\vec{P}', s) | S | f(\vec{P}, s) \rangle \quad (5.4)$$

$$= -i\kappa \langle f(\vec{P}', s) | \hat{T}_{\lambda\rho}(0) | f(\vec{P}, s) \rangle \int d^4x e^{-iq \cdot x} h^{\lambda\rho}(x), \quad (5.5)$$

where we have used the relation

$$\hat{T}_{\lambda\rho}(x) = e^{i\hat{P} \cdot x} \hat{T}_{\lambda\rho}(0) e^{-i\hat{P} \cdot x}, \quad (5.6)$$

$\hat{P}_\mu$ being the momentum operator, and also defined

$$q = p - p'. \quad (5.7)$$

Here $q_0 = E_f - E'_f$, each of the energies satisfying the dispersion relation of charged leptons in thermal QED.

At zeroth order in e , the matrix element of Eq. (5.5) equals $\bar{u}_s(p') V_{\lambda\rho}(p, p') u_s(p)$ with $V_{\lambda\rho}(p, p')$ given by Eq. (3.95). When higher orders in e are taken into account, we write

$$\langle f(\vec{P}', s) | \hat{T}_{\lambda\rho}(0) | f(\vec{P}, s) \rangle = \sqrt{Z_f(P) Z_f(P')} \bar{U}_s(\vec{P}') \Gamma_{\lambda\rho}(p, p') U_s(\vec{P}) \quad (5.8)$$

following the LSZ reduction formula of vacuum field theory. Here $\Gamma_{\lambda\rho}(p, p')$ is the 1PI vertex function: it is the sum of all connected, amputated diagrams contributing to the fermion-fermion-graviton vertex. Though the vertex function can be defined in general for off-shell momenta (see Eq. (C.11)), the arguments of $\Gamma_{\lambda\rho}$ in Eq. (5.8) are on-shell momenta, and satisfy the dispersion relation of the charged lepton. It may be noted that the replacement of the tree-level spinor $u_s(p)$ with $\sqrt{Z_f(P)}U_s(\vec{P})$ in going beyond the lowest order, as done in Eq. (5.8), also gives the residue of the one-particle pole of the dressed propagator in Eq. (4.38) [50].

Using Eq. (5.8) in Eq. (5.5), and also taking $h^{\lambda\rho}$ to be independent of time, we obtain

$$S_{ff} = -i\kappa(2\pi)\delta(E_f - E'_f)Z_f(P)\overline{U}_s(\vec{P}')\Gamma_{\lambda\rho}(p, p')U_s(\vec{P})h^{\lambda\rho}(\vec{Q}). \quad (5.9)$$

Here we have denoted the three-vector part of q^μ by upper case letter, and also defined the Fourier transform by

$$f(\vec{x}) = \int \frac{d^3Q}{(2\pi)^3} f(\vec{Q}) e^{-i\vec{Q}\cdot\vec{x}} \quad (5.10)$$

for any function $f(\vec{x})$. Note that $E_f = E'_f$ implies that $P = P'$, so that the wavefunction renormalization factors for the initial and the final states are the same.

Now we relate $h^{\lambda\rho}$ to an external gravitational potential ϕ^{ext} . First consider Poisson's equation

$$\nabla^2 \phi^{\text{ext}}(\vec{x}) = 4\pi G \rho^{\text{ext}}(\vec{x}), \quad (5.11)$$

the static mass density $\rho^{\text{ext}}(\vec{x})$ being the source of the gravitational potential. In momentum space, this becomes

$$-2Q^2 \phi^{\text{ext}}(\vec{Q}) = \kappa^2 \rho^{\text{ext}}(\vec{Q}). \quad (5.12)$$

Next, the energy-momentum tensor corresponding to ρ^{ext} is

$$T_{\lambda\rho} = v_\lambda v_\rho \rho^{\text{ext}} \quad (5.13)$$

where v_λ is the four-velocity of the mass density. This $T_{\lambda\rho}$ determines the desired $h_{\lambda\rho}$ through the linearized Einstein field equations, which, in the harmonic gauge, read

$$\square h_{\lambda\rho} = -\kappa(T_{\lambda\rho} - \frac{1}{2}\eta_{\lambda\rho}T^\mu{}_\mu). \quad (5.14)$$

(To obtain Eq. (5.14), one starts with the field equations in the form

$$R_{\lambda\rho} = -8\pi G(T_{\lambda\rho} - \frac{1}{2}g_{\lambda\rho}T^\mu{}_\mu), \quad (5.15)$$

and evaluates each side to the lowest order in κ . Thus, one puts $R_{\lambda\rho}$ correct to $O(\kappa)$ (from Eqs. (3.7) and (3.13)) on the L.H.S. Since the R.H.S. is already of $O(\kappa^2)$, one can put $g_{\lambda\rho} = \eta_{\lambda\rho}$ on this side. Then the gauge condition (3.23) is used.) After noting that $\square$ becomes $-\nabla^2$ for the static case, we write Eq. (5.14) in momentum space, and use first Eq. (5.13) and then Eq. (5.12) in it. In this way, we finally arrive at

$$\kappa h^{\lambda\rho}(\vec{Q}) = (2v^\lambda v^\rho - \eta^{\lambda\rho})\phi^{\text{ext}}(\vec{Q}) \quad (5.16)$$

It may be noted that in the frame in which the mass distribution is at rest, i.e. for $v^\lambda = (1, \vec{0})$, Eq. (5.16) immediately leads to $g_{00} = 1 + 2\phi^{\text{ext}}$, which can also be deduced by demanding that the geodesic equation reduces to Newton's equation of motion in the appropriate limit [51]. Another important point is that we shall take v^λ equal to the four-velocity of the medium, which occurs in the thermal propagators. This means that we shall take the medium to be at rest relative to the mass distribution producing the external gravitational field (a special case being that the medium itself is the source of the gravitational field). Eqs. (5.9) and (5.16) constitute our end-result for the S -matrix element.

We now define the gravitational mass of the fermion. The mass density operator for the fermion, $\hat{\rho}_f(t, \vec{x})$, is determined by writing another effective Lagrangian

$$\mathcal{L}'_{\text{eff}} = -\hat{\rho}_f(t, \vec{x})\phi^{\text{ext}}(\vec{x}) \quad (5.17)$$

such that it reproduces S_{ff} as given by Eqs. (5.9) and (5.16). The gravitational mass M'_f is then defined through the matrix element of the mass operator (obtained by integrating $\hat{\rho}_f$ at any particular instant t_ρ over all space) in the zero-momentum limit as follows:

$$\lim_{\vec{P} \rightarrow 0} \langle f(\vec{P}', s) | \int d^3x \hat{\rho}_f(t_\rho, \vec{x}) | f(\vec{P}, s) \rangle = (2\pi)^3 [\delta^{(3)}(\vec{P} - \vec{P}')]_{\vec{P} \rightarrow 0} M'_f. \quad (5.18)$$

A number of remarks on this definition are in order. Firstly, the normalization of one-particle states adopted by us, as given by Eq. (B.8), is consistent with the definition given in Eq. (5.18). Secondly, the mass operator is independent of $\vec{x}$ (as it involves integration over all space) and so carries zero momentum. Consequently, on the L.H.S. of Eq. (5.18) we have $P = \vec{P}'$. The delta function on the R.H.S. embodies this fact (see also the

derivation of Eq. (5.21) below.) Thirdly, we shall see that M'_f is independent of the choice of t_ρ .

The remaining task is to find an expression for the gravitational mass. Eqs. (5.3), (5.4) and (5.17) give, to lowest order in interaction,

$$S_{ff} = -i2\pi\delta(E'_f - E_f)\phi^{\text{ext}}(\vec{P} - \vec{P}')\langle f(\vec{P}', s)|\hat{\rho}_f(t_\rho, \vec{0})|f(\vec{P}, s)\rangle, \quad (5.19)$$

where we used

$$\hat{\rho}_f(t, \vec{x}) = e^{i\hat{H}(t-t_\rho)-i\hat{\vec{P}}\cdot\vec{x}}\hat{\rho}_f(t_\rho, \vec{0})e^{-i\hat{H}(t-t_\rho)+i\hat{\vec{P}}\cdot\vec{x}}, \quad (5.20)$$

and the definition of Fourier transform given in Eq. (5.10). Again Eq. (5.18) gives (on using Eq. (5.20) with $t = t_\rho$)

$$M'_f = \lim_{\vec{P} \rightarrow 0} \left[\langle f(\vec{P}', s)|\hat{\rho}_f(t_\rho, \vec{0})|f(\vec{P}, s)\rangle \right]_{\vec{P}' = \vec{P}}. \quad (5.21)$$

Now $\vec{P}' = \vec{P}$ implies that $E'_f = E_f$. So the matrix element occurring on the R.H.S. of Eq. (5.21) can be determined by using Eq. (5.19) on one hand, and Eqs. (5.9) and (5.16) on the other hand. We are thus led to

$$M'_f = (2v^\lambda v^\rho - \eta^{\lambda\rho}) \lim_{\vec{P} \rightarrow 0} Z_f(P) \left\{ \overline{U}_s(\vec{P}') \Gamma_{\lambda\rho}(p, p') U_s(\vec{P}) \right\}_{\substack{E'_f = E_f \\ \vec{P}' \rightarrow P}}. \quad (5.22)$$

Note that in the last step we have relaxed the equality $\vec{P}' = \vec{P}$ to a limiting procedure, while retaining the weaker condition $E'_f = E_f$. The significance of this exercise will be explained in detail at the end of Sec. 5.3.

We end this section with the comment that our aim was only to get the expression for the gravitational mass in a medium [52]. Calculation of scattering cross-section, which also involves modification of phase space factors at finite temperature, does not concern us here.

5.3 Operational definition at $O(e^2)$

We write the complete 1PI vertex function at one-loop in the form

$$\Gamma_{\lambda\rho} = V_{\lambda\rho} + \Gamma'_{\lambda\rho}. \quad (5.23)$$

Since $V_{\lambda\rho}$ denotes the tree-level vertex function, $\Gamma'_{\lambda\rho}$ is the $O(e^2)$ contribution. Then, using Eqs. (5.23) and (4.43), the formula given by Eq. (5.22) can be rewritten at $O(e^2)$

in the form

$$M'_f = (2v^\lambda v^\rho - \eta^{\lambda\rho}) \times \lim_{\vec{P} \rightarrow 0} \left\{ \bar{U}_s(\vec{P}) V_{\lambda\rho}(p, p) U_s(\vec{P}) + \zeta_f \bar{u}_s(\vec{P}) V_{\lambda\rho}(p, p) u_s(\vec{P}) \right. \\ \left. + \left[\bar{u}_s(\vec{P}') \Gamma'_{\lambda\rho}(p, p') u_s(\vec{P}) \right]_{\substack{E'_f = E_f \\ \vec{P}' \rightarrow P}} \right\}. \quad (5.24)$$

Since ζ_f and $\Gamma'_{\lambda\rho}$ are $O(e^2)$, in any term that contains either of these factors we have substituted the tree-level expressions for the other quantities. Now, the terms involving $V_{\lambda\rho}$ can be evaluated immediately with the help of the identities given in Eqs. (4.32) and (4.33). Remembering that $E_f(0) = M_f$, we finally obtain the operational definition to $O(e^2)$

$$M'_f = 3M_f - 2m_f + \zeta_f m_f + (2v^\lambda v^\rho - \eta^{\lambda\rho}) \lim_{\vec{P} \rightarrow 0} \left\{ \left[\bar{u}_s(\vec{P}') \Gamma'_{\lambda\rho}(p, p') u_s(\vec{P}) \right]_{\substack{E'_f = E_f \\ \vec{P}' \rightarrow P}} \right\}, \quad (5.25)$$

where we can set $E_f = \sqrt{P^2 + m_f^2}$ in the last term.

We now turn to the derivation of an operational definition at $O(e^2)$ for the gravitational mass of the antilepton. We first aim to write down an expression for $\langle \bar{f}(\vec{P}, s) | S | \bar{f}(\vec{P}', s) \rangle$ from Eq. (5.9) by crossing. To this end, let us note the action of the field operators ψ and $\bar{\psi}$, contained, through $\mathcal{L}_{\text{eff}}$, in S . In the S -matrix element of Eq. (5.9), ψ kills $f(\vec{P}, s)$ and gives $\sqrt{Z_f(P)} U_s(\vec{P}) e^{-ip \cdot x}$, while in $\langle \bar{f}(\vec{P}, s) | S | \bar{f}(\vec{P}', s) \rangle$, ψ creates $\bar{f}(\vec{P}, s)$ and gives $\sqrt{Z_{\bar{f}}(P)} V_s(\vec{P}) e^{ip \cdot x}$ (see the tree-level Fourier expansion for ψ given in Eq. (2.37)). This does not only identify the external leg factors, but also shows that the incoming momentum p in the amputated vertex corresponding to the first case gets replaced in the second case with an outgoing momentum p , equivalent to an incoming momentum $-p$. Similarly, one can go through the action of $\bar{\psi}$. Note also that the graviton comes out with three-momentum $\vec{P}' - \vec{P}$ in $\langle \bar{f}(\vec{P}, s) | S | \bar{f}(\vec{P}', s) \rangle$, while the fermion exchange rule gives an extra minus sign. We therefore have

$$\langle \bar{f}(\vec{P}, s) | S | \bar{f}(\vec{P}', s) \rangle = (-1)(-i\kappa) 2\pi \delta(E_{\bar{f}} - E'_{\bar{f}}) \\ \times Z_{\bar{f}}(P) \bar{V}_s(\vec{P}') \Gamma_{\lambda\rho}(-p, -p') V_s(\vec{P}) h^{\lambda\rho}(\vec{P}' - \vec{P}). \quad (5.26)$$

Note that p and p' , of which $\Gamma_{\lambda\rho}$ is a function of in Eq. (5.26), are on-shell, and satisfy the antifermion dispersion relation.

Next, we simply interchange $\vec{P}$ and $\vec{P}'$ in Eq. (5.26) to obtain the antiparticle equation corresponding to Eq. (5.9):

$$\begin{aligned} S_{\bar{f}\bar{f}} &\equiv \langle \bar{f}(\vec{P}', s) | S | \bar{f}(\vec{P}, s) \rangle \\ &= (-1)(-i\kappa)2\pi\delta(E_{\bar{f}} - E'_{\bar{f}})Z_{\bar{f}}(P) \bar{V}_s(\vec{P})\Gamma_{\lambda\rho}(-p', -p)V_s(\vec{P}')h^{\lambda\rho}(\vec{P} - \vec{P}') \end{aligned} \quad (5.27)$$

This leads to an equation that is analogous to Eq. (5.24), but with an extra minus sign in front and some obvious changes in the corresponding symbols, which in turn lead to the $O(e^2)$ formula

$$\begin{aligned} M'_{\bar{f}} &= 3M_{\bar{f}} - 2m_f + \zeta_{\bar{f}}m_f \\ &\quad - (2v^\lambda v^\rho - \eta^{\lambda\rho}) \lim_{\vec{P} \rightarrow 0} \left\{ \left[\bar{v}_s(\vec{P})\Gamma'_{\lambda\rho}(-p', -p)v_s(\vec{P}') \right]_{\substack{E'_{\bar{f}}=E_{\bar{f}} \\ \vec{P}' \rightarrow P}} \right\}. \end{aligned} \quad (5.28)$$

Using the usual relation between the free particle and antiparticle spinors given in Eq. (4.77), and the properties of the charge conjugation matrix $\mathbf{C}$ given by Eqs. (4.61), (4.62) and (4.63), the spinor matrix element that appears in Eq. (5.28) can be rewritten in the form

$$\bar{v}_s(\vec{P})\Gamma'_{\lambda\rho}(-p', -p)v_s(\vec{P}') = -\bar{u}_s(\vec{P}')\mathbf{C}\Gamma_{\lambda\rho}^T(-p', -p)\mathbf{C}^{-1}u_s(\vec{P}). \quad (5.29)$$

Finally, using Eq. (C.20) (which actually holds at all orders and even for off-shell momenta) in Eq. (5.29), and substituting the relation in Eq. (5.28), we arrive at

$$\begin{aligned} M'_{\bar{f}} &= 3M_{\bar{f}} - 2m_f + \zeta_{\bar{f}}m_f \\ &\quad + (2v^\lambda v^\rho - \eta^{\lambda\rho}) \lim_{\vec{P} \rightarrow 0} \left\{ \left[\left[\bar{u}_s(\vec{P}')\Gamma'_{\lambda\rho}(p, p')u_s(\vec{P}) \right]_{\mu \rightarrow -\mu} \right]_{\substack{E'_f=E_f \\ \vec{P}' \rightarrow P}} \right\}. \end{aligned} \quad (5.30)$$

From Eqs. (5.25) and (5.30), it follows that $M'_{\bar{f}}$ and M'_f are related by

$$[M'_f]_{\mu \rightarrow -\mu} = M'_{\bar{f}}. \quad (5.31)$$

Since the gravitational mass of the antiparticle should equal that of the particle for C symmetric Lagrangian and medium, Eq. (5.31) is expected from the discussion of Sec. 4.4 for C symmetric Lagrangian and C violating background. The consequence of Eq. (5.31) is that we can just obtain $M'_{\bar{f}}$ from M'_f by implementing the prescription of Eq. (4.115) in the $O(e^2)$ correction.

We take the opportunity to emphasize the following point. In the calculations that follow, we will find expressions for the various contributions to $\Gamma'_{\lambda\rho}(p, p')$, which are given as integrals over the propagators and thermal distribution functions. In general, such expressions do not have a unique limiting value as we let $p' \rightarrow p$ in an arbitrary way [53]. More importantly, some of the integrals involve ill-defined expressions if the limit is not taken properly. In our case, the precise order in which the various limits must be taken has been dictated by the physical issue at hand. Thus, since we are interested in the interaction of the particle with a static gravitational potential, the quantity that enters is $\Gamma'_{\lambda\rho}(p, p')$ evaluated for $E'_f = E_f$. Next we set $\vec{P}' = \vec{P}$ since we want the gravitational field to vary slowly over a macroscopic region. Finally we set $\vec{P} \rightarrow 0$ to obtain the coupling at zero momentum, which determines the gravitational mass. This justifies the somewhat cumbersome notation regarding the limits in Eq. (5.25), but it is meant to indicate precisely what we have just explained. We will see that this prescription allowed us to evaluate all the integrals involved (including those that superficially seem to be ill-defined) in a unique and well-defined way, without having to introduce any special regularization technique. Our result for the case of the photon background agrees with the result of DHR [20], who used a special regularization technique, while our result for the case of the electron background agrees with the answer obtained by the use of a regularization technique similar to that of DHR (see Appendix E.4).

5.4 Gravitational vertex

5.4.1 Irreducible diagrams

The irreducible one-loop diagrams for the vertex function are given in Figs. 5.1 and 5.2. The four-vector q^μ , which has been defined in Eq. (5.7), gives the momentum of the outgoing graviton.

We calculate only the terms that contribute to the dispersive part of the vertex function. To determine the condition for this, let us return to Eq. (5.1), and demand that $\mathcal{L}_{\text{eff}}$ be hermitian. This means $\hat{T}_{\lambda\rho}^\dagger(x) = \hat{T}_{\lambda\rho}(x)$. Next consider Eq. (5.8). Exchange of p and p' in this equation, followed by complex conjugation, gives

$$\langle f(\vec{P}, s) | \hat{T}_{\lambda\rho}(0) | f(\vec{P}', s) \rangle^* = \sqrt{Z_f(P)Z_f(P')} \bar{U}_s(\vec{P}') \gamma_0 \Gamma_{\lambda\rho}^\dagger(p', p) \gamma_0 U_s(\vec{P}). \quad (5.32)$$




Figure 5.1: Bubble diagrams for the one-loop gravitational vertex of charged leptons. These diagrams contribute in a background of photons and electrons. The braided line represents the graviton.




Figure 5.2: Tadpole diagrams for the one-loop gravitational vertex of charged leptons. These diagrams contribute to the vertex of any charged lepton in a background of electrons and nucleons.

But since $\hat{T}_{\lambda\rho}$ is a hermitian operator, the left hand sides of Eqs. (5.8) and (5.32) are equal. So equating the right hand sides of these equations, we are led to the desired condition

$$\Gamma_{\lambda\rho}(p, p') = \gamma_0 \Gamma_{\lambda\rho}^\dagger(p', p) \gamma_0 \quad (5.33)$$

for the dispersive part of the vertex function. The absorptive part, on the other hand, corresponds to antihermitian $\mathcal{L}_{\text{eff}}$ and satisfies $\Gamma_{\lambda\rho}(p, p') = -\gamma_0 \Gamma_{\lambda\rho}^\dagger(p', p) \gamma_0$. It contributes to fermion damping, with which we are not concerned in the present work.

When the formulas given in Eqs. (2.48) and (2.52) for the propagators are substituted in the expressions corresponding to the diagrams, we obtain terms of different kind. One of them is independent of the background medium, in which we are not interested. Those involving two factors of the thermal part of the propagators contribute to the absorptive part of the vertex, while those involving three factors of the thermal part vanish because of the various δ -functions appearing in it. Thus, the background induced contribution to the dispersive part of the vertex contains the thermal part of only one of the propagators, and they are the only kind of term that we retain.

We have omitted the one-particle reducible diagrams in which the graviton line comes out from one of the external fermion legs, because they do not contribute to $\Gamma_{\lambda\rho}$. The proper way to take them into account in the calculation of the amplitude for any given process, is to choose the external spinor to be the solution of the effective Dirac equation for the propagating fermion mode in the medium (instead of the spinor representing the free-particle solution of the equation in vacuum), and multiply it with the square-root of the wavefunction renormalization factor, as discussed in Sec. 5.2.

5.4.2 Bubble diagrams given in Fig. 5.1

Diagram 5.1A

The amplitude of the diagram in Fig. 5.1A can be written as

$$-i\kappa\Gamma_{\lambda\rho}^{(A)}(p, p') = \int \frac{d^4k}{(2\pi)^4} ie\gamma_\alpha iS_\ell(k') (-i\kappa)V_{\lambda\rho}(k, k') iS_\ell(k) ie\gamma_\beta iD^{\alpha\beta}(k-p) \quad (5.34)$$

where

$$k' \equiv k - q. \quad (5.35)$$

As already explained, to determine the contribution to the dispersive part of the vertex function we need to retain the terms that contain the thermal part of only one of the

propagators. Any of them contains some combination of the form

$$\Lambda_{\lambda\rho}(k_1, k_2) \equiv \gamma_\alpha(\not{k}_2 + m_\ell)V_{\lambda\rho}(k_1, k_2)(\not{k}_1 + m_\ell)\gamma^\alpha. \quad (5.36)$$

After some straightforward algebra, this can be written as

$$\begin{aligned} \Lambda_{\lambda\rho}(k_1, k_2) = & -\frac{1}{2} \left[(k_1 + k_2)_\rho (\not{k}_1 \gamma_\lambda \not{k}_2 + m_\ell^2 \gamma_\lambda) + (k_1 + k_2)_\lambda (\not{k}_1 \gamma_\rho \not{k}_2 + m_\ell^2 \gamma_\rho) \right] \\ & + \eta_{\lambda\rho} \left[(k_1^2 - m_\ell^2)(\not{k}_2 - 2m_\ell) + (k_2^2 - m_\ell^2)(\not{k}_1 - 2m_\ell) \right] \\ & + 2m_\ell (k_1 + k_2)_\lambda (k_1 + k_2)_\rho. \end{aligned} \quad (5.37)$$

For the sake of convenience, we divide the total contribution into two parts

$$\Gamma_{\lambda\rho}^{(A)}(p, p') = \Gamma_{\lambda\rho}^{\prime(A1)}(p, p') + \Gamma_{\lambda\rho}^{\prime(A2)}(p, p'), \quad (5.38)$$

where $\Gamma_{\lambda\rho}^{\prime(A1)}$ contains the distribution function of the photon and therefore contributes to the gravitational vertex for all charged leptons, and $\Gamma_{\lambda\rho}^{\prime(A2)}$ contains the distribution function of the electrons and contributes only to the vertex for the electrons. Changing the integration variable from k to $k + p$, we obtain

$$\Gamma_{\lambda\rho}^{\prime(A1)}(p, p') = -e^2 \int \frac{d^4 k}{(2\pi)^3} \frac{\delta(k^2) \eta_\gamma(k)}{[(k + p')^2 - m_\ell^2][(k + p)^2 - m_\ell^2]} \Lambda_{\lambda\rho}(k + p, k + p') \quad (5.39)$$

and similarly,

$$\begin{aligned} \Gamma_{\lambda\rho}^{\prime(A2)}(p, p') = & e^2 \int \frac{d^4 k}{(2\pi)^3} \delta(k^2 - m_\ell^2) \eta_\ell(k) \\ & \times \left(\frac{\Lambda_{\lambda\rho}(k, k - q)}{[(k - q)^2 - m_\ell^2](k - p)^2} + \frac{\Lambda_{\lambda\rho}(k + q, k)}{[(k + q)^2 - m_\ell^2](k - p')^2} \right). \end{aligned} \quad (5.40)$$

Diagram 5.1B

For this diagram

$$-i\kappa \Gamma_{\lambda\rho}^{(B)}(p, p') = \int \frac{d^4 k}{(2\pi)^4} i e \gamma^\alpha i S_\ell(p - k) i e \gamma^\beta (-i\kappa) C_{\mu\nu\lambda\rho}(k, k') i D^{\nu\alpha}(k) i D^{\mu\beta}(k'), \quad (5.41)$$

and we decompose it in analogy with Eq. (5.38). The part that contains the photon distribution function is

$$\Gamma_{\lambda\rho}^{\prime(B1)}(p, p') = e^2 \int \frac{d^4 k}{(2\pi)^4} \gamma^\nu S_{F\ell}(p - k) \gamma^\mu C_{\mu\nu\lambda\rho}(k, k') [\Delta_F(k) \Delta_T(k') + \Delta_F(k') \Delta_T(k)] \quad (5.42)$$

Making a change of the integration variable in one of the terms, this can be written as

$$\begin{aligned}\Gamma_{\lambda\rho}'^{(B1)}(p, p') &= e^2 \int \frac{d^4 k}{(2\pi)^3} \delta(k^2) \eta_\gamma(k) \left[\frac{\gamma^\nu(\not{p}' - \not{k} + m_\ell) \gamma^\mu C_{\mu\nu\lambda\rho}(k + q, k)}{[(p' - k)^2 - m_\ell^2](k + q)^2} \right. \\ &\quad \left. + \frac{\gamma^\nu(\not{p} - \not{k} + m_\ell) \gamma^\mu C_{\mu\nu\lambda\rho}(k, k - q)}{[(p - k)^2 - m_\ell^2](k - q)^2} \right],\end{aligned}\quad (5.43)$$

while

$$\begin{aligned}\Gamma_{\lambda\rho}'^{(B2)}(p, p') &= e^2 \int \frac{d^4 k}{(2\pi)^4} \gamma^\nu S_{T\ell}(k) \gamma^\mu C_{\mu\nu\lambda\rho}(p - k, p' - k) \Delta_F(p - k) \Delta_F(p' - k) \\ &= -e^2 \int \frac{d^4 k}{(2\pi)^3} \delta(k^2 - m_\ell^2) \eta_\ell(k) \\ &\quad \times \gamma^\nu (\not{k} + m_\ell) \gamma^\mu \frac{C_{\mu\nu\lambda\rho}(p - k, p' - k)}{(p - k)^2 (p' - k)^2}\end{aligned}\quad (5.44)$$

gives the lepton background part.

Diagrams 5.1C and 5.1D

For these two diagrams the manipulations are similar and, omitting the details, the results are

$$\begin{aligned}\Gamma_{\lambda\rho}'^{(C1+D1)}(p, p') &= -e^2 a_{\mu\nu\lambda\rho} \int \frac{d^4 k}{(2\pi)^3} \delta(k^2) \eta_\gamma(k) \\ &\quad \times \left[\frac{\gamma^\mu (\not{k} + \not{p}' + m_\ell) \gamma^\nu}{(k + p')^2 - m_\ell^2} + \frac{\gamma^\nu (\not{k} + \not{p} + m_\ell) \gamma^\mu}{(k + p)^2 - m_\ell^2} \right],\end{aligned}\quad (5.45)$$

and

$$\begin{aligned}\Gamma_{\lambda\rho}'^{(C2+D2)}(p, p') &= e^2 a_{\mu\nu\lambda\rho} \int \frac{d^4 k}{(2\pi)^3} \delta(k^2 - m_\ell^2) \eta_\ell(k) \\ &\quad \times \left[\frac{\gamma^\mu (\not{k} + m_\ell) \gamma^\nu}{(k - p')^2} + \frac{\gamma^\nu (\not{k} + m_\ell) \gamma^\mu}{(k - p)^2} \right].\end{aligned}\quad (5.46)$$

5.4.3 Tadpole diagrams given in Fig. 5.2

The question of the photon tadpole

The calculation of the diagrams of Fig. 5.2 leads to an immediate problem. While the contributions from the diagrams 5.2B, 5.2C and 5.2D appear to vanish due to the charge neutrality of the medium (see Sec. 4.5), the contribution from the diagram 5.2A appears to diverge for $q = 0$ (due to the divergence of the photon propagator in this limit). The following discussion is devoted to the setting up of a framework of calculation which leads to the disappearance of the aforementioned divergence.




Figure 5.3: The one-loop diagrams that contribute to order κ to the photon tadpole in a medium, in the presence of a static and homogeneous ($q = 0$) gravitational potential. The fermion loop involves a sum over all the species of fermions present in the medium.

Let us recall from the discussion of Sec. 4.5 that the electrical neutrality of the medium requires the vanishing of the photon tadpole. The one-loop diagrams that contribute to the photon tadpole *in the presence of a static and homogeneous gravitational potential* are shown in Fig. 5.3, where the graviton line represents the $q = 0$ background field. In the absence of the background field, only the diagram 5.3A contributes to the photon tadpole. In that case, the requirement that the tadpole vanishes yields the familiar condition (see Eq. (4.91))

$$\mathcal{Q}^{(5.3A)} \equiv \sum_f \mathcal{Q}_f \left[2 \int \frac{d^3 K}{(2\pi)^3} [f_f(E_K) - f_{\bar{f}}(E_K)] \right] = 0. \quad (5.47)$$

and the quantity $\mathcal{Q}^{(5.3A)}$ is identified with the total charge of the medium. However, in the presence of the background field, and to the order that we are calculating, we have to take into account the contributions of the diagrams 5.3B and 5.3C to the photon tadpole or, equivalently, to the total charge of the system. If we denote them by $\mathcal{Q}^{(5.3B)}$ and $\mathcal{Q}^{(5.3C)}$ respectively, the condition for the vanishing of the photon tadpole is

$$\mathcal{Q}^{(5.3A)} + \mathcal{Q}^{(5.3B)} + \mathcal{Q}^{(5.3C)} = 0, \quad (5.48)$$

instead of Eq. (5.47). Physically, this means that the number density of the particles are not determined by their free distribution functions. The particle distributions rearrange themselves in a way that depends on the background gravitational field.

Consider now the problem of finding out the dispersion relation of the fermion in an external gravitational field. The dispersion relation is determined by the poles of the function given in Eq. (4.1), with $\Sigma_f(p)$ now denoting the fermion self-energy *in an external gravitational field*. Thus, the one-loop diagrams contributing to $\Sigma_f(p)$ at $O(\kappa)$ are the diagrams of Fig. 4.1, Fig. 5.1 and Fig. 5.2. Let us confine our attention to the tadpole diagrams, i.e., the diagram of Fig. 4.1B and the diagrams of Fig. 5.2, and consider the implications of the charge neutrality condition of Eq. (5.48) for the sum of the contributions arising from these diagrams.

Firstly, the unadorned tadpole of Fig. 5.3A is now itself of order κ because of the condition (5.48). Since the diagrams 5.2C and 5.2D contain an explicit factor of κ apart from the unadorned tadpole, their contribution is actually of order κ^2 and therefore we can neglect them. Secondly, the diagram 4.1B cancels the q -independent contributions from the diagrams 5.2A and 5.2B. Since the loop in diagram 5.2B is independent of q , this diagram is totally cancelled.

In summary, the only contribution from the various tadpole diagrams arises from the q -dependent part of the tadpole subdiagram of Fig. 5.2A (by subdiagram, we mean that the lower part, beginning with the photon propagator, is amputated). It seems reasonable to assume that this holds true also when we consider the gravitational vertex function and its contribution to the gravitational mass. It then turns out that the q -dependent part of the tadpole subdiagram of Fig. 5.2A, when multiplied with $2v^\lambda v^\rho - \eta^{\lambda\rho}$, and evaluated at $q_0 = 0$ and $\vec{Q} \rightarrow 0$, is proportional to $v_\alpha Q^2$ (see Eqs. (5.82), (E.40) (E.41) and (E.49)). When multiplied with the photon propagator at $q_0 = 0$, it gives zero for the $\delta(Q^2)$ part of the propagator, and cancels the $1/Q^2$ in the other part. This latter contribution will be labeled by the letter ‘X’ in order not to confuse it with the contributions of Fig. 5.1A.

The non-vanishing contribution

We denote the vertex contribution coming from Fig. 5.2A by

$$\Gamma_{\lambda\rho}^{(X)}(p, p') = \frac{e^2 \gamma^\alpha}{q^2} X_{\lambda\rho\alpha}(q), \quad (5.49)$$

where $X_{\lambda\rho\alpha}(q)$ is the photon-graviton mixing diagram with external momentum q

$$X_{\lambda\rho\alpha}(q) = \sum_f \int \frac{d^4 k}{(2\pi)^4} \text{Tr} \left[V_{\lambda\rho}(k, k') i S_f(k) i \mathcal{Q}_f \gamma_\alpha i S_f(k') \right]. \quad (5.50)$$

Then, taking the above discussion above into account, the quantity which will appear in the expression for the gravitational mass is given by

$$\tilde{\Gamma}_{\lambda\rho}^{(X)}(p, p') = \frac{e^2\gamma^\alpha}{q^2} [X_{\lambda\rho\alpha}(q) - X_{\lambda\rho\alpha}(0)]. \quad (5.51)$$

(As mentioned in Sec. 4.5, the sum in Eq. (5.50) is over all species of fermions in the medium, the charge of each species being denoted by $\mathcal{Q}_f$ with the convention that $\mathcal{Q}_e = -1$.) The medium-dependent contribution to $X_{\lambda\rho\alpha}(q)$ can be written as

$$X_{\lambda\rho\alpha}(q) = \sum_f \mathcal{Q}_f \int \frac{d^4k}{(2\pi)^3} \delta(k^2 - m_f^2) \eta_f(k) \left[\frac{A_{\lambda\rho\alpha}(k, k-q)}{q^2 - 2k \cdot q} + \frac{A_{\lambda\rho\alpha}(k+q, k)}{q^2 + 2k \cdot q} \right], \quad (5.52)$$

where, for arbitrary 4-momenta k_1 and k_2 , we have defined

$$\begin{aligned} A_{\lambda\rho\alpha}(k_1, k_2) &= \text{Tr} [V_{\lambda\rho}(k_1, k_2)(\not{k}_1 + m_f)\gamma_\alpha(\not{k}_2 + m_f)] \\ &= [(2k_{1\lambda}k_{1\rho} + k_{1\lambda}k_{2\rho} + k_{2\lambda}k_{1\rho})k_{2\alpha} + (m_f^2 - k_1 \cdot k_2)(\eta_{\lambda\alpha}k_{1\rho} + \eta_{\rho\alpha}k_{1\lambda}) \\ &\quad - 2\eta_{\lambda\rho}(k_1^2 - m_f^2)k_{2\alpha}] + [k_1 \leftrightarrow k_2]. \end{aligned} \quad (5.53)$$

Putting $k^2 = m_f^2$, we obtain

$$\begin{aligned} A_{\lambda\rho\alpha}(k, k-q) &= [8k_\lambda k_\rho - 4(k_\lambda q_\rho + k_\rho q_\lambda) + 2q_\lambda q_\rho]k_\alpha - [4k_\lambda k_\rho - (k_\lambda q_\rho + k_\rho q_\lambda)]q_\alpha \\ &\quad + k \cdot q [\eta_{\lambda\alpha}(2k - q)_\rho + \eta_{\rho\alpha}(2k - q)_\lambda] - 2\eta_{\lambda\rho}(q^2 - 2k \cdot q)k_\alpha. \end{aligned} \quad (5.54)$$

Since $A_{\lambda\rho\alpha}(k_1, k_2) = A_{\lambda\rho\alpha}(k_2, k_1)$ by definition, $A_{\lambda\rho\alpha}(k+q, k)$ is obtained by changing the sign of q in this expression.

5.4.4 Additional remarks

This is an appropriate point to consider the cases when 2-type vertices can occur in the diagrams contributing to the gravitational vertex. Since the vertices connected to the external legs can only be of 1-type, a 2-type vertex cannot occur in any of the diagrams of Fig. 5.1 or in the diagram of Fig. 5.2B. However, in Figs. 5.2A, 5.2C and 5.2D, the vertex where the photon line connects to the fermion loop can be of 2-type.

Now, due to the reason given in Sec. 4.5.1 (see after Eq. (4.92)), the unadorned tadpole with a 2-type vertex is just the negative of the unadorned tadpole with a 1-type vertex, and hence of $O(\kappa)$ because of the charge neutrality condition. So the diagrams of Figs. 5.2C and 5.2D, even with a 2-type vertex, are of $O(\kappa^2)$, and can be neglected.

In Fig. 5.2A, the occurrence of the 2-type vertex makes the photon propagator a “12” propagator. So the photon propagator is proportional to $\delta(q^2)$, which becomes $\delta(Q^2)$ for $q^0 = 0$. However, the value of the graviton momentum is taken to be $\vec{Q} \rightarrow 0$, and not $Q = 0$ (having $Q = 0$ in addition to $q^0 = 0$ would mean a potential constant in space and time, resulting in zero gravitational field). Consequently, $\delta(Q^2)$ in the “12” propagator causes this contribution to vanish. Therefore, 2-type vertices do not play any role in the present calculation.

Next, we make some comments on the expressions deduced in this section.

Firstly, one can easily check that the expressions given in Eqs. (5.39), (5.40), (5.43), (5.44), (5.45), (5.46), and (5.49) (coupled with (5.52)), all satisfy the dispersive condition of Eq. (5.33). (Similarly, it can be checked that the terms involving the thermal parts of two propagators from $\Gamma_{\lambda\rho}^{(A)}$, $\Gamma_{\lambda\rho}^{(B)}$, $\Gamma_{\lambda\rho}^{(C+D)}$ and $\Gamma_{\lambda\rho}^{(X)}$ are all absorptive.)

Secondly, it can be verified that the different contributions to the dispersive part of $\Gamma'_{\lambda\rho}$ also satisfy the charge conjugation relation given by Eq. (C.20).

Thirdly, the complete one-loop vertex function satisfies the transversality condition, which is implied by the gravitational gauge invariance of the theory. This is shown in Appendix D.

5.5 Calculation of the gravitational mass

As seen in Eq. (5.25), there are three types of $O(e^2)$ correction to the gravitational mass. One of them is proportional to the inertial mass that was calculated in Sec. 4.5.2, and another one involves the wave function renormalization factor derived in Sec. 4.5.3. In this section we find the contributions from the irreducible one-loop vertex diagrams. Since the expressions for those already have an explicit factor of e^2 outside the integral, to evaluate them we can use the tree-level values for the dispersion relation and the spinors associated with the external lepton.

5.5.1 Terms with the photon distribution from Fig. 5.1

We first evaluate those terms obtained in Sec. 5.4 that contain the photon distribution function. In fact, if the temperature of the ambient medium is low ($T \ll m_e$) and the chemical potential of the background electrons is zero, these are the only terms that contribute and they are precisely the ones that were calculated in Ref. [20]. Since we have

performed the calculations in a different way, using 1-particle irreducible diagrams only, the following results serve as a good checkpoint between the earlier calculations of Ref. [20] and ours.

Contribution (A1)

From the formula for the gravitational mass given in Eq. (5.25), it follows that we need to calculate the vertex only for $p = p'$, in which case

$$\Gamma'_{\lambda\rho}(A1)(p, p) = -e^2 \int \frac{d^4 k}{(2\pi)^3} \frac{\delta(k^2) \eta_\gamma(k)}{4(k \cdot p)^2} \Lambda_{\lambda\rho}(k + p, k + p). \quad (5.55)$$

From Eq. (5.37) it follows that, for any 4-vector y^μ ,

$$\Lambda_{\lambda\rho}(y, y) = -4y_\lambda y_\rho (\not{y} - 2m_\ell) + (y^2 - m_\ell^2) \left[(\gamma_\lambda y_\rho + \gamma_\rho y_\lambda) + 2\eta_{\lambda\rho} (\not{y} - 2m_\ell) \right], \quad (5.56)$$

which leads to

$$\begin{aligned} \left[\bar{u}_s(\vec{P}) \Gamma'_{\lambda\rho}(A1)(p, p) u_s(\vec{P}) \right]_{p^\mu=(m_\ell, \vec{0})} &= -e^2 \int \frac{d^4 k}{(2\pi)^3} \frac{\delta(k^2) \eta_\gamma(k)}{(k \cdot p)^2} \bar{u}_s(\vec{P}) \left[-\frac{k \cdot p}{m_\ell} k_\lambda p_\rho \right. \\ &\quad \left. + m_\ell (k_\lambda k_\rho + p_\lambda p_\rho) + \frac{(k \cdot p)^2}{m_\ell} \eta_{\lambda\rho} \right] u_s(\vec{P}), \end{aligned} \quad (5.57)$$

where we have used Eq. (4.58) and omitted the terms odd in k , which integrate to zero.

Using the notation

$$m'_{(A1)} = (2v^\lambda v^\rho - \eta^{\lambda\rho}) \left[\bar{u}_s(\vec{P}) \Gamma'_{\lambda\rho}(A1)(p, p) u_s(\vec{P}) \right]_{p^\mu=(m_\ell, \vec{0})}, \quad (5.58)$$

we obtain

$$\begin{aligned} m'_{(A1)} &= e^2 \int \frac{d^3 K}{(2\pi)^3} f_\gamma(K) \left[\frac{1}{m_\ell K} - \frac{m_\ell}{K^3} \right] \\ &= \frac{e^2 T^2}{12m_\ell} - \frac{e^2 m_\ell}{2\pi^2} \int_0^\infty \frac{dK}{K} f_\gamma(K). \end{aligned} \quad (5.59)$$

The remaining integral is infrared divergent, but its contribution to the gravitational mass is canceled by a similar term that arises from the wavefunction renormalization, as we show below.

Contribution (B1)

This term has to be treated carefully because the denominators in the integrand of Eq. (5.43) vanish for $q = 0$. Indeed, by using the formulas

$$\begin{aligned} \mathcal{P}r\left(\frac{1}{x}\right) &= \lim_{\epsilon \rightarrow 0} \frac{x}{x^2 + \epsilon^2}, \\ \delta(x) &= \lim_{\epsilon \rightarrow 0} \frac{1}{\pi} \frac{\epsilon}{x^2 + \epsilon^2} \end{aligned} \quad (5.60)$$

(with $\mathcal{P}r$ denoting the principal value), one can show that [54]

$$\mathcal{P}r\left(\frac{1}{k^2 - m^2}\right)\delta(k^2 - m^2) = \frac{1}{2}\frac{d}{dm^2}\delta(k^2 - m^2) \quad (5.61)$$

which is ill-defined even for $m \neq 0$, whereas for $m = 0$, one can interpret the R.H.S. of Eq. (5.61) only as a limiting form with $m \rightarrow 0$. However, a careful evaluation of the term given in Eq. (5.43), following the procedure indicated in Eq. (5.25), shows that the limit exists. Denoting

$$m'_{(B1)} \equiv (2v^\lambda v^\rho - \eta^{\lambda\rho}) \lim_{\vec{P} \rightarrow 0} \left\{ \left[\bar{u}_s(\vec{P}') \Gamma'_{\lambda\rho}{}^{(B1)}(p, p') u_s(\vec{P}) \right]_{\substack{E'_\ell = E_\ell \\ \vec{P}' \rightarrow P}} \right\}, \quad (5.62)$$

the result is

$$m'_{(B1)} = -\frac{e^2 T^2}{3m_\ell}. \quad (5.63)$$

The details of the derivation of this result are given in Appendix E.1. This result can also be obtained by means of a special regularization at $q = 0$ (following Eq. (5.61)), as shown in Appendix E.4.

Contributions (C1+D1)

We can proceed as in the evaluation of $m'_{(A1)}$ above. Thus, from Eq. (5.45),

$$\begin{aligned} \left[\bar{u}_s(\vec{P}) \Gamma'_{\lambda\rho}{}^{(C1+D1)}(p, p) u_s(\vec{P}) \right]_{p^\mu = (m_\ell, \vec{0})} &= -e^2 a_{\mu\nu\lambda\rho} \int \frac{d^4 k}{(2\pi)^3} \frac{\delta(k^2) \eta_\gamma(k)}{k \cdot p} \\ &\times \left[\bar{u}_s(\vec{P}) \gamma^\mu \not{k} \gamma^\nu u_s(\vec{P}) \right], \end{aligned} \quad (5.64)$$

using the fact that $a_{\mu\nu\lambda\rho}$ is symmetric in the indices μ, ν . Then using

$$(2v^\lambda v^\rho - \eta^{\lambda\rho}) a_{\mu\nu\lambda\rho} = -\eta_{\mu\nu} - 2v_\mu v_\nu \quad (5.65)$$

it follows that

$$\begin{aligned} m'_{(C1+D1)} &\equiv (2v^\lambda v^\rho - \eta^{\lambda\rho}) \left[\bar{u}_s(\vec{P}) \Gamma'_{\lambda\rho}{}^{(C1+D1)}(p, p) u_s(\vec{P}) \right]_{p^\mu = (m_\ell, \vec{0})} \\ &= 0. \end{aligned} \quad (5.66)$$

5.5.2 Terms with the electron distribution from Fig. 5.1

These terms contribute only to the gravitational vertex involving electrons and positrons. The integration over k_0 and the angular variables can be done exactly. The remaining integral can be evaluated analytically only for special cases of the distribution functions, some of which we consider afterwards.

Contribution (A2)

As can be seen from Eq. (5.40), the denominators of the integrand of this term vanish as $q \rightarrow 0$. Consequently, the prescription indicated in Eq. (5.25) has to be followed carefully in this case. As we show in detail in Appendix E.2, defining

$$m'_{(A2)} \equiv (2v^\lambda v^\rho - \eta^{\lambda\rho}) \lim_{\vec{P} \rightarrow 0} \left\{ \left[\bar{u}_s(\vec{P}') \Gamma_{\lambda\rho}'^{(A2)}(p, p') u_s(\vec{P}) \right]_{\substack{E'_\ell = E_\ell \\ \vec{P}' \rightarrow P}} \right\}, \quad (5.67)$$

the final result for this term is

$$\begin{aligned} m'_{(A2)} = & \frac{e^2}{m_e} \int \frac{d^3 K}{(2\pi)^3 2E_K} \left\{ \frac{2E_K^2 - m_e^2}{E_K} \left(\frac{E_K - 2m_e}{E_K - m_e} \frac{\partial f_e}{\partial E_K} + \frac{E_K + 2m_e}{E_K + m_e} \frac{\partial f_{\bar{e}}}{\partial E_K} \right) \right. \\ & + \frac{2E_K^4 - E_K^3 m_e - 5E_K^2 m_e^2 + 2E_K m_e^3 - 2m_e^4}{m_e E_K^2 (E_K - m_e)} f_e \\ & \left. - \frac{2E_K^4 + E_K^3 m_e - 5E_K^2 m_e^2 - 2E_K m_e^3 - 2m_e^4}{m_e E_K^2 (E_K + m_e)} f_{\bar{e}} \right\}, \quad (5.68) \end{aligned}$$

where E_K is defined in Eq. (2.6). This result is also derived in Appendix E.4 by means of a special regularization at $q = 0$.

Contribution (B2)

From Eq. (5.44) it is seen that the integrand is not singular in the limit $q \rightarrow 0$. Therefore we can evaluate directly

$$\Gamma_{\lambda\rho}'^{(B2)}(p, p) = -e^2 \int \frac{d^4 k}{(2\pi)^3} \delta(k^2 - m_e^2) \eta_e(k) \gamma^\nu(\not{k} + m_e) \gamma^\mu \frac{C_{\mu\nu\lambda\rho}(p - k, p - k)}{(p - k)^4} \quad (5.69)$$

and the contribution to the gravitational mass is given by

$$m'_{(B2)} = (2v^\lambda v^\rho - \eta^{\lambda\rho}) \left[\bar{u}_s(\vec{P}) \Gamma_{\lambda\rho}'^{(B2)}(p, p) u_s(\vec{P}) \right]_{p^\mu = (m_e, \vec{0})}. \quad (5.70)$$

In the expression for $C_{\mu\nu\lambda\rho}$, any term having a factor of $(p - k)_\mu$ or $(p - k)_\nu$ does not contribute to the integral. This is because, within the spinors, we can write

$$\gamma^\nu(\not{k} + m_e) \gamma^\mu(p - k)_\mu = \gamma^\nu(\not{k} + m_e)(m_e - \not{k}) = \gamma^\nu(m_e^2 - k^2), \quad (5.71)$$

which vanishes because of the δ -function. The argument is similar for $(p - k)_\nu$. Thus,

$$\begin{aligned} & (2v^\lambda v^\rho - \eta^{\lambda\rho}) \bar{u}_s(\vec{P}) \gamma^\nu(\not{k} + m_e) \gamma^\mu C_{\mu\nu\lambda\rho}(p - k, p - k) u_s(\vec{P}) \\ & = \bar{u}_s(\vec{P}) \left\{ 8(\not{k} - 2m_e)(p \cdot v - k \cdot v)^2 + 4(m_e - 2k \cdot v \not{p})(p - k)^2 \right\} u_s(\vec{P}), \quad (5.72) \end{aligned}$$

ignoring all terms which have a factor of $k^2 - m_e^2$. Using Eqs. (4.58) and (4.59), we then obtain

$$\begin{aligned} m'_{(B2)} &= -\frac{2e^2}{m_e^2} \int \frac{d^4k}{(2\pi)^3} \delta(k^2 - m_e^2) \eta_e(k) \left(k_0 + \frac{m_e^2}{k_0 - m_e} \right) \\ &= -\frac{e^2}{m_e^2} \int \frac{d^3K}{(2\pi)^3} (f_e - f_{\bar{e}}) - \frac{e^2}{2\pi^2} \int dK \frac{K^2}{E_K} \left[\frac{f_e(E_K)}{E_K - m_e} - \frac{f_{\bar{e}}(E_K)}{E_K + m_e} \right]. \end{aligned} \quad (5.73)$$

Contributions (C2+D2)

Similarly, for this term we can evaluate directly

$$m'_{(C2+D2)} = (2v^\lambda v^\rho - \eta^{\lambda\rho}) \left[\bar{u}_s(\vec{P}) \Gamma'_{\lambda\rho}{}^{(C2+D2)}(p, p) u_s(\vec{P}) \right]_{p^\mu=(m_e, \vec{0})}, \quad (5.74)$$

with

$$\Gamma'_{\lambda\rho}{}^{(C2+D2)}(p, p) = 2e^2 a_{\mu\nu\lambda\rho} \int \frac{d^4k}{(2\pi)^3} \delta(k^2 - m_e^2) \eta_e(k) \frac{\gamma^\mu(\not{k} + m_e) \gamma^\nu}{(k - p)^2}. \quad (5.75)$$

By straight forward algebra

$$(2v^\lambda v^\rho - \eta^{\lambda\rho}) a_{\mu\nu\lambda\rho} \gamma^\mu(\not{k} + m_e) \gamma^\nu = 4\not{k} - 6m_e - 4k \cdot v \not{v}, \quad (5.76)$$

and using Eqs. (4.58) and (4.59),

$$\begin{aligned} m'_{(C2+D2)} &= 6e^2 \int \frac{d^4k}{(2\pi)^3} \delta(k^2 - m_e^2) \eta_e(k) \frac{1}{k_0 - m_e} \\ &= \frac{3e^2}{2\pi^2} \int dK \frac{K^2}{E_K} \left[\frac{f_e(E_K)}{E_K - m_e} - \frac{f_{\bar{e}}(E_K)}{E_K + m_e} \right]. \end{aligned} \quad (5.77)$$

5.5.3 Terms from Fig. 5.2

The contribution to the gravitational mass due to this term is

$$m'_{(X)} \equiv (2v^\lambda v^\rho - \eta^{\lambda\rho}) \lim_{\vec{P} \rightarrow 0} \left\{ \left[\bar{u}_s(\vec{P}') \tilde{\Gamma}_{\lambda\rho}^{(X)}(p, p') u_s(\vec{P}) \right]_{\substack{q^0=0 \\ \vec{Q} \rightarrow 0}} \right\}, \quad (5.78)$$

where, from Eq. (5.51),

$$\tilde{\Gamma}_{\lambda\rho}^{(X)}(p, p') \Big|_{q^0=0} = -\frac{e^2 \gamma^\alpha}{Q^2} \left[X_{\lambda\rho\alpha}(q) \Big|_{q^0=0} - X_{\lambda\rho\alpha}(0) \right]. \quad (5.79)$$

Using the expression for $A_{\lambda\rho\alpha}$ from Eq. (5.54) we obtain

$$\begin{aligned} (2v^\lambda v^\rho - \eta^{\lambda\rho}) X_{\lambda\rho\alpha}(q) \Big|_{q^0=0} &= 8 \sum_f \mathcal{Q}_f \int \frac{d^4k}{(2\pi)^3} \delta(k^2 - m_f^2) \eta_f(k) k_0 v_\alpha \left[(2k_0^2 - m_f^2 - \frac{1}{2}Q^2) \right. \\ &\quad \times \left. \left(\frac{1}{2\vec{K} \cdot \vec{Q} - Q^2} - \frac{1}{2\vec{K} \cdot \vec{Q} + Q^2} \right) \right], \end{aligned} \quad (5.80)$$

where we have omitted the terms that vanish by symmetric integration over $\vec{K}$, as well as all those terms that are independent of q , because they drop out of Eq. (5.79), and in addition all the terms that are proportional to q_α , because in Eq. (5.78) they yield a factor of $\not{q}$ which vanishes between spinors. Performing the integration over k_0 ,

$$\begin{aligned} (2v^\lambda v^\rho - \eta^{\lambda\rho})X_{\lambda\rho\alpha}(q)\Big|_{q^0=0} &= 4v_\alpha \sum_f \mathcal{Q}_f \int \frac{d^3K}{(2\pi)^3} (f_f - f_{\bar{f}})(2E_K^2 - m_f^2 - \frac{1}{2}Q^2) \\ &\quad \times \left(\frac{1}{2\vec{K} \cdot \vec{Q} - Q^2} - \frac{1}{2\vec{K} \cdot \vec{Q} + Q^2} \right). \end{aligned} \quad (5.81)$$

For the term that contains an explicit factor of Q^2 in the numerator we use the angular integration formula of Eq. (E.9), which yields

$$\begin{aligned} (2v^\lambda v^\rho - \eta^{\lambda\rho})X_{\lambda\rho\alpha}(q)\Big|_{q^0=0} &= v_\alpha \sum_f \mathcal{Q}_f \left[\frac{Q^2}{2\pi^2} \int dK (f_f - f_{\bar{f}}) + 4 \int \frac{d^3K}{(2\pi)^3} (f_f - f_{\bar{f}}) \right. \\ &\quad \left. \times (2E_K^2 - m_f^2) \left(\frac{1}{2\vec{K} \cdot \vec{Q} - Q^2} - \frac{1}{2\vec{K} \cdot \vec{Q} + Q^2} \right) \right]. \end{aligned} \quad (5.82)$$

The evaluation of the rest of the integral is presented in Appendix E.3. Substituting the results into Eq. (5.79), the contribution of this diagram to the gravitational mass is found to be given by

$$m'_{(X)} = -e^2 \sum_f \frac{\mathcal{Q}_f}{6\pi^2} \int dK \left[(f_f - f_{\bar{f}}) - \frac{2E_K^2 - m_f^2}{2E_K} \frac{\partial}{\partial E_K} (f_f - f_{\bar{f}}) \right]. \quad (5.83)$$

5.5.4 Summary

Starting from Eq. (5.25), the total contribution to the gravitational mass of charged leptons can be written in the form

$$M'_\ell = m_\ell + m'_{\ell 1} + m'_{\ell 2} + m'_{(X)}, \quad (5.84)$$

where $m'_{(X)}$ is the contribution from Eq. (5.83), which is the same for all charged leptons, $m'_{\ell 1}$ represents the terms that contain the photon distribution function, and $m'_{\ell 2}$ contains the terms that depend on the electron distribution function. They are given as follows.

Substituting into Eq. (5.25) the results given in Eqs. (5.59), (5.63) and (5.66), and using the expression for the wave-function normalization and the correction to the inertial mass given in Eqs. (4.120) and (4.105), we find

$$m'_{\ell 1} = -\frac{e^2 T^2}{12m_\ell}, \quad (5.85)$$

in agreement with the DHR result[20], quoted in Eq. (1.2). Notice that the infrared divergence contained in the $m'_{(A1)}$ term cancels with a similar one that arises from the wave function renormalization correction $\zeta_{\ell 1}$.

The terms from the diagrams in Fig. 5.1 that involve the fermion distribution function contribute only to the gravitational mass of the electron, and therefore

$$m'_{\mu 2} = m'_{\tau 2} = 0. \quad (5.86)$$

The individual contributions of this type to the electron gravitational mass appear in Eqs. (5.68), (5.73) and (5.77). Substituting those results into Eq. (5.25), and using the results for the inertial mass and the wave-function normalization factor, given in Eqs. (4.113) and (4.123) respectively, we obtain

$$\begin{aligned} m'_{e2} = & \frac{e^2}{\pi^2 m_e} \int_0^\infty dK \frac{K^2}{2E_K} \left\{ \left(\frac{3}{2} + \frac{m_e^2}{E_K^2} - \frac{m_e}{E_K - m_e} \right) f_e(E_K) \right. \\ & + \left(\frac{3}{2} + \frac{m_e^2}{E_K^2} + \frac{m_e}{E_K + m_e} \right) f_{\bar{e}}(E_K) \\ & \left. + \frac{2E_K^2 - m_e^2}{2E_K} \left(\frac{E_K - 2m_e}{E_K - m_e} \frac{\partial f_e}{\partial E_K} + \frac{E_K + 2m_e}{E_K + m_e} \frac{\partial f_{\bar{e}}}{\partial E_K} \right) \right\}. \end{aligned} \quad (5.87)$$

The corresponding formulas for the antileptons are obtained by making the substitution given in Eq. (4.115), as explained after Eq. (5.31). Thus,

$$M'_\ell = m_\ell + m'_{\ell 1} + m'_{\ell 2} - m'_X, \quad (5.88)$$

where

$$m'_{\bar{\mu} 2} = m'_{\bar{\tau} 2} = 0, \quad (5.89)$$

while the result for $m'_{\bar{e} 2}$ is obtained from Eq. (5.87) by making the substitution $f_e \leftrightarrow f_{\bar{e}}$.

5.6 Results for particular cases

In contrast with the $m_{\ell 1}$ and $m'_{\ell 1}$ terms, which depend on the photon momentum distribution, m_{e2} , $m_{\bar{e} 2}$, m'_{e2} , $m'_{\bar{e} 2}$ and $m'_{(X)}$ depend on the fermion distribution functions and cannot be evaluated exactly in the general case. Therefore, for illustration, we consider in detail their calculation for the specific situation in which the background is composed of non-relativistic protons and electrons. In this case we can set (for $f = e, p$)

$$f_f(E) \approx 0 \quad (5.90)$$

and

$$E_K \approx m_f + \frac{K^2}{2m_f}. \quad (5.91)$$

We consider in detail two cases separately, according to whether the electron gas is classical or degenerate.

5.6.1 Classical electron gas and classical proton gas

In this situation we can set

$$f_f(E) = e^{-\beta(E-\mu_f)} \quad (5.92)$$

for both $f = e, p$. This implies the relation

$$\frac{\partial f_f}{\partial E} = -\beta f_f, \quad (5.93)$$

as well as the integration formula

$$\int dK K^{2r} f_f = 2\pi^{3/2} \Gamma\left(r + \frac{1}{2}\right) \left(\frac{\beta}{2m_f}\right)^{1-r} n_f, \quad (5.94)$$

where n_f is the number density, given by

$$n_f = 2 \int \frac{d^3K}{(2\pi)^3} f_f(E) \approx 2 \left(\frac{m_f}{2\pi\beta}\right)^{3/2} e^{-\beta(m_f-\mu_f)}. \quad (5.95)$$

Let us consider m_{e2} and $m_{\bar{e}2}$, given in Eqs. (4.113) and (4.114), respectively. Setting $f_{\bar{e}} = 0$ and using Eq. (5.91) to expand the co-efficients of f_e in the integrands in powers of K , the remaining integrals are evaluated by means of Eq. (5.94) to yield

$$\begin{aligned} m_{e2} &= -\frac{e^2 n_e}{2m_e T} + O(e^2 n_e / m_e^2), \\ m_{\bar{e}2} &= \frac{3e^2 n_e}{8m_e^2} + O(e^2 n_e T^2 / m_e^4). \end{aligned} \quad (5.96)$$

(Note that the subleading terms are small because they are suppressed by powers of T/m_e , and $T < m_e$ for the electrons to be non-relativistic.) Similarly, from Eq. (5.83) we obtain for this case

$$\begin{aligned} m'_{(X)} &= -\frac{7e^2}{24T} \sum_{f=e,p} \frac{Q_f n_f}{m_f} + O(e^2 n_e T / m_e^3) \\ &\approx \frac{7e^2 n_e}{24m_e T}, \end{aligned} \quad (5.97)$$

where we have used the charge-neutrality condition which, neglecting terms $O(\kappa)$, is simply $\sum_f \mathcal{Q}_f n_f = 0$. Applying the same procedure in Eq. (5.87), the leading contribution, in powers of T/m_e , comes from the $\partial f_e / \partial E_K$ term in that equation, and leads to

$$m'_{e2} = \frac{e^2 n_e}{2T^2} + O(e^2 n_e / T m_e). \quad (5.98)$$

By the substitution indicated in Eq. (4.115), the corresponding result for the positron is

$$m'_{\bar{e}2} = -\frac{3e^2 n_e}{8m_e T} + O(e^2 n_e / m_e^2). \quad (5.99)$$

Therefore, using Eqs. (4.104), (5.84) and (5.88), the inertial and gravitational masses for charged leptons ℓ other than the electron are obtained as

$$\begin{aligned} M_{\bar{\ell}} = M_{\ell} &= m_{\ell} + \frac{e^2 T^2}{12m_{\ell}}, \\ M'_{\ell, \bar{\ell}} &= m_{\ell} - \frac{e^2 T^2}{12m_{\ell}} \pm \frac{7e^2 n_e}{24m_e T}, \end{aligned} \quad (5.100)$$

where the upper sign corresponds to the leptons and the lower one to the anti-leptons. The corresponding formulas for the electron are

$$\begin{aligned} M_e &= m_e + \frac{e^2 T^2}{12m_e} - \frac{e^2 n_e}{2m_e T}, \\ M'_e &= m_e - \frac{e^2 T^2}{12m_e} + \frac{e^2 n_e}{2T^2}, \end{aligned} \quad (5.101)$$

and for the positron they are

$$\begin{aligned} M_{\bar{e}} &= m_e + \frac{e^2 T^2}{12m_e} + \frac{3e^2 n_e}{8m_e^2}, \\ M'_{\bar{e}} &= m_e - \frac{e^2 T^2}{12m_e} - \frac{2e^2 n_e}{3m_e T}. \end{aligned} \quad (5.102)$$

We now estimate how large these corrections could be for the electron. Those due to the photon background were estimated by DHR [20] and were found to be very small. This is because the fractional changes in the inertial and gravitational masses in that case are $O(e^2 T^2 / m_e^2)$, and $T < m_e$. Therefore, neglecting that contribution, the fractional changes are given by

$$\left| \frac{M_e - m_e}{m_e} \right| = \frac{e^2 n_e}{2m_e^2 T} \quad (5.103)$$

$$\left| \frac{M'_e - m_e}{m_e} \right| = \frac{e^2 n_e}{2m_e T^2}. \quad (5.104)$$

Although it may seem that the effects are more noticeable as the temperature decreases, they are bounded by the conditions that the electron gas is non-degenerate and non-interacting, which require that

$$T > \frac{n_e^{2/3}}{m_e} \quad (5.105)$$

and

$$T > \frac{e^2}{r_{\text{av}}} \sim e^2 n_e^{1/3}, \quad (5.106)$$

since $r_{\text{av}} \sim n_e^{-1/3}$ [55]. Using the fact that $n_p = n_e$, it follows that the corresponding conditions for the proton gas do not imply further restrictions, because they are automatically satisfied whenever Eqs. (5.105) and (5.106) hold.

By writing the right-hand side of Eq. (5.103) in the alternative forms

$$\begin{aligned} \frac{e^2 n_e}{2m_e^2 T} &= \frac{1}{2} \left(\frac{e^2 n_e^{1/3}}{T} \right) \left(\frac{n_e^{1/3}}{m_e} \right)^2 \\ &= \left(\frac{n_e^{2/3}}{m_e T} \right)^2 \left(\frac{T}{2m_e} \right) \left(\frac{e^2 m_e}{n_e^{1/3}} \right), \end{aligned} \quad (5.107)$$

it is seen that

$$\left| \frac{M_e - m_e}{m_e} \right| < \begin{cases} e^4/2 & \text{if } n_e^{1/3} < e^2 m_e, \\ T/2m_e & \text{if } n_e^{1/3} > e^2 m_e. \end{cases} \quad (5.108)$$

Similarly, writing

$$\begin{aligned} \frac{e^2 n_e}{2m_e T^2} &= \frac{1}{2} \left(\frac{e^2 n_e^{1/3}}{T} \right)^2 \left(\frac{n_e^{1/3}}{e^2 m_e} \right) \\ &= \frac{1}{2} \left(\frac{n_e^{2/3}}{m_e T} \right)^2 \left(\frac{e^2 m_e}{n_e^{1/3}} \right), \end{aligned} \quad (5.109)$$

it follows that

$$\left| \frac{M'_e - m_e}{m_e} \right| < \frac{1}{2} \quad (5.110)$$

in either case. Therefore, while the fractional correction to the electron's inertial mass is likely to be small in most situations with the conditions that we are presently considering, the fractional change in the gravitational mass could be substantial. For example, if we use

the temperature and density at the solar core, i.e., $T = 1.57 \times 10^7$ K, $n_e = 9.5 \times 10^{25} \text{ cm}^{-3}$, we obtain

$$\begin{aligned} \left| \frac{M_e - m_e}{m_e} \right| &= 9.8 \times 10^{-5}, \\ \left| \frac{M'_e - m_e}{m_e} \right| &= 3.5 \times 10^{-2}, \end{aligned} \quad (5.111)$$

which shows that the correction to the gravitational mass of the electron can at least be appreciable in realistic physical situations.

5.6.2 Degenerate electron gas and classical proton gas

For a degenerate electron gas,

$$T < \frac{n_e^{2/3}}{m_e} \sim \frac{K_F^2}{m_e}, \quad (5.112)$$

where K_F is the Fermi momentum of the electron gas. We assume

$$K_F < m_e \quad (5.113)$$

so that the electrons are non-relativistic. We also assume the electron gas to be non-interacting. This requires that the average kinetic energy $\sim K_F^2/m_e$ of an electron is larger than the average Coulomb interaction energy $\sim e^2 n_e^{1/3} \sim e^2 K_F$. (Unlike the case of the classical gas, T contributes negligibly to the average kinetic energy in the present case; see Eq. (5.112).) This implies that [56]

$$K_F > e^2 m_e. \quad (5.114)$$

We now show that, under these conditions, the protons can be treated as a non-interacting, non-relativistic, classical gas [57]. The only additional condition we have to impose is the condition for non-interaction for a classical proton gas:

$$T > e^2 n_p^{1/3} \sim e^2 K_F \quad (5.115)$$

(using $n_p = n_e$). Since Eqs. (5.112) and (5.113) only imply that $T < K_F$, it is still possible to satisfy Eq. (5.115). Eqs. (5.112) and (5.113) also imply that $T < m_e$. So the non-relativistic condition for the classical proton gas is automatically satisfied. Finally, since $K_F < m_e \ll m_p$, Eq. (5.115) gives $T \gg e^2 K_F^2/m_p$, so that the non-degeneracy condition

$$T > \frac{K_F^2}{m_p} \sim \frac{n_p^{2/3}}{m_p} \quad (5.116)$$

is satisfied as well.

Therefore, Eq. (5.92) applies to the proton, while for the electron

$$f_e = \Theta(K_F - K) \quad (5.117)$$

with

$$K_F = (3\pi^2 n_e)^{1/3}, \quad (5.118)$$

which in turn imply the relation

$$\frac{df_e}{dK} = -\delta(K_F - K). \quad (5.119)$$

We repeat the calculation of the quantities m_{e2} , $m_{\bar{e}2}$, m'_{e2} , $m'_{\bar{e}2}$ and m'_X for this case, neglecting the terms that are a factor $\sim O(K_F^2/m_e^2)$ smaller than the ones that we retain. From Eqs. (4.113) and (4.114), setting $f_{\bar{e}} = 0$ and using Eq. (5.91), we obtain

$$\begin{aligned} m_{e2} &= -\frac{e^2 K_F}{2\pi^2}, \\ m_{\bar{e}2} &= \frac{e^2 K_F^3}{8\pi^2 m_e^2}. \end{aligned} \quad (5.120)$$

From Eq. (5.83),

$$m'_{(X)} = \frac{e^2}{6\pi^2} \int dK \left[f_e - \left(K + \frac{m_e^2}{2K} \right) \frac{df_e}{dK} \right] - \frac{7e^2}{24T} \frac{n_p}{m_p}, \quad (5.121)$$

where we have borrowed the result for the proton contribution from Eq. (5.97), while in the electron term we have expressed E_K in terms of K and used

$$\frac{d}{dE_K} = \frac{E_K}{K} \frac{d}{dK} \quad (5.122)$$

for any function of E_K . Using Eqs. (5.117) and (5.119) this finally yields

$$m'_{(X)} = \frac{e^2 m_e^2}{12\pi^2 K_F}. \quad (5.123)$$

Here we have neglected the proton contribution because it is $\sim e^2 K_F^3 / (T m_p) < e^2 K_F$ from Eq. (5.116). In a similar fashion, from Eq. (5.87),

$$m'_{e2} = \frac{e^2 m_e^2}{2\pi^2 K_F}, \quad (5.124)$$

and by the substitution indicated in Eq. (4.115), the corresponding result for the positron is

$$m'_{\bar{e}2} = -\frac{3e^2 K_F}{8\pi^2}. \quad (5.125)$$

Thus, substituting these results into Eqs. (4.104), (5.84) and (5.88), we obtain the following expressions for the inertial and gravitational masses, retaining only the leading terms in powers of K_F/m_e . For the charged leptons ℓ other than the electron,

$$\begin{aligned} M_{\bar{\ell}} = M_{\ell} &= m_{\ell} + \frac{e^2 T^2}{12m_{\ell}} \\ M'_{\ell, \bar{\ell}} &= m_{\ell} - \frac{e^2 T^2}{12m_{\ell}} \pm \frac{e^2 m_e^2}{12\pi^2 K_F}, \end{aligned} \quad (5.126)$$

with the upper sign corresponding to the leptons and the lower one to the anti-leptons, while for the electron

$$\begin{aligned} M_e &= m_e + \frac{e^2 T^2}{12m_e} - \frac{e^2 K_F}{2\pi^2}, \\ M'_e &= m_e - \frac{e^2 T^2}{12m_e} + \frac{7e^2 m_e^2}{12\pi^2 K_F}, \end{aligned} \quad (5.127)$$

and for the positron

$$\begin{aligned} M_{\bar{e}} &= m_e + \frac{e^2 T^2}{12m_e} + \frac{e^2 K_F^3}{8\pi^2 m_e^2}, \\ M'_{\bar{e}} &= m_e - \frac{e^2 T^2}{12m_e} - \frac{e^2 m_e^2}{12\pi^2 K_F}. \end{aligned} \quad (5.128)$$

It is interesting to note that Eqs. (5.112) and (5.113) imply that the photon contributions in Eqs. (5.126)-(5.128) are much smaller than the contribution due to the electron background in each case. In fact, using Eq. (5.114), we see that the fractional corrections to the gravitational mass can be as large as about $7/12\pi^2$ for the electron and $1/12\pi^2$ for the positron and the other leptons.

CHAPTER 6

Some aspects of high-temperature QED

6.1 Introduction

So far in this work, we have considered low temperatures and densities. Such thermal backgrounds ensure the validity of a straightforward expansion in e^2 , something we have always assumed. A negative illustration of this point is provided by Eqs. (4.104) and (4.105) for the inertial mass of the charged lepton in photon background. It is clear from these equations that if T exceeds m_ℓ by a large amount, the change in inertial mass can no longer be treated as a perturbation.

In this chapter, we turn to an entirely different kind of thermal background, namely, a background at high temperature. More specifically, we will now consider high-temperature QED with exact chiral invariance and at zero chemical potential. In this case, the medium effects are not, in general, of $O(e^2)$ (see, for example, Sec. 6.5.1). Exact chiral invariance means that the full Dirac action of Eq. (4.71) is invariant under $\psi \rightarrow e^{i\gamma_5\theta}\psi$ for any real constant θ . So, the electron is massless in vacuum, and the self-energy cannot have the constant term on the R.H.S. of Eq. (4.6). (Chiral symmetry also rules out the $\sigma^{\mu\nu}p_\mu v_\nu$ term in the self-energy, which, anyway, is absent at one-loop, beyond which we will not be interested in.) Thus, we now have the general form

$$\Sigma(k) = a\not{k} + b\not{p} \tag{6.1}$$

for the self-energy of the electron. We shall use k^μ to denote the electron four-momentum in this chapter [58]. The scalars a and b are then functions of

$$\omega \equiv k \cdot v, \quad K \equiv [(k.v)^2 - k^2]^{1/2} \tag{6.2}$$

which satisfy

$$k^2 = \omega^2 - K^2. \quad (6.3)$$

The pole in the full propagator is given by

$$f(\omega, K) \equiv (\omega - K)(1 - a) - b = 0. \quad (6.4)$$

(This is nothing but Eq. (4.34) at zero mass. Since we now take the chemical potential to be zero, C symmetry is unbroken, and the particle and antiparticle dispersion relations are the same. So, instead of E_f and $E_{\bar{f}}$, we just use ω to denote the energy.) To leading order in T and at one-loop, it was shown by Weldon in his seminal work [5] that

$$\begin{aligned} a &= -\frac{e^2 T^2}{8K^2} \left[1 - \frac{\omega}{2K} \ln \left(\frac{\omega + K}{\omega - K} \right) \right], \\ b &= \frac{e^2 T^2}{8K} \left[\frac{\omega}{K} - \frac{1}{2} \left(\frac{\omega^2}{K^2} - 1 \right) \ln \left(\frac{\omega + K}{\omega - K} \right) \right]. \end{aligned} \quad (6.5)$$

Substitution of Eq. (6.5) into Eq. (6.4) then gives the electron dispersion relation

$$\omega - K = \frac{e^2 T^2}{8K} \left[1 + \frac{1}{2} \left(1 - \frac{\omega}{K} \right) \ln \left(\frac{\omega + K}{\omega - K} \right) \right]. \quad (6.6)$$

An important point is that if we write $\omega - K = b/(1 - a) \approx b$ from Eq. (6.4) by neglecting a^2 and ab , and then substitute Eq. (6.5), we do not get the correct answer given in Eq. (6.6). This is an example of what we meant by the invalidity of a straightforward expansion in e^2 at the beginning of this section.

In the same work, Weldon demonstrated the gauge independence of this leading order dispersion relation, by showing that gauge dependence appears in the self-energy only at subleading order in temperature. However, one expects that even this subleading order gauge dependence should not show up in the dispersion relation. We are therefore going to investigate the gauge independence of the dispersion relation after taking the subleading temperature dependence into account [23].

6.2 One-loop electron self-energy in a general linear covariant gauge

$\text{Re } \Sigma$ is defined through the equations

$$a(\omega, K) = a_R(\omega, K) + ia_I(\omega, K), \quad (6.7)$$

$$b(\omega, K) = b_R(\omega, K) + ib_I(\omega, K), \quad (6.8)$$

$$\text{Re } \Sigma = a_R \not{k} + b_R \not{\epsilon}. \quad (6.9)$$

To obtain the dispersion relation, we need to consider only $\text{Re } \Sigma$ and, consequently, only the “11” propagators, at one-loop [59]. As in Chapter 4, *we will denote $\text{Re } \Sigma$ by just Σ* . Because of the additional features of masslessness of the fermion, zero chemical potential, and inclusion of gauge dependence in the photon propagator, it will be convenient for us to rewrite the “11” propagators for the purpose of the calculations in this chapter. Thus, we are going to use the free-particle massless fermion propagator at zero chemical potential given by

$$S(p) = \not{p} \left[\frac{1}{p^2 + i\epsilon} + 2\pi i \delta(p^2) f_F(p) \right], \quad (6.10)$$

$$f_F(p) = [e^{|p_0|/T} + 1]^{-1}, \quad (6.11)$$

and the free photon propagator in a general covariant gauge given by [60]

$$D^{\mu\nu}(p) = D_{\text{FG}}^{\mu\nu}(p) + D_{\xi}^{\mu\nu}(p), \quad (6.12)$$

$$D_{\text{FG}}^{\mu\nu}(p) = -\eta^{\mu\nu} \left[\frac{1}{p^2 + i\epsilon} - 2\pi i \delta(p^2) f_B(p) \right], \quad (6.13)$$

$$D_{\xi}^{\mu\nu}(p) = -\xi p^{\mu} p^{\nu} \left[\frac{1}{(p^2 + i\epsilon)^2} + 2\pi i f_B(p) \frac{d\delta(p^2)}{dp^2} \right], \quad (6.14)$$

$$f_B(p) = [e^{|p_0|/T} - 1]^{-1}. \quad (6.15)$$

Here ξ is the gauge parameter and FG denotes the Feynman gauge ($\xi = 0$). Since *the calculation will be performed in the rest-frame of the medium*, we have used p_0 in the place of $p \cdot v$ in Eqs. (6.11) and (6.15). In this frame, we define the components of the electron four-momentum by writing

$$k^{\mu} = (k^0, \vec{K}). \quad (6.16)$$

Note that this choice of frame will not lead to any loss of generality in the proof of gauge independence, since the dispersion relation is Lorentz invariant.

Let us write

$$\Sigma = \Sigma^{T=0} + \Sigma'. \quad (6.17)$$

We now evaluate the real parts of these functions by putting the expressions for the fermion and the photon propagators in the expression for the self-energy given by Eq. (4.93). Let us first consider the $T = 0$ part

$$\Sigma^{T=0} = a^{T=0}(k^2) \not{k}, \quad (6.18)$$

and write $a^{T=0} = a_{\text{FG}}^{T=0} + a_{\xi}^{T=0}$. Dimensional regularization of $\Sigma_{\text{FG}}^{T=0}$ in $4 - \epsilon'$ dimensions gives

$$a_{\text{FG}}^{T=0} = -\frac{e^2}{(4\pi)^2} \left[\frac{2}{\epsilon'} - \gamma - 1 + \ln(4\pi) - 2 \int_0^1 dx \, x \ln(x(1-x)k^2) + O(\epsilon') \right] \quad (6.19)$$

where γ is the Euler-Mascheroni constant. On the other hand, one easily obtains

$$\Sigma_{\xi}^{T=0} = -i\xi e^2 \int \frac{d^4 p}{(2\pi)^4} \frac{1}{(p^2 + i\epsilon)^2} \left[\not{p} - \frac{p^2 \not{k} + k^2 \not{p}}{(p+k)^2 + i\epsilon} \right]. \quad (6.20)$$

The odd $\not{p}$ term vanishes on integration. Evaluation of the rest leads to

$$a_{\xi}^{T=0} = -\frac{\xi e^2}{(4\pi)^2} \left[\frac{2}{\epsilon'} - \gamma - 1 + \ln(4\pi) - \int_0^1 dx \, \ln(x(1-x)k^2) + O(\epsilon') \right]. \quad (6.21)$$

Finally adding the counter-term to $a^{T=0}$, as fixed by the renormalization condition

$$a_{\text{ren}}^{T=0}(k^2 = \sigma^2) = 0, \quad (6.22)$$

we arrive at

$$a_{\text{ren}}^{T=0}(k^2) = (1 + \xi) \frac{e^2}{(4\pi)^2} \ln \frac{k^2}{\sigma^2}. \quad (6.23)$$

where σ is the renormalization scale.

We now turn to the T -dependent part, and write

$$\Sigma' = \Sigma'_{\text{FG}} + \Sigma'_{\xi}. \quad (6.24)$$

To obtain Σ'_{FG} , one has to put Eqs. (6.10) and (6.13) in the expression (4.93) for the self-energy, and consider the relevant terms. It is then convenient to change p to $-p - K$ in the f_F -containing term. Finally setting $p^2 = 0$, as allowed by $\delta(p^2)$, yields

$$\Sigma'_{\text{FG}} = 2e^2 \int \frac{d^4 p}{(2\pi)^4} [(\not{p} + \not{k})f_B(p) + \not{p}f_F(p)] 2\pi\delta(p^2) \text{Re} \frac{1}{2p \cdot k + k^2 + i\epsilon}. \quad (6.25)$$

Putting (6.10) and (6.14) in (4.93), one arrives at Σ'_{ξ} . Its f_F -containing part and f_B -containing part will be indicated explicitly:

$$\Sigma'_{\xi} = \Sigma'_{\xi,F} + \Sigma'_{\xi,B}. \quad (6.26)$$

In $\Sigma'_{\xi,F}$, changing p to $-p - K$ and then setting $p^2 = 0$ (allowed by the delta function) gives us

$$\Sigma'_{\xi,F} = \xi e^2 \int \frac{d^4 p}{(2\pi)^4} [k^2 \not{p} - 2p \cdot k \not{k}] 2\pi\delta(p^2) f_F(p) \text{Re} \frac{1}{(2p \cdot k + k^2 + i\epsilon)^2}. \quad (6.27)$$

In $\Sigma'_{\xi,B}$, simplification leads to

$$\Sigma'_{\xi,B} = \xi e^2 \int \frac{d^4 p}{(2\pi)^3} f_B(p) \frac{d\delta(p^2)}{dp^2} \left[\not{p} - (p^2 \not{k} + k^2 \not{p}) \operatorname{Re} \frac{1}{(p+k)^2 + i\epsilon} \right]. \quad (6.28)$$

The $\not{p}$ term, being odd, drops out on integration. To deal with the remaining part, we first write down

$$\frac{d\delta(p^2)}{dp^2} = \lim_{\epsilon \rightarrow 0} \frac{1}{2\pi i} \left[\frac{1}{(p^2 + i\epsilon)^2} - \frac{1}{(p^2 - i\epsilon)^2} \right] \quad (6.29)$$

and then use the regularization

$$\frac{1}{(p^2 \pm i\epsilon)^2} = \lim_{\lambda \rightarrow 0} \frac{\partial}{\partial \lambda^2} \frac{1}{p^2 - \lambda^2 \pm i\epsilon} \quad (6.30)$$

so that

$$\frac{d\delta(p^2)}{dp^2} = - \lim_{\lambda \rightarrow 0} \frac{\partial}{\partial \lambda^2} \delta(p^2 - \lambda^2). \quad (6.31)$$

Let us now make use of Eq. (6.31) in Eq. (6.28). We then commute the integration over p_0 with the limit and the differentiation involving λ , and set $p^2 = \lambda^2$ (allowed by the delta function). This gives

$$\begin{aligned} \Sigma'_{\xi,B} &= \xi e^2 \int \frac{d^3 P}{(2\pi)^3} \lim_{\lambda \rightarrow 0} \frac{\partial}{\partial \lambda^2} \int dp_0 f_B(p) \delta(p^2 - \lambda^2) (k^2 \not{p} + \lambda^2 \not{k}) \\ &\quad \times \operatorname{Re} \frac{1}{k^2 + 2p \cdot k + \lambda^2 + i\epsilon}. \end{aligned} \quad (6.32)$$

After integrating over p_0 , the operations involving λ are carried out. While this last step is easily performed for the part proportional to λ^2 (since

$$\lim_{\lambda \rightarrow 0} \frac{\partial}{\partial \lambda^2} (\lambda^2 f(\lambda^2)) = f(0) \quad (6.33)$$

with f denoting a function with finite $\partial f / \partial \lambda^2$ at $\lambda = 0$ [61]), a more tedious algebra is to be worked out for the remaining part, finally giving

$$\begin{aligned} \Sigma'_{\xi,B} &= \xi \frac{e^2}{2} \int \frac{d^3 P}{(2\pi)^3} \frac{1}{e^{\beta P} - 1} \frac{1}{P} \left[\left(\not{k} + \frac{k^2 \vec{P} \cdot \vec{\gamma}}{2P^2} \right) \operatorname{Re} \left(\frac{1}{D_+} + \frac{1}{D_-} \right) \right. \\ &\quad \left. - k^2 P \gamma_0 \operatorname{Re} \left(\frac{1 + k_0/P}{D_+^2} - \frac{1 - k_0/P}{D_-^2} \right) + k^2 \vec{P} \cdot \vec{\gamma} \operatorname{Re} \left(\frac{1 + k_0/P}{D_+^2} + \frac{1 - k_0/P}{D_-^2} \right) \right] \\ &\quad - \xi \frac{e^2 \beta}{4} \int \frac{d^3 P}{(2\pi)^3} \frac{e^{\beta P}}{(e^{\beta P} - 1)^2} \frac{1}{P^2} \left[k^2 P \gamma_0 \operatorname{Re} \left(\frac{1}{D_+} - \frac{1}{D_-} \right) \right. \\ &\quad \left. - k^2 \vec{P} \cdot \vec{\gamma} \operatorname{Re} \left(\frac{1}{D_+} + \frac{1}{D_-} \right) \right]. \end{aligned} \quad (6.34)$$

Here

$$D_{\pm} = k^2 \pm 2Pk_0 - 2\vec{P} \cdot \vec{K} + i\epsilon. \quad (6.35)$$

While the expressions (6.25) and (6.27) are the same as those in Ref. [5] (except that we have carefully incorporated the $i\epsilon$, which will be needed for the calculations in Appendix F), the expression (6.34) is a different one. We have found this form of $\Sigma'_{\xi,B}$ convenient for actually carrying out the integration (this, again, is done in Appendix F).

Let us now investigate the high- T behaviour of (6.25), (6.27) and (6.34). Following Ref. [5], this behaviour can be inferred from the degree of ultraviolet divergence of the integral in each expression in the absence of the cut-off of $O(T)$ provided by f_B or f_F or $e^{\beta P}/(e^{\beta P} - 1)^2$. One then finds that Σ'_{FG} goes like T^2 , while $\Sigma'_{\xi,F}$ and $\Sigma'_{\xi,B}$ go like T . It may be noted that one cannot obtain the correct high- T behaviour of $\Sigma'_{\xi,B}$ without removing the regulator λ , and this can be done only after performing the p_0 -integration, as in (6.34).

At high T , one can neglect the renormalized $\Sigma^{T=0}$ compared to Σ' [46]. Actually $\Sigma^{T=0}$ depends on the renormalization scale σ , but since the dependence is logarithmic, we can still ignore it vis-a-vis the power law dependences on T in the various parts of Σ' . We shall, however, see that it is *not necessary* to neglect $\Sigma^{T=0}$ for proving gauge independence at $K \gg eT$. We shall also use (6.23) to incorporate the σ -dependence in the equation for the effective mass.

6.3 Equation governing gauge dependence of dispersion relation at one loop, and general considerations

Inverting Eq. (6.1), one obtains

$$a = \frac{1}{4K^2} [-\text{Tr}(\not{K}\Sigma) + \omega \text{Tr}(\not{p}\Sigma)], \quad (6.36)$$

$$b = \frac{1}{4K^2} [\omega \text{Tr}(\not{K}\Sigma) - (\omega^2 - K^2) \text{Tr}(\not{p}\Sigma)]. \quad (6.37)$$

The dispersion relation is obtained by putting a , b in Eq. (6.4). Let us now write the function f in Eq. (6.4) as

$$f = f_{\text{FG}} + f_{\xi}, \quad (6.38)$$

$$f_{\text{FG}} = (\omega - K)(1 - a_{\text{FG}}) - b_{\text{FG}}, \quad (6.39)$$

$$f_{\xi} = -(\omega - K)a_{\xi} - b_{\xi}. \quad (6.40)$$

Next using Eqs. (6.26), (6.27) and (6.34), one readily sees that k^2 factors out from the expression for $\text{Tr}(k\Sigma'_\xi)$. Therefore in view of Eq. (6.3), one can write from Eq. (6.37) that

$$b_\xi = (\omega - K)(\text{1-loop function of } \omega, K) \quad (6.41)$$

(note that b , being zero at $T = 0$, is determined by the Σ' part only). Consequently

$$f_\xi = -(\omega - K)[a_\xi + (\text{1-loop function of } \omega, K)]. \quad (6.42)$$

This is the equation governing the gauge-dependence of the electron dispersion relation at one-loop. This equation is of the form

$$(f_\xi)_{\text{1-loop}} = f_{\text{tree}} \times [\text{1-loop } \xi\text{-dependent function}]. \quad (6.43)$$

As we shall see in the next section, *the fact that f_{tree} factors out on the R.H.S. of Eq. (6.43) is crucial to the proof of gauge independence.*

Ref. [24] contains a general, nonperturbative derivation of the identities determining the gauge dependence of the gluon dispersion relations (see Eqs. (16) and (17) of [24]). To one loop these gauge dependence identities are shown to reduce to relations (see Eq. (20) of [24]) which are analogous to Eq. (6.43) above. *It is therefore likely that a general gauge dependence identity for the electron dispersion relation, reducing to Eq. (6.43) at one loop, can be derived and used to arrive at a general proof of gauge independence of the dispersion relation* (as in the gluon case). In this paper, however, we confine ourselves to one-loop calculations.

6.4 Gauge independence of one-loop dispersion relation at momenta much larger than eT

Since we shall consider $K \gg eT$ in this section, let us first note that for *sufficiently small* e there always exists a domain $eT \ll K \ll T$, so that the restriction $K \gg eT$ does not contradict the high- T approximation.

The basic premise for the considerations of this section is that when the finite-temperature effects are small, the dispersion relation should involve a deviation from $\omega = K$ primarily by powers of e^2 . We have used the phrase “primarily” because the terms in the deviation may come multiplied with powers of $\ln(1/e)$ (originating from powers of $\ln(\omega - K)$; see, for example, Eqs. (6.5) and (6.6)). Now, to first order in e^2 , Eq.

(6.6) becomes $\omega - K = e^2 T^2 / 8K$. This, when used in the remaining term on the R.H.S. of Eq. (6.6), results in [5]

$$\omega = K + \frac{M_0^2}{K} - \frac{M_0^4}{2K^3} \ln \frac{2K^2}{M_0^2} + \dots \quad (6.44)$$

where $M_0^2 = e^2 T^2 / 8$. The terms on the R.H.S. of the above expansion clearly indicate that the expansion is valid for $K \gg M_0$ i.e. $K \gg eT$. This can be understood in the following way. As shown in Ref. [5] and as we shall see in Sec. 6.5.3, M_0 is the leading-order effective electron mass [62] and thus can be considered a measure of finite-temperature effects. So, $K \gg eT$ ensures that the finite-temperature *effects* are small (even though T is large), and only then the expansion in e^2 given by Eq. (6.44) is valid. (As we will see in Sec. 6.5.1, such an expansion does not exist for $K \ll eT$.)

Now we turn to gauge independence. The expected general form of the one-loop dispersion relation for $K \gg eT$ *when the terms subleading in T are kept* is

$$\omega = K + e^2 f_1(e, K, T) \quad (6.45)$$

where the e -dependence of f_1 involves only powers of $\ln(1/e)$. Equation (6.44) (without the last term on its R.H.S., which is actually of order $e^4 \ln(1/e)$) is a special case of Eq. (6.45), where f_1 turned out to be totally e -independent. Let us now take Eq. (6.45) to be the relation in the Feynman gauge i.e. assume that it satisfies $f_{\text{FG}} = 0$. Then to prove that (6.45) is gauge independent we have to show that it also satisfies $f_{\text{FG}} + f_\xi = 0$. It suffices to show that, at (6.45), f_ξ is of order e^4 (with or without powers of $\ln(1/e)$) [63]. This follows readily from Eq. (6.42), since the portion of the R.H.S. of Eq. (6.42) within the square brackets involves terms of the order of e^2 and $e^2(\ln(1/e))^n$ (n is an integer), and so also does $(\omega - K)$ at (6.45).

Digressing briefly from *high-temperature* QED, we mention that *the above demonstration of gauge independence is actually sufficient also for the case which involves not large T but large momenta*. Thus for $K \gg T$, a one-loop relation of the key form (6.45) should still hold. Note that the rest of the proof, namely, arriving at Eq. (6.42), did not assume large T (in particular, we did not neglect the vacuum contribution to a). The proof of gauge independence continues to be valid if, in addition, we have a chemical potential μ such that $K \gg \mu$ as well. Then the relevant changes are that f_1 in (6.45) depends on μ as well, f_F is modified and we have $f_F(-p)$ in place of $f_F(p)$ in (6.27), but none of these affect the proof outlined above.

6.5 Gauge independence of effective mass at one-loop

6.5.1 Subtleties at zero momentum

The effective electron mass is the value of ω at $K = 0$. (Here we are following the nomenclature used in much of the literature on high-temperature quantum field theory, including Ref. [5], though this quantity has been called the inertial mass earlier in this work.) The analysis of the previous section does not prove the gauge independence of the effective mass because the form (6.45) does not hold near $K = 0$. For example, to leading order in T , the one-loop dispersion relation for $K \ll eT$ is [5]

$$\omega = M_0 + \frac{K}{3} + \frac{K^2}{3M_0} + \dots \quad (6.46)$$

(This is obtained by substituting in Eq. (6.6) a trial solution of the form (6.53) and determining the coefficients.) Eq. (6.46) suggests that *the leading order values of a and b are not of $O(e^2)$ (with or without powers of $\ln(1/e)$) in the $K \ll eT$ limit*; indeed the values at $K = 0$ are $a = -1/3$ and $b = -2M_0/3$ [64]. This apparently surprising behaviour can be understood in a simple way, without performing a detailed calculation, as follows.

At leading order a and b are $e^2 T^2$ times some function of ω and K (T^2 being deduced from ultraviolet power counting in Σ'). Therefore, to have the correct dimensions, $a_{K=0} \sim e^2 T^2 / M_0^2$ and $b_{K=0} \sim e^2 T^2 / M_0$ (remembering $M_0 = \omega_{K=0}$ at leading order). Putting these in the dispersion relation (6.4) at $K = 0$, namely,

$$M_0 - M_0 a_{K=0} - b_{K=0} = 0, \quad (6.47)$$

gives us $M_0 \sim eT$. Using this, it follows that $a_{K=0}$ is of $O(1)$ and $b_{K=0}$ is of $O(eT)$.

6.5.2 Formula for the effective mass

The traces $\text{Tr}(\not{k}\Sigma)$ and $\text{Tr}(\not{p}\Sigma)$ are functions of ω and K . The expressions for them are obtained from Eqs. (6.18), (6.23), (6.25), (6.27) and (6.34). First of all we show that these traces are both even functions of K . For the $\Sigma^{T=0}$ part, this is obvious from (6.3). For the Σ' part, first note that k^μ can occur in the expressions for the traces only through $k \cdot p$, $k \cdot v$ and k^2 (p^μ being the integration variable). Next, let us go to the rest frame of the medium (in which K is the magnitude of the three-momentum) and note that $\vec{K}$ can now occur in the expressions for the traces only through $\vec{K} \cdot \vec{P}$ and K^2 . Odd power of

K can come from $\vec{K} \cdot \vec{P} = KP \cos \theta$ where θ is the angle between $\vec{K}$ and $\vec{P}$. This is also the only place where θ occurs in the integrand. So changing θ to $\pi - \theta$ together with the change $K \rightarrow -K$, we establish the even nature as a function of K .

For small K , therefore, we can write

$$\frac{1}{4} \text{Tr}(\not{k} \Sigma) = h_0 + h_1 K^2 + h_2 K^4 + \dots \quad (6.48)$$

$$\frac{1}{4} \text{Tr}(\not{p} \Sigma) = g_0 + g_1 K^2 + g_2 K^4 + \dots \quad (6.49)$$

where h_i and g_i are functions of ω . Substitution of the above expressions into Eqs. (6.36) and (6.37) gives

$$a = -\frac{1}{K^2}(h_0 - \omega g_0) - (h_1 - \omega g_1) + O(K^2), \quad (6.50)$$

$$b = \frac{\omega}{K^2}(h_0 - \omega g_0) + (\omega h_1 - \omega^2 g_1 + g_0) + O(K^2). \quad (6.51)$$

For the form factors a and b to remain analytic at $K = 0$, the relation

$$h_0 = \omega g_0 \quad (6.52)$$

must hold. We now put (6.50) and (6.51), subject to the constraint (6.52), in the dispersion relation (6.4). Then we use the fact that for any small K , the solution

$$\omega = M + \omega_1 K + \omega_2 K^2 + \dots \quad (6.53)$$

where M and ω_i are constants, must satisfy the dispersion relation. We also expand h_i and g_i in K by first doing a Taylor expansion of them around $\omega = M$ as functions of ω , and then putting (6.53). Thus

$$h_i(\omega) = h_i(M) + h'_i(M)\omega_1 K + O(K^2) \quad (6.54)$$

with a similar equation for g_i . The equation that now results from the dispersion relation (6.4) has a series in powers of K alone on the L.H.S. Equating the coefficient of each power of K separately to zero will determine the constants M and ω_i of (6.53). Thus, equating the constant term to zero gives

$$M = g_0(M). \quad (6.55)$$

Multiplying both sides by M , we make use of

$$h_0(M) = M g_0(M) \quad (6.56)$$

(which follows from (6.52)) to finally arrive at

$$M^2 = h_0(M). \quad (6.57)$$

One can use either (6.57) or (6.55) as the formula for the effective mass. We shall use (6.57), which, by using Eq. (6.48), can be written as [65]:

$$M^2 = \frac{1}{4} \lim_{K \rightarrow 0, \omega \rightarrow M} \text{Tr}[\not{K}\Sigma]. \quad (6.58)$$

We comment that one should not attempt to compare this formula with the formula for the inertial mass of Sec. 4.1, since the latter was deduced by assuming a straightforward expansion in e^2 and neglecting various terms accordingly.

6.5.3 Investigation of gauge independence

Let us now substitute the expression for the self-energy in Eq. (6.58). On using Eqs. (6.17) and (6.18) in Eq. (6.58), we obtain

$$M^2 = M^2 a^{T=0}(M^2) + M'^2, \quad (6.59)$$

$$M'^2 \equiv \frac{1}{4} \lim_{K \rightarrow 0, \omega \rightarrow M} \text{Tr}[\not{K}\Sigma']. \quad (6.60)$$

At $T = 0$, $M' = 0$ and so Eq. (6.59) is correctly satisfied by $M = 0$.

Putting in Eq. (6.60) the expressions for Σ'_{FG} , $\Sigma'_{\xi,F}$ and $\Sigma'_{\xi,B}$ given by Eqs. (6.25), (6.27) and (6.34), and noting that the limit $\omega \rightarrow M$ translates to $k_0 \rightarrow M$ in the rest-frame of the medium, we obtain

$$M'^2 = M'^2_{\text{FG}} + M'^2_{\xi,F} + M'^2_{\xi,B}, \quad (6.61)$$

$$M'^2_{\text{FG}} = 2e^2 \int \frac{d^4 p}{(2\pi)^3} [(p_0 + M)f_B(p) + p_0 f_F(p)] \delta(p^2) \text{Re} \frac{1}{2p_0 + M + i\frac{\epsilon}{M}}, \quad (6.62)$$

$$M'^2_{\xi,F} = -\xi e^2 M \int \frac{d^4 p}{(2\pi)^3} p_0 f_F(p) \delta(p^2) \text{Re} \frac{1}{(2p_0 + M + i\frac{\epsilon}{M})^2}, \quad (6.63)$$

$$M'^2_{\xi,B} = M'^2_{\xi,B(\text{I})} + M'^2_{\xi,B(\text{II})} + M'^2_{\xi,B(\text{III})}, \quad (6.64)$$

$$M'^2_{\xi,B(\text{I})} = \frac{\xi e^2 M}{2} \int \frac{d^3 P}{(2\pi)^3} \frac{1}{P} \frac{1}{e^{\beta P} - 1} \text{Re} \left(\frac{1}{E_+} + \frac{1}{E_-} \right), \quad (6.65)$$

$$M'^2_{\xi,B(\text{II})} = -\frac{\xi e^2 M}{2} \int \frac{d^3 P}{(2\pi)^3} \frac{1}{e^{\beta P} - 1} \text{Re} \left[\frac{M}{P} \left(\frac{1}{E_+^2} + \frac{1}{E_-^2} \right) + \left(\frac{1}{E_+^2} - \frac{1}{E_-^2} \right) \right], \quad (6.66)$$

$$M'^2_{\xi,B(\text{III})} = -\frac{\xi e^2 M^2 \beta}{4} \int \frac{d^3 P}{(2\pi)^3} \frac{1}{P} \frac{e^{\beta P}}{(e^{\beta P} - 1)^2} \text{Re} \left(\frac{1}{E_+} - \frac{1}{E_-} \right). \quad (6.67)$$

Here

$$E_{\pm} = M \pm 2P + \frac{i\epsilon}{M}. \quad (6.68)$$

We now investigate the high- T behaviour of the three terms on the R.H.S. of Eq. (6.61). This can be arrived at from the ultraviolet behaviour, as explained towards the end of Sec. 6.2. However there is a constraint that each term on the R.H.S. of Eq. (6.61) is an even function of M . (This can be seen by changing p_0 to $-p_0$ together with the change $M \rightarrow -M$ in (6.62), in (6.63) and in

$$M'_{\xi,B}{}^2 = -\xi e^2 M^2 \int \frac{d^4 p}{(2\pi)^3} (p^2 + p_0 M) f_B(p) \frac{d\delta(p^2)}{dp^2} \text{Re} \frac{1}{p^2 + 2p_0 M + M^2 + i\epsilon}, \quad (6.69)$$

the last equation having been obtained by putting (6.28) in (6.60).) These considerations, plus dimensional analysis, tell us that at high T , $M'_{\text{FG}}{}^2$ goes like T^2 , $M'_{\xi,F}{}^2$ like $M^2(\ln |T/M|)^{n_1}$ and $M'_{\xi,B}{}^2$ like $M^2(\ln |T/M|)^{n_2}$ (n_1, n_2 being positive integers). Since after T^2 , the next allowed term in $M'_{\text{FG}}{}^2$ is $M^2(\ln |T/M|)^{n_3}$ (n_3 a positive integer), the general expression for M'^2 at high T can be written as

$$M'^2 = e^2 \left[c_0 T^2 + c_3 M^2 \left(\ln \frac{T}{M} \right)^{n_3} + \xi M^2 \left(c_1 \left(\ln \frac{T}{M} \right)^{n_1} + c_2 \left(\ln \frac{T}{M} \right)^{n_2} \right) \right] \quad (6.70)$$

where c_0, c_1, c_2, c_3 are constants, and we have dropped the modulus of the arguments of the logarithms since T/M is positive. Terms independent of T have been neglected on the R.H.S. of Eq. (6.70).

In principle there could be further constraints ruling out some term(s) in Eq. (6.70), but a detailed calculation, described in Appendix F, reveals that all these terms are indeed present and that $n_1 = 1 = n_2 = n_3$. Thus we actually have the high temperature equation

$$M'^2 = e^2 \left[\frac{T^2}{8} + \frac{M^2}{8\pi^2} \ln \frac{T}{M} + \xi \frac{M^2}{8\pi^2} \ln \frac{T}{M} \right]. \quad (6.71)$$

The detailed calculation may also be viewed as a check on the argument involving the degree of ultraviolet divergence mentioned before.

Using Eqs. (6.23) and (6.71) in Eq. (6.59), we then have

$$M^2 = e^2 \left[\frac{T^2}{8} + (1 + \xi) \frac{M^2}{8\pi^2} \ln \frac{T}{\sigma} \right]. \quad (6.72)$$

On the R.H.S. of Eq. (6.72) (as also in Eq. (6.71)), we have not given the term $\sim e^2 M^2$; all other terms contributing to the R.H.S. of Eq. (6.72) (and (6.71)) vanish in the limit of

large T/M . It is interesting to note that in Eq. (6.72), the $\ln(M/\sigma)$ term from $\Sigma^{T=0}$ and the $\ln(T/M)$ term from Σ' have exactly combined to yield just a $\ln(T/\sigma)$ term, because it has been observed that similar combination also takes place in the case of the gauge boson self-energy in the Yang-Mills theory [66]. A discussion of similar behaviour in the case of three-point function, and, in general, N -point function of gauge boson in the Yang-Mills theory, is to be found in Ref. [67].

The first important observation from Eq. (6.72) is that $M^2 = e^2 T^2/8$ to leading order in T , which, of course, is a well-known result [5]. Now, the gauge dependent part of Σ' goes like T . Therefore in view of Eq. (6.58), one would expect an $e^2 \xi MT$ term in Eq. (6.72). The absence of such a term shows that *the part of Σ'_ξ leading in T does not contribute to M .*

To quantify the effect of this absence let us consider the equation

$$M^2 = e^2 \left(\frac{T^2}{8} + cMT \right) \quad (6.73)$$

where c is a constant. To leading order in T , the second term on the R.H.S. is negligible, so that $M = O(eT)$. This can now be used as an approximation in the second term, to give

$$M^2 = e^2 \frac{T^2}{8} [1 + O(e)] \quad (6.74)$$

showing that there is a correction of $O(e^2 T)$ to M . (This conclusion can also be arrived at by solving Eq. (6.73) for M .) So the absence of the $e^2 MT$ term means that *there is no $O(e^2 T)$ correction to the effective mass in any linear covariant gauge.* It may be noted that the consequences of the presence or absence of various *powers* of T in Eq. (6.72) are to be taken seriously despite the presence of the scale σ , since the σ dependence is only logarithmic and is not multiplied with any power of T .

Finally, (6.72) shows that only the subleading $\ln T$ dependence of Σ'_ξ contributes to M . Using the leading order result $M = O(eT)$ as before to approximate the remaining terms on the R.H.S. of Eq. (6.72), we infer that the ξ -dependence in Eq. (6.72) is $O(e^4 T^2)$, apart from the logarithm. As one certainly expects $O(e^4 T^2)$ contribution to Eq. (6.72) from two loops, *it is possible that the ξ -dependence which we have obtained will be canceled by two-loop contribution.* But it seems that this issue can be decided only by an actual two-loop calculation. It is however clear that an $e^2 \xi MT$ term, being of $O(e^3 T^2)$, was less likely to be canceled by two-loop contribution. This shows the significance of the absence of such a term.

CHAPTER 7

Conclusions

In this thesis, we studied some aspects of particle physics in the presence of a background medium, using the real-time formulation of thermal field theory. We studied the $O(e^2)$ corrections to the gravitational interactions of a charged lepton in the presence of a matter background. We also investigated gauge independence of one-loop electron dispersion relation in high-temperature QED. These calculations complement and extend previous calculations along similar lines, in various useful ways. The usefulness, as we will explain, is both theoretical and phenomenological.

We first discuss the importance of the various aspects of our work from a theoretical point of view. The calculation of gravitational couplings in a medium, as presented by us, employed several thermal field theoretic techniques that can be useful in other contexts also. For example, we made explicit use of only the 1PI diagrams in our calculation of the gravitational mass. The effect of the one-particle reducible diagrams were taken into account by choosing each external spinor to be the solution of the effective Dirac equation in a medium, and multiplying the spinor with the square-root of the wavefunction renormalization factor. It was then seen that the wavefunction renormalization factors are instrumental in cancelling an infrared divergent contribution in the 1PI gravitational vertex function.

Again, we dealt extensively with the limiting case of the finite-temperature gravitational vertex function in which the graviton carries zero momentum. The ambiguity of the finite-temperature Green's functions evaluated at zero momentum is a well-known problem in thermal field theory. This property is usually due to the fact that the different mathematical limits correspond to different physical situations, so that the resolution of the apparent paradox lies in recognizing the appropriate correspondence with the physical situation at hand. In our case also, the 1PI gravitational vertex at zero graviton momentum turned out to contain contributions which superficially seemed to be ill-defined, and

the problem was resolved by taking the relevant physical situation into account. Thus, taking the various limits in the precise order indicated in the operational formula for the gravitational mass, given by Eq. (5.25), we obtained unique and well-defined expressions for all the contributions. We also reproduced these expressions by using the special regularization technique given in Eq. (E.50).

On the other hand, our work on high-temperature QED gives credence to the idea that any gauge dependence present in the self-energy should not show up in the dispersion relation. This idea motivated us to focus on the subleading part of the self-energy of the electron at one-loop, which (unlike the leading, $O(T^2)$ part) depends on the choice of gauge.

We obtained an equation which governs the gauge dependence of the one-loop dispersion relation, stressed its analogy with the corresponding equation in the gluon case, and consequently pointed out the possibility of a generalization of our equation to all orders. From this equation obtained by us, the gauge independence in the $K \gg eT$ limit followed in a straightforward way. We then showed that the effective mass is not affected by the leading part (going like T) of the gauge-dependent part of the self-energy and hence does not receive $O(e^2T)$ correction in any gauge. While the effective mass was found to be influenced by the subleading part (going like $\ln T$) of the gauge-dependent part of the self-energy, it is possible that this will be canceled by two-loop contribution. Exploration of this possibility can be a problem for further investigation.

Our calculations and results are also important from a phenomenological point of view. We found that the contribution to the gravitational mass of the electron from a matter background with a non-zero chemical potential, such as the Sun, can be quite appreciable. An interesting problem, combining the various aspects covered in this thesis, would be the study of the gravitational interactions of charged leptons in a background at high temperature. The results can then be estimated for the case of a supernova, which is an example of such a background.

Moreover, the matter-induced corrections to the gravitational mass are different for the various charged lepton flavors, and are not the same for the corresponding antiparticles. There are situations in which mass differences, intrinsic or induced, have important physical implications, such as the neutron-proton mass difference in the context of the nucleosynthesis calculations in the Early Universe. Although our work has focused in the case of the charged leptons, similar considerations can be applied to the other fermions

as well. Our calculations have provided a necessary ingredient for being able to consider them in a systematic manner, and set the stage for their further study on a firm basis.

APPENDIX A

Field quantization using discrete momenta

In this Appendix, we relate the quantization with discrete momenta, used in Sec. 2.2, to the quantization with continuous momenta, used in Sec. 2.3 onward [68].

Beginning with the complex scalar field, we consider the Fourier expansion given in Eq. (2.19). The equal-time commutation relations then force

$$[a(\vec{P}), a^\dagger(\vec{P}')]_- = \delta^{(3)}(\vec{P} - \vec{P}'), \quad [b(\vec{P}), b^\dagger(\vec{P}')]_- = \delta^{(3)}(\vec{P} - \vec{P}'), \quad (\text{A.1})$$

all other commutators being zero. Also, Eq. (2.19) leads to

$$\begin{aligned} H &= \int d^3P E_P \left(a^\dagger(\vec{P})a(\vec{P}) + b^\dagger(\vec{P})b(\vec{P}) \right), \\ Q &= \int d^3P \left(a^\dagger(\vec{P})a(\vec{P}) - b^\dagger(\vec{P})b(\vec{P}) \right). \end{aligned} \quad (\text{A.2})$$

If we instead use quantization in a volume V , the plane-wave states can only possess the discrete momenta

$$\vec{P} = \frac{2\pi}{V^{1/3}} \vec{n}, \quad (\text{A.3})$$

where n_1, n_2, n_3 are integers. Integrals are then replaced by sums

$$\int \frac{d^3P}{(2\pi)^3} \rightarrow \frac{1}{V} \sum_{\vec{P}} \quad (\text{A.4})$$

and delta functions by Kronecker symbols

$$(2\pi)^3 \delta^{(3)}(\vec{P} - \vec{P}') \rightarrow V \delta_{\vec{P}, \vec{P}'} . \quad (\text{A.5})$$

Note that the left-hand sides of Eqs. (A.4) and (A.5) taken together give unity, and so also do the right-hand sides.

Let us now redefine the ladder operators through

$$a(\vec{P}) = \sqrt{\frac{V}{(2\pi)^3}} a_{\vec{P}}, \quad b(\vec{P}) = \sqrt{\frac{V}{(2\pi)^3}} b_{\vec{P}}. \quad (\text{A.6})$$

Eqs. (A.1), (A.5) and (A.6) give

$$[a_{\vec{P}}, a_{\vec{P}'}]_- = \delta_{\vec{P}, \vec{P}'}, \quad [b_{\vec{P}}, b_{\vec{P}'}]_- = \delta_{\vec{P}, \vec{P}'}, \quad (\text{A.7})$$

while Eqs. (A.2), (A.4) and (A.6) give

$$\begin{aligned} H &= \sum_{\vec{P}} E_P (a_{\vec{P}}^\dagger a_{\vec{P}} + b_{\vec{P}}^\dagger b_{\vec{P}}), \\ Q &= \sum_{\vec{P}} (a_{\vec{P}}^\dagger a_{\vec{P}} - b_{\vec{P}}^\dagger b_{\vec{P}}). \end{aligned} \quad (\text{A.8})$$

(Alternatively, one can use Eqs. (2.19), (A.4) and (A.6) to arrive at

$$\phi(x) = \sum_{\vec{P}} \frac{1}{\sqrt{2E_P V}} (a_{\vec{P}} e^{-ip \cdot x} + b_{\vec{P}}^\dagger e^{ip \cdot x}), \quad (\text{A.9})$$

the discretized Fourier expansion for the field. Eqs. (A.7) and (A.8) then follow from this expansion.) Eqs. (A.7) and (A.8) are the key expressions used in Sec. 2.2.

Turning next to the fermion field, we consider the Fourier expansion of Eq. (2.37) and the spinor normalization given in Eq. (2.38). The equal-time canonical anticommutation relations then force

$$[c_s(\vec{P}), c_{s'}^\dagger(\vec{P}')]_+ = \delta_{ss'} \delta^{(3)}(\vec{P} - \vec{P}'), \quad [d_s(\vec{P}), d_{s'}^\dagger(\vec{P}')]_+ = \delta_{ss'} \delta^{(3)}(\vec{P} - \vec{P}'), \quad (\text{A.10})$$

all other anticommutators being zero. (In more general terms, the choice $N_1 = 1 = N_2$ forces $N_3 = 1$ in Eqs. (B.16) – (B.18).) Also, Eq. (2.37) leads to

$$\begin{aligned} H &= \int d^3P E_P \sum_s (c_s^\dagger(\vec{P}) c_s(\vec{P}) + d_s^\dagger(\vec{P}) d_s(\vec{P})), \\ Q &= \int d^3P \sum_s (c_s^\dagger(\vec{P}) c_s(\vec{P}) - d_s^\dagger(\vec{P}) d_s(\vec{P})). \end{aligned} \quad (\text{A.11})$$

Use of Eqs. (A.4) and (A.5), and redefinitions similar to those in Eq. (A.6) then give the fermionic counterparts of Eqs. (A.7) and (A.8), which were used in Sec. 2.2.

APPENDIX B

Full propagator and wavefunction renormalization factor in vacuum

We start with the definition of the full fermion propagator in vacuum

$$iS'_{f\alpha\beta}(x) \equiv \langle 0|T[\psi_\alpha(x)\bar{\psi}_\beta(0)]|0\rangle \quad (\text{B.1})$$

$$= \theta(x_0)\langle 0|\psi_\alpha(x)\bar{\psi}_\beta(0)|0\rangle - \theta(-x_0)\langle 0|\bar{\psi}_\beta(0)\psi_\alpha(x)|0\rangle \quad (\text{B.2})$$

where $\psi(x)$ is the Heisenberg picture field, and $|0\rangle$ the vacuum of the full interacting theory. The wavefunction renormalization factor Z is defined through

$$\langle 0|\psi_\alpha(0)|f(\vec{P}, s)\rangle = \sqrt{Z} u_{s\alpha}(\vec{P}), \quad (\text{B.3})$$

where the 1-particle state $|f(\vec{P}, s)\rangle$ is again an eigenstate of the full interacting hamiltonian. The tree-level spinor $u_s(\vec{P})$ is actually the analogue of $U_s(\vec{P})$ (see Eq. (4.26)), since the sum of the bare mass and the self-energy correction in vacuum is set equal to the physical mass of the fermion.

Eq. (B.3) implies that

$$\langle 0|\psi_\alpha(x)|f(\vec{P}, s)\rangle = \sqrt{Z} u_{s\alpha}(\vec{P})e^{-ip\cdot x}, \quad (\text{B.4})$$

with $p_0 = E_P \equiv \sqrt{P^2 + m_f^2}$. (We used an identity similar to the one in Eq. (5.6).) It then follows that

$$\langle f(\vec{P}, s)|\bar{\psi}_\alpha(x)|0\rangle = \sqrt{Z} \bar{u}_{s\alpha}(\vec{P})e^{ip\cdot x}. \quad (\text{B.5})$$

Similar relations involving the 1-antiparticle state are $|\bar{f}(\vec{P}, s)\rangle$ are

$$\langle \bar{f}(\vec{P}, s)|\psi_\alpha(x)|0\rangle = \sqrt{Z} v_{s\alpha}(\vec{P})e^{ip\cdot x}, \quad (\text{B.6})$$

$$\langle 0|\bar{\psi}_\alpha(x)|\bar{f}(\vec{P}, s)\rangle = \sqrt{Z} \bar{v}_{s\alpha}(\vec{P})e^{-ip\cdot x}. \quad (\text{B.7})$$

In Eqs. (B.6) and (B.7) also, $p_0 = E_P$. It may be noted that in vacuum, Z is independent of momentum and also the same for the fermion and the antifermion, properties which may not hold in a medium.

We shall take the normalization of 1-particle states to be

$$\langle f(\vec{P}', s') | f(\vec{P}, s) \rangle = (2\pi)^3 \delta^{(3)}(\vec{P} - \vec{P}') \delta_{s, s'}, \quad (\text{B.8})$$

with a similar normalization for $|\bar{f}(\vec{P}, s)\rangle$ as well. The completeness relation is then given by

$$1 = |0\rangle\langle 0| + \sum_s \int \frac{d^3 P}{(2\pi)^3} |f(\vec{P}, s)\rangle\langle f(\vec{P}, s)| + \sum_s \int \frac{d^3 P}{(2\pi)^3} |\bar{f}(\vec{P}, s)\rangle\langle \bar{f}(\vec{P}, s)| + \dots, \quad (\text{B.9})$$

where the ellipsis stands for multiparticle state contributions, which we shall not be interested in.

Let us now insert Eq. (B.9) in each of the two terms of Eq. (B.2). A look at Eqs. (B.4)-(B.7) then reveals that the 1-particle intermediate states contribute to the first term in Eq. (B.2) and the 1-antiparticle intermediate states to the second term (there is no contribution from $|0\rangle\langle 0|$ since the fermion field cannot have a vacuum expectation value). Thus,

$$iS'_{f\alpha\beta}(x) \Big|_{1\text{-particle}} = \theta(x_0) \sum_s \int \frac{d^3 P}{(2\pi)^3} \langle 0 | \psi_\alpha(x) | f(\vec{P}, s) \rangle \langle f(\vec{P}, s) | \bar{\psi}_\beta(0) | 0 \rangle, \quad (\text{B.10})$$

$$iS'_{f\alpha\beta}(x) \Big|_{1\text{-antiparticle}} = -\theta(-x_0) \sum_s \int \frac{d^3 P}{(2\pi)^3} \langle 0 | \bar{\psi}_\beta(0) | \bar{f}(\vec{P}, s) \rangle \langle \bar{f}(\vec{P}, s) | \psi_\alpha(x) | 0 \rangle. \quad (\text{B.11})$$

In Eq. (B.10), use Eqs. (B.4) and (B.5), and also

$$\theta(x_0) = i \int \frac{dp_0}{2\pi} \frac{e^{-ip_0 x_0}}{p_0 + i\epsilon}. \quad (\text{B.12})$$

Changing p_0 to $p_0 - E_P$ then leads to

$$S'_f(x) \Big|_{1\text{-particle}} = \int \frac{d^4 p}{(2\pi)^4} e^{-ip \cdot x} \frac{Z \sum_s u_s(\vec{P}) \bar{u}_s(\vec{P})}{p_0 - E_p + i\epsilon}. \quad (\text{B.13})$$

In Eq. (B.11), use Eqs. (B.6) and (B.7), and also put $\theta(-x_0)$ from Eq. (B.12). Changing p_0 to $-p_0 - E_P$ and $\vec{P}$ to $-\vec{P}$ then leads to

$$S'_f(x) \Big|_{1\text{-antiparticle}} = \int \frac{d^4 p}{(2\pi)^4} e^{-ip \cdot x} \frac{Z \sum_s v_s(-\vec{P}) \bar{v}_s(-\vec{P})}{p_0 + E_p - i\epsilon}. \quad (\text{B.14})$$

We are now going to check Eqs. (B.13) and (B.14) at the tree-level, for which $Z = 1$. We first note that Eqs. (B.4) and (B.8) imply that

$$u_s^\dagger(\vec{P})u_s(\vec{P}) = 1. \quad (\text{B.15})$$

(This can be seen in the following way. Let us introduce the factors N_1, N_2, N_3, N_4 through

$$u_s^\dagger(\vec{P})u_{s'}(\vec{P}) = N_1\delta_{ss'}, \quad (\text{B.16})$$

$$\psi(x) = \sum_s \int \frac{d^3P}{\sqrt{(2\pi)^3 N_2}} \left[c_s(\vec{P})u_s(\vec{P})e^{-ip \cdot x} + \dots \right], \quad (\text{B.17})$$

$$\left[c_s(\vec{P}), c_{s'}^\dagger(\vec{P}') \right]_+ = N_3\delta_{ss'}\delta^3(\vec{P} - \vec{P}'), \quad (\text{B.18})$$

$$|f(\vec{P}, s)\rangle = \sqrt{N_4}c_s^\dagger(\vec{P})|0\rangle, \quad (\text{B.19})$$

omitting the antiparticle parts for brevity. Then, $[\psi(\vec{x}, t), \psi(\vec{y}, t)]_+ = \delta^3(\vec{x} - \vec{y})$ gives $N_1 N_3 = N_2$, Eq. (B.4) (with $Z = 1$) gives $N_3^2 N_4 = (2\pi)^3 N_2$, and Eq. (B.8) gives $N_3 N_4 = (2\pi)^3$. It then follows that $N_1 = 1$.)

Similarly, $v_s^\dagger(p)v_s(\vec{P}) = 1$. For these normalizations, the spin-sum relations are given by Eq. (2.46). Since

$$S_f(p_0, \vec{P}) = \frac{p_0\gamma_0 - \vec{P} \cdot \vec{\gamma} + m_f}{p_0^2 - E_P^2 + i\epsilon} \approx \frac{E_P\gamma_0 - \vec{P} \cdot \vec{\gamma} + m_f}{2E_P(p_0 - E_P + i\epsilon)} \quad (\text{B.20})$$

near the pole $p_0 = E_P - i\epsilon$, Eq. (B.13) (with $Z = 1$) checks out. Similarly one can check Eq. (B.14).

In a medium, Eqs. (B.13) and (B.14) are generalized to Eqs. (4.38) and (4.52) respectively, while maintaining the relations given by Eq. (B.8) (see remark after Eq. (5.18)) and Eq. (B.15) (see Eq. (4.27)) which do not depend on the wavefunction renormalization factor.

APPENDIX C

Charge conjugation and the 1PI gravitational vertex

The aim of this Appendix is to derive the charge conjugation property of the 1PI gravitational vertex, given by Eq. (C.20), or Eq. (C.21), below.

We begin by considering the case of a C invariant Lagrangian and a C symmetric background. Since $\mu = 0$, Eqs. (2.2) – (2.4) reduce to

$$\langle \mathcal{O} \rangle = \frac{\text{Tr}(e^{-\beta H} \mathcal{O})}{\text{Tr} e^{-\beta H}}. \quad (\text{C.1})$$

Using $C^2 = 1$, this can be rewritten as

$$\langle \mathcal{O} \rangle = \frac{\text{Tr}(C e^{-\beta H} C C \mathcal{O} C)}{\text{Tr}(C e^{-\beta H} C)}. \quad (\text{C.2})$$

Use of $CHC = H$ (see Eq. (4.81)) and $C^2 = 1$ then gives

$$\langle \mathcal{O} \rangle = \langle C \mathcal{O} C \rangle. \quad (\text{C.3})$$

It will be instructive to first apply this equation to the definition of the full fermion propagator $S'_f(p)$, given by

$$i(2\pi)^4 \delta^{(4)}(p - p') S'_{f_{nm}}(p) = \int d^4 y d^4 z e^{ip \cdot y} e^{-ip' \cdot z} \langle T[\psi_n(y) \bar{\psi}_m(z)] \rangle, \quad (\text{C.4})$$

where the fermion fields are the Heisenberg picture fields. (This is equivalent to the definition

$$\langle T[\psi_n(y) \bar{\psi}_m(z)] \rangle = i \int \frac{d^4 k}{(2\pi)^4} e^{-ik \cdot (y-z)} S'_{f_{nm}}(k) \quad (\text{C.5})$$

used elsewhere in this work.) Using Eq. (C.3) and $C^2 = 1$, we can write

$$\langle T[\psi_n(y) \bar{\psi}_m(z)] \rangle = \langle T[C \psi_n(y) C C \bar{\psi}_m(z) C] \rangle. \quad (\text{C.6})$$

Now, $C\psi C$ is given by Eq (4.60). Also, $C\bar{\psi}C = (C\psi C)^\dagger \gamma_0$, since (as explained below Eq. (4.83)) C is hermitian. So, using Eqs. (4.60), (4.61) and (4.62), we obtain

$$C\bar{\psi}C = -\psi^T C^{-1}. \quad (C.7)$$

One can now use Eqs. (4.60) and (C.7) to determine the R.H.S. of Eq. (C.6). Consequently, the R.H.S. of Eq. (C.4) equals

$$C_{nb}(C^{-1})_{am} \int d^4z d^4y e^{-ip'\cdot z} e^{ip\cdot y} \langle T[\psi_a(z) \bar{\psi}_b(y)] \rangle. \quad (C.8)$$

The double integral is just $i(2\pi)^4 \delta^{(4)}(p - p') S'_{fab}(-p)$, by the definition (C.4). So the expression given in Eq. (C.8) equals

$$i(2\pi)^4 \delta^{(4)}(p - p') [CS_f'^T(-p)C^{-1}]_{nm}. \quad (C.9)$$

Since this equals the L.H.S. of Eq. (C.4), we arrive at

$$S'_f(p) = CS_f'^T(-p)C^{-1}. \quad (C.10)$$

As a check, we note that the use of Eq. (4.1) reduces Eq. (C.10) to Eq. (4.74), which was deduced by demanding the action to be C invariant.

Now we shall apply Eq. (C.3) to the definition of the 1PI gravitational vertex function $\Gamma^{\lambda\rho}(p, p')$, given by

$$\begin{aligned} & (2\pi)^4 \delta^{(4)}(p - p' - q) S'_{fnn'}(p) \Gamma_{n'm'}^{\lambda\rho}(p, p') S'_{fm'm}(p') \\ &= \int d^4x d^4y d^4z e^{-iq\cdot x} e^{ip\cdot y} e^{-ip'\cdot z} \langle T[\hat{T}^{\lambda\rho}(x) \psi_n(y) \bar{\psi}_m(z)] \rangle. \end{aligned} \quad (C.11)$$

Using Eq. (C.3), $C^2 = 1$, and $C\hat{T}^{\lambda\rho}C = \hat{T}^{\lambda\rho}$, we can write

$$\langle T[\hat{T}^{\lambda\rho}(x) \psi_n(y) \bar{\psi}_m(z)] \rangle = \langle T[\hat{T}^{\lambda\rho}(x) C\psi_n(y) C C\bar{\psi}_m(z) C] \rangle. \quad (C.12)$$

One can now use Eqs. (4.60) and (C.7) to determine the R.H.S. of Eq. (C.12). Consequently, the R.H.S. of Eq. (C.11) equals

$$C_{nb}(C^{-1})_{am} \int d^4x d^4y d^4z e^{-iq\cdot x} e^{-ip'\cdot z} e^{ip\cdot y} \langle T[\hat{T}^{\lambda\rho}(x) \psi_a(z) \bar{\psi}_b(y)] \rangle. \quad (C.13)$$

The triple integral is just $(2\pi)^4 \delta^{(4)}(p - p' - q) S'_{fan'}(-p') \Gamma_{n'm'}^{\lambda\rho}(-p', -p) S'_{fm'b}(-p)$, by the definition (C.11). So the expression given in Eq. (C.13) equals

$$(2\pi)^4 \delta^{(4)}(p - p' - q) [CS_f'^T(-p) \Gamma^{\lambda\rho T}(-p', -p) S_f'^T(-p') C^{-1}]_{nm}. \quad (C.14)$$

Since this equals the L.H.S. of Eq. (C.11), we arrive at

$$S'_f(p)\Gamma_{\lambda\rho}(p,p')S'_f(p') = \mathbf{C}S'^{\text{T}}_f(-p)\Gamma_{\lambda\rho}^{\text{T}}(-p',-p)S'^{\text{T}}_f(-p')\mathbf{C}^{-1}. \quad (\text{C.15})$$

Using Eq. (C.10), this leads to

$$\Gamma_{\lambda\rho}(p,p') = \mathbf{C}\Gamma_{\lambda\rho}^{\text{T}}(-p',-p)\mathbf{C}^{-1}. \quad (\text{C.16})$$

Now, consider the case of C symmetric Lagrangian and C asymmetric background. Then

$$\langle \mathcal{O} \rangle = \frac{\text{Tr}(e^{-\beta(H-\mu Q)}\mathcal{O})}{\text{Tr} e^{-\beta(H-\mu Q)}}, \quad (\text{C.17})$$

instead of Eq. (C.1). Since $CQC = -Q$ (see Eq. (4.81)), we now have

$$\langle \mathcal{O} \rangle = [\langle COC \rangle]_{\mu \rightarrow -\mu} \quad (\text{C.18})$$

in the place of Eq. (C.3). It is easy to see that this results in

$$S'_f(p) = [\mathbf{C}S'^{\text{T}}_f(-p)\mathbf{C}^{-1}]_{\mu \rightarrow -\mu}, \quad (\text{C.19})$$

$$\Gamma_{\lambda\rho}(p,p') = [\mathbf{C}\Gamma_{\lambda\rho}^{\text{T}}(-p',-p)\mathbf{C}^{-1}]_{\mu \rightarrow -\mu}. \quad (\text{C.20})$$

Since μ enters through the thermal propagator of the fermion given by Eqs. (2.50) and (2.51), and $\mu \rightarrow -\mu$ in the propagator is equivalent to $v \rightarrow -v$, one can also write Eq. (C.20) in the form

$$\Gamma_{\lambda\rho}(p,p',v) = \mathbf{C}\Gamma_{\lambda\rho}^{\text{T}}(-p',-p,-v)\mathbf{C}^{-1}, \quad (\text{C.21})$$

where we have explicitly indicated the dependence of the vertex function on the vector v^μ .

APPENDIX D

Transversality of the gravitational vertex

It is useful to verify that the complete vertex function to $O(e^2)$, obtained in Sec. 5.4, satisfies the transversality conditions

$$q^\lambda \bar{U}(\vec{P}') \Gamma_{\lambda\rho}(p, p') U(\vec{P}) = 0 = q^\rho \bar{U}(\vec{P}') \Gamma_{\lambda\rho}(p, p') U(\vec{P}) \quad (\text{D.1})$$

to this order. The wavefunction renormalization factors associated with the external lines, being scalars, do not play any role in these relations (in other words, they just cancel out). Also, since the vertex is symmetric in the Lorentz indices λ, ρ , either of these relations guarantees the other. In order to simplify the notation, in this Appendix we omit the subscript s in the spinors.

In order to verify this relation, the important point is that we must include all the terms upto $O(e^2)$. Since the one-loop terms in the induced vertex are already $O(e^2)$, for them we can adopt the tree-level definition of the spinors, i.e.

$$\not{p} u(\vec{P}) = m u(\vec{P}), \quad \bar{u}(\vec{P}') \not{p}' = m \bar{u}(\vec{P}'), \quad (\text{D.2})$$

as well as the tree-level on-shell conditions

$$p^2 = p'^2 = m^2. \quad (\text{D.3})$$

In this appendix, as well as in Appendix E, we use the tree level mass m without any subscript, implying m_ℓ , m_e or m_f which should be understood from the context. Also note that the photon distribution function as well as the associated δ -function are even in k , and therefore those terms which are odd in k in the rest of the integrand do not contribute.

We first show that the vertex contribution from Fig. 5.2A is transverse by itself. From Eq. (5.54),

$$q^\lambda A_{\lambda\rho\alpha}(k, k-q) = -(q^2 - 2k \cdot q)(4k_\rho k_\alpha + k \cdot q \eta_{\rho\alpha}), \quad (\text{D.4})$$

where we omit the terms that are proportional to q_α because, in Eq. (5.49), they will yield $\not{q}$ which vanishes between the spinors. Changing the sign of q in the last equation yields

$$q^\lambda A_{\lambda\rho\alpha}(k+q, k) = (q^2 + 2k \cdot q)(4k_\rho k_\alpha - k \cdot q \eta_{\rho\alpha}), \quad (\text{D.5})$$

and as a result $q^\lambda \bar{u}(\vec{P}') \Gamma_{\lambda\rho}^{(X)}(p, p') u(\vec{P})$ turns out to be proportional to $\sum_f Q_f(n_f - n_{\bar{f}})$, which is zero to this order.

As for the other diagrams, straightforward algebra gives the following results:

$$\begin{aligned} q^\lambda \bar{u}(\vec{P}') \Gamma_{\lambda\rho}^{(A1)}(p, p') u(\vec{P}) &= -\frac{e^2}{4} \int \frac{d^4 k}{(2\pi)^3} \delta(k^2) \eta_\gamma(k) \\ &\times \bar{u}(\vec{P}') \left[\frac{4mk_\rho - (p+5p')_\rho \not{k} + 2k \cdot p \gamma_\rho}{k \cdot p'} - (p \leftrightarrow p') \right] u(\vec{P}) \\ q^\lambda \bar{u}(\vec{P}') \Gamma_{\lambda\rho}^{(A2)}(p, p') u(\vec{P}) &= \frac{e^2}{4} \int \frac{d^4 k}{(2\pi)^3} \delta(k^2 - m^2) \eta_f(k) \\ &\times \bar{u}(\vec{P}') \left[\frac{4(\not{k} - 2m)k_\rho + \not{k}(p+p')_\rho - 2k \cdot p' \gamma_\rho}{m^2 - k \cdot p} \right. \\ &\quad \left. - (p \leftrightarrow p') \right] u(\vec{P}), \\ q^\lambda \bar{u}(\vec{P}') \Gamma_{\lambda\rho}^{(B1)}(p, p') u(\vec{P}) &= e^2 \int \frac{d^4 k}{(2\pi)^3} \delta(k^2) \eta_\gamma(k) \\ &\times \bar{u}(\vec{P}') \left[\frac{mk_\rho - \not{k}p'_\rho}{k \cdot p'} - \frac{mk_\rho - \not{k}p_\rho}{k \cdot p} \right] u(\vec{P}) \\ q^\lambda \bar{u}(\vec{P}') \Gamma_{\lambda\rho}^{(B2)}(p, p') u(\vec{P}) &= -e^2 \int \frac{d^4 k}{(2\pi)^3} \delta(k^2 - m^2) \eta_f(k) \\ &\times \bar{u}(\vec{P}') \left[\frac{(\not{k} - 2m)k_\rho + mp_\rho}{m^2 - k \cdot p} - \frac{(\not{k} - 2m)k_\rho + mp'_\rho}{m^2 - k \cdot p'} \right] u(\vec{P}), \\ q^\lambda \bar{u}(\vec{P}') \Gamma_{\lambda\rho}^{(C1+D1)}(p, p') u(\vec{P}) &= \frac{e^2}{2} \int \frac{d^4 k}{(2\pi)^3} \delta(k^2) \eta_\gamma(k) \left(\frac{1}{k \cdot p'} + \frac{1}{k \cdot p} \right) \\ &\times \bar{u}(\vec{P}') [\not{k}q_\rho + k \cdot q \gamma_\rho] u(\vec{P}), \\ q^\lambda \bar{u}(\vec{P}') \Gamma_{\lambda\rho}^{(C2+D2)}(p, p') u(\vec{P}) &= -\frac{e^2}{2} \int \frac{d^4 k}{(2\pi)^3} \delta(k^2 - m^2) \eta_f(k) \\ &\times \left(\frac{1}{m^2 - k \cdot p'} + \frac{1}{m^2 - k \cdot p} \right) \\ &\times \bar{u}(\vec{P}') [(\not{k} - 3m)q_\rho + k \cdot q \gamma_\rho] u(\vec{P}). \end{aligned} \quad (\text{D.6})$$

Therefore, adding all the one-loop contributions to the vertex, we obtain

$$q^\lambda \bar{u}(\vec{P}') \Gamma_{\lambda\rho}^{(1)}(p, p') u(\vec{P}) = \frac{e^2}{4} \int \frac{d^4 k}{(2\pi)^3} \delta(k^2) \eta_\gamma(k) \times \bar{u}(\vec{P}') \left[\frac{\not{k}}{k \cdot p'} (3p_\rho - p'_\rho) - (p \leftrightarrow p') \right] u(\vec{P}), \quad (\text{D.7})$$

$$q^\lambda \bar{u}(\vec{P}') \Gamma_{\lambda\rho}^{(2)}(p, p') u(\vec{P}) = \frac{e^2}{4} \int \frac{d^4 k}{(2\pi)^3} \delta(k^2 - m^2) \eta_f(k) \times \bar{u}(\vec{P}') \left[\frac{(\not{k} - 2m)(3p'_\rho - p_\rho) - 2k \cdot p \gamma_\rho}{m^2 - k \cdot p} - (p \leftrightarrow p') \right] u(\vec{P}). \quad (\text{D.8})$$

We need to add to these the tree-level contribution to the gravitational vertex that appears in Eq. (3.95). In this case, we must include the $O(e^2)$ corrections to the equation for the spinors, which arise from the self-energy diagrams of Sec. 4.5. Thus, for this part, using Eq. (4.26) and its hermitian conjugate

$$\bar{U}(\vec{P}') (\not{p}' - m - \Sigma(p')) = 0, \quad (\text{D.9})$$

we obtain

$$\bar{U}(\vec{P}') \not{p}' U(\vec{P}) = \bar{U}(\vec{P}') (\Sigma'(p) - \Sigma'(p')) U(\vec{P}), \quad (\text{D.10})$$

which in turn yields

$$q^\lambda \bar{U}(\vec{P}') V_{\lambda\rho}(p, p') U(\vec{P}) = \frac{1}{4} \bar{U}(\vec{P}') \left[(3p' - p)_\rho \Sigma'(p) + p^2 \gamma_\rho - (p \leftrightarrow p') \right] U(\vec{P}). \quad (\text{D.11})$$

This can be cast in a different form by multiplying Eq. (4.26) from the left by $\bar{U}(\vec{P}') \gamma_\rho (\not{p} + m)$ and Eq. (D.9) from the right by $(\not{p}' + m) \gamma_\rho U(\vec{P})$ and taking the difference of the resulting equations. This gives

$$(p^2 - p'^2) \bar{U}(\vec{P}') \gamma_\rho U(\vec{P}) = \bar{U}(\vec{P}') \left[\gamma_\rho (\not{p} + m) \Sigma'(p) - \Sigma'(p') (\not{p}' + m) \gamma_\rho \right] U(\vec{P}), \quad (\text{D.12})$$

and substituting this result into Eq. (D.11), we obtain

$$q^\lambda \bar{U}(\vec{P}') V_{\lambda\rho}(p, p') U(\vec{P}) = \frac{1}{4} \bar{U}(\vec{P}') \left[(3p' - p)_\rho \Sigma'(p) - (3p - p')_\rho \Sigma'(p') + \gamma_\rho (\not{p} + m) \Sigma'(p) - \Sigma'(p') (\not{p}' + m) \gamma_\rho \right] U(\vec{P}). \quad (\text{D.13})$$

Since Σ' is explicitly of $O(e^2)$ while we are interested in results to $O(e^2)$ only, we can use the tree-level spinors on the right-hand side. Using Eq. (D.3) in Eq. (4.95), we can write

the self-energy contribution involving the photon distribution function as

$$\Sigma'_1(p) = e^2 \int \frac{d^4 k}{(2\pi)^3} \delta(k^2) \eta_\gamma(k) \frac{\not{k}}{k \cdot p}, \quad (\text{D.14})$$

disregarding terms odd in k . Similarly, from Eq. (4.96), the part containing the Fermi distribution function can be written as

$$\Sigma'_2(p) = -e^2 \int \frac{d^4 k}{(2\pi)^3} \delta(k^2 - m^2) \eta_f(k) \frac{\not{k} - 2m}{m^2 - k \cdot p}. \quad (\text{D.15})$$

Substituting these forms into Eq. (D.13) and using the identities

$$\begin{aligned} \bar{u}(\vec{P}') \gamma_\rho (\not{p} + m) \not{k} u(\vec{P}) &= 2k \cdot p \bar{u}(\vec{P}') \gamma_\rho u(\vec{P}) \\ \bar{u}(\vec{P}') \not{k} (\not{p}' + m) \gamma_\rho u(\vec{P}) &= 2k \cdot p' \bar{u}(\vec{P}') \gamma_\rho u(\vec{P}), \end{aligned} \quad (\text{D.16})$$

we see that Eq. (D.13) cancels the contribution from the loop diagrams given in Eqs. (D.7) and (D.8) to this order. This proves the transversality of the effective vertex.

APPENDIX E

Apparently ill-defined contributions to the gravitational vertex

E.1 The B1 contribution

We start from the formula given in Eq. (5.43), from which it follows that

$$\begin{aligned} \Gamma_{\lambda\rho}^{\prime(B1)}(p, p) = & -\frac{e^2}{2} \lim_{\vec{Q} \rightarrow 0} \int \frac{d^4 k}{(2\pi)^3} \delta(k^2) \eta_\gamma(k) \left[\frac{\gamma^\nu (\not{p} - \not{k} + m) \gamma^\mu C_{\mu\nu\lambda\rho}(k, k - q)}{k \cdot p (2\vec{K} \cdot \vec{Q} - Q^2)} \right. \\ & \left. - \frac{\gamma^\nu (\not{p} - \not{k} - \not{q} + m) \gamma^\mu C_{\mu\nu\lambda\rho}(k + q, k)}{(k \cdot p + \vec{K} \cdot \vec{Q})(2\vec{K} \cdot \vec{Q} + Q^2)} \right], \end{aligned} \quad (\text{E.1})$$

where we have put

$$q^\mu = (0, \vec{Q}) \quad k^\mu = (k_0, \vec{K}). \quad (\text{E.2})$$

In order to take the limit $\vec{Q} \rightarrow 0$, our strategy is to expand the coefficients of the factors $1/(2\vec{K} \cdot \vec{Q} \pm Q^2)$ in powers of $\vec{Q}$. Of the resulting terms in the coefficients, those which are quadratic in $\vec{Q}$ do not contribute in the $\vec{Q} \rightarrow 0$ limit and therefore we need to keep only the terms that are at most linear in $\vec{Q}$.

Using the property $C_{\mu\nu\lambda\rho}(k + q, k) = C_{\nu\mu\lambda\rho}(k, k + q)$, we can write

$$\begin{aligned} C_{\mu\nu\lambda\rho}(k, k - q) &= C_{\mu\nu\lambda\rho}(k, k) - C'_{\mu\nu\lambda\rho}(k, q) \\ C_{\mu\nu\lambda\rho}(k + q, k) &= C_{\mu\nu\lambda\rho}(k, k) + C'_{\nu\mu\lambda\rho}(k, q), \end{aligned} \quad (\text{E.3})$$

where

$$\begin{aligned} C'_{\mu\nu\lambda\rho}(k, q) = & \eta_{\lambda\rho}(\eta_{\mu\nu} k \cdot q - q_\mu k_\nu) - \eta_{\mu\nu}(k_\lambda q_\rho + q_\lambda k_\rho) + k_\nu(\eta_{\lambda\mu} q_\rho + \eta_{\rho\mu} q_\lambda) \\ & + q_\mu(\eta_{\lambda\nu} k_\rho + \eta_{\rho\nu} k_\lambda) - k \cdot q(\eta_{\lambda\mu} \eta_{\rho\nu} + \eta_{\lambda\nu} \eta_{\rho\mu}). \end{aligned} \quad (\text{E.4})$$

To first order in Q , we can also put

$$\frac{1}{k \cdot p + \vec{K} \cdot \vec{Q}} = \frac{1}{k \cdot p} - \frac{\vec{K} \cdot \vec{Q}}{(k \cdot p)^2}. \quad (\text{E.5})$$

This enables us to decompose $\Gamma_{\lambda\rho}^{(B1)}(p, p)$ in the following four terms:

$$\begin{aligned} \Gamma_{\lambda\rho}^{(B1a)}(p) &= -\frac{e^2}{2} \lim_{\vec{Q} \rightarrow 0} \int \frac{d^4 k}{(2\pi)^3} \delta(k^2) \eta_\gamma(k) \frac{C_{\mu\nu\lambda\rho}(k, k) \gamma^\nu (\not{p} - \not{k} + m) \gamma^\mu}{k \cdot p} \\ &\quad \times \left[\frac{1}{2\vec{K} \cdot \vec{Q} - Q^2} - \frac{1}{2\vec{K} \cdot \vec{Q} + Q^2} \right] \\ \Gamma_{\lambda\rho}^{(B1b)}(p) &= -\frac{e^2}{2} \lim_{\vec{Q} \rightarrow 0} \int \frac{d^4 k}{(2\pi)^3} \delta(k^2) \eta_\gamma(k) \frac{C_{\mu\nu\lambda\rho}(k, k) \gamma^\nu \not{Q} \gamma^\mu}{k \cdot p (2\vec{K} \cdot \vec{Q} + Q^2)} \\ \Gamma_{\lambda\rho}^{(B1c)}(p) &= -\frac{e^2}{2} \lim_{\vec{Q} \rightarrow 0} \int \frac{d^4 k}{(2\pi)^3} \delta(k^2) \eta_\gamma(k) \frac{C_{\mu\nu\lambda\rho}(k, k) \gamma^\nu (\not{p} - \not{k} + m) \gamma^\mu}{(k \cdot p)^2} \left[\frac{\vec{K} \cdot \vec{Q}}{2\vec{K} \cdot \vec{Q} + Q^2} \right] \\ \Gamma_{\lambda\rho}^{(B1d)}(p) &= -\frac{e^2}{2} \lim_{\vec{Q} \rightarrow 0} \int \frac{d^4 k}{(2\pi)^3} \delta(k^2) \eta_\gamma(k) \frac{\gamma^\nu (\not{p} - \not{k} + m) \gamma^\mu}{k \cdot p} \\ &\quad \times \left[-\frac{C'_{\mu\nu\lambda\rho}(k, q)}{2\vec{K} \cdot \vec{Q} - Q^2} - \frac{C'_{\nu\mu\lambda\rho}(k, q)}{2\vec{K} \cdot \vec{Q} + Q^2} \right]. \end{aligned} \quad (\text{E.6})$$

We carry out these integrals one by one.

Eliminating the manifestly k -odd terms from the integrand and performing the k_0 -integration, we obtain

$$\Gamma_{\lambda\rho}^{(B1a)}(p) = e^2 \lim_{\vec{Q} \rightarrow 0} \int \frac{d^3 K}{(2\pi)^3 2K} f_\gamma(K) \frac{4k_\lambda k_\rho \not{k}}{k \cdot p} \left[\frac{1}{2\vec{K} \cdot \vec{Q} - Q^2} - \frac{1}{2\vec{K} \cdot \vec{Q} + Q^2} \right], \quad (\text{E.7})$$

using $k^2 = 0$. The expression within the square brackets is finite for $Q \rightarrow 0$. Therefore, in the spinors we can set $p = p'$, and using Eq. (4.58) we then obtain

$$m'_{(B1a)} = \frac{4e^2}{m} \lim_{\vec{Q} \rightarrow 0} \int \frac{d^3 K}{(2\pi)^3} f_\gamma(K) K \left[\frac{1}{2\vec{K} \cdot \vec{Q} - Q^2} - \frac{1}{2\vec{K} \cdot \vec{Q} + Q^2} \right]. \quad (\text{E.8})$$

We can perform the integration over the angular variables in $\vec{K}$, the integral being understood, as usual, in terms of the principal value part. That gives

$$\int d\Omega \frac{1}{2\vec{K} \cdot \vec{Q} - Q^2} = - \int d\Omega \frac{1}{2\vec{K} \cdot \vec{Q} + Q^2} = -\frac{\pi}{K^2} + \mathcal{O}(Q^2), \quad (\text{E.9})$$

so that

$$m'_{(B1a)} = -\frac{e^2}{\pi^2 m} \int dK f_\gamma(K) K = -\frac{e^2 T^2}{6m}. \quad (\text{E.10})$$

As for the next contribution, it is straightforward to verify that

$$(2v^\lambda v^\rho - \eta^{\lambda\rho})\gamma^\nu \not{q} \gamma^\mu C_{\mu\nu\lambda\rho}(k, k) = 4\vec{K} \cdot \vec{Q}(\not{k} - 2k \cdot v \not{p}), \quad (\text{E.11})$$

using $q \cdot v = q_0 = 0$. So

$$(2v^\lambda v^\rho - \eta^{\lambda\rho})\Gamma_{\lambda\rho}^{\prime(B1b)}(p) = e^2 \int \frac{d^4 k}{(2\pi)^3} \delta(k^2) \eta_\gamma(k) \frac{(-\not{k} + 2k \cdot v \not{p})}{k \cdot p}. \quad (\text{E.12})$$

Now using Eqs. (4.58) and (4.59), carrying out the integral over k_0 , and finally putting $P = 0$, we get

$$m'_{(B1b)} = \frac{2e^2}{m} \int \frac{d^3 K}{(2\pi)^3 2K} f_\gamma(K) = \frac{e^2 T^2}{12m}. \quad (\text{E.13})$$

Similarly,

$$\Gamma_{\lambda\rho}^{\prime(B1c)}(p) = -\frac{e^2}{2} \int \frac{d^3 K}{(2\pi)^3 2K} f_\gamma(K) \gamma^\nu (\not{p} + m) \gamma^\mu C_{\mu\nu\lambda\rho}(k, k) \frac{1}{(k \cdot p)^2}, \quad (\text{E.14})$$

and

$$\begin{aligned} (2v^\lambda v^\rho - \eta^{\lambda\rho})\Gamma_{\lambda\rho}^{\prime(B1c)}(p) &= -\frac{2e^2}{m} \int \frac{d^3 K}{(2\pi)^3 2K} f_\gamma(K) \frac{1}{(k \cdot p)^2} \\ &\times \left[-(k \cdot p)^2 - 2m^2(k \cdot v)^2 + 4(k \cdot p)(k \cdot v)(p \cdot v) \right] \end{aligned} \quad (\text{E.15})$$

so that

$$m'_{(B1c)} = -\frac{2e^2}{m} \int \frac{d^3 K}{(2\pi)^3 2K} f_\gamma(K) = -\frac{e^2 T^2}{12m}. \quad (\text{E.16})$$

For $\Gamma_{\lambda\rho}^{\prime(B1d)}$ we first perform the integral over k_0 . Remembering that in the remaining integral we can change $\vec{K}$ to $-\vec{K}$ and using the fact that $C'_{\mu\nu\lambda\rho}(-k, q) = -C'_{\mu\nu\lambda\rho}(k, q)$, we obtain

$$\begin{aligned} \Gamma_{\lambda\rho}^{\prime(B1d)}(p) &= \frac{e^2}{2} \lim_{\vec{Q} \rightarrow 0} \int \frac{d^3 K}{(2\pi)^3 2K} f_\gamma(K) \left(\frac{1}{k \cdot p} \right) \\ &\times \left\{ \gamma^\nu (\not{p} + m) \gamma^\mu \left[\frac{1}{2\vec{K} \cdot \vec{Q} - Q^2} - \frac{1}{2\vec{K} \cdot \vec{Q} + Q^2} \right] [C'_{\mu\nu\lambda\rho}(k, q) - C'_{\nu\mu\lambda\rho}(k, q)] \right. \\ &\left. + \gamma^\nu (-\not{k}) \gamma^\mu \left[\frac{1}{2\vec{K} \cdot \vec{Q} - Q^2} + \frac{1}{2\vec{K} \cdot \vec{Q} + Q^2} \right] [C'_{\mu\nu\lambda\rho}(k, q) + C'_{\nu\mu\lambda\rho}(k, q)] \right\} \\ &= -\frac{e^2}{2} \lim_{\vec{Q} \rightarrow 0} \int \frac{d^3 K}{(2\pi)^3 2K} f_\gamma(K) \frac{1}{k \cdot p} \\ &\times \left\{ \frac{\gamma^\nu \not{k} \gamma^\mu}{\vec{K} \cdot \vec{Q}} [C'_{\mu\nu\lambda\rho}(k, q) + C'_{\nu\mu\lambda\rho}(k, q)] + O(Q) \right\}. \end{aligned} \quad (\text{E.17})$$

We now use

$$(2v^\lambda v^\rho - \eta^{\lambda\rho})\gamma^\nu \not{k} \gamma^\mu [C'_{\mu\nu\lambda\rho}(k, q) + C'_{\nu\mu\lambda\rho}(k, q)] = 8(k \cdot v)(\vec{K} \cdot \vec{Q})\not{p}. \quad (\text{E.18})$$

Then, using Eq. (4.58) and putting $\vec{P} = 0$, we get

$$m'_{(B2d)} = -\frac{4e^2}{m} \int \frac{d^3 K}{(2\pi)^3 2K} f_\gamma(K) = -\frac{e^2 T^2}{6m}. \quad (\text{E.19})$$

Adding the results given in Eqs. (E.10), (E.13), (E.16) and (E.19), we get the total contribution from the B1 term presented in Eq. (5.63).

E.2 The A2 contribution

For this contribution, we start from Eq. (5.40). Using Eq. (E.2), we can write it as

$$\begin{aligned} \Gamma_{\lambda\rho}'^{(A2)}(p, p) &= \frac{e^2}{2} \lim_{\vec{Q} \rightarrow 0} \int \frac{d^4 k}{(2\pi)^3} \delta(k^2 - m^2) \eta_f(k) \\ &\times \left[\frac{\Lambda_{\lambda\rho}(k, k - q)}{(2\vec{K} \cdot \vec{Q} - Q^2)(m^2 - k \cdot p)} - \frac{\Lambda_{\lambda\rho}(k + q, k)}{(2\vec{K} \cdot \vec{Q} + Q^2)(m^2 - k \cdot p')} \right] \end{aligned} \quad (\text{E.20})$$

Following the strategy stated below Eq. (E.2), let us now write

$$\begin{aligned} \Lambda_{\lambda\rho}(k, k - q) &= \Lambda_{\lambda\rho}(k, k) + \Lambda'_{\lambda\rho}(k, q), \\ \Lambda_{\lambda\rho}(k + q, k) &= \Lambda_{\lambda\rho}(k, k) + \Lambda''_{\lambda\rho}(k, q), \end{aligned} \quad (\text{E.21})$$

and, in the denominator, expand $m^2 - k \cdot p'$ in powers of $\vec{Q}$:

$$\frac{1}{m^2 - k \cdot p'} = \frac{1}{m^2 - k \cdot p} + \frac{\vec{K} \cdot \vec{Q}}{(m^2 - k \cdot p)^2} + O(Q^2). \quad (\text{E.22})$$

Then we can decompose $\Gamma_{\lambda\rho}'^{(A2)}(p, p)$ into the following terms, omitting higher powers of Q which anyway will not contribute:

$$\begin{aligned} \Gamma_{\lambda\rho}'^{(A2a)}(p) &= \frac{e^2}{2} \lim_{\vec{Q} \rightarrow 0} \int \frac{d^4 k}{(2\pi)^3} \delta(k^2 - m^2) \eta_f(k) \\ &\times \frac{\Lambda_{\lambda\rho}(k, k)}{m^2 - k \cdot p} \left(\frac{1}{2\vec{K} \cdot \vec{Q} - Q^2} - \frac{1}{2\vec{K} \cdot \vec{Q} + Q^2} \right), \\ \Gamma_{\lambda\rho}'^{(A2b)}(p) &= -\frac{e^2}{4} \int \frac{d^4 k}{(2\pi)^3} \delta(k^2 - m^2) \eta_f(k) \frac{\Lambda_{\lambda\rho}(k, k)}{(m^2 - k \cdot p)^2}, \\ \Gamma_{\lambda\rho}'^{(A2c)}(p) &= \frac{e^2}{2} \lim_{\vec{Q} \rightarrow 0} \int \frac{d^4 k}{(2\pi)^3} \delta(k^2 - m^2) \eta_f(k) \\ &\times \frac{1}{m^2 - k \cdot p} \left(\frac{\Lambda'_{\lambda\rho}(k, q)}{2\vec{K} \cdot \vec{Q} - Q^2} - \frac{\Lambda''_{\lambda\rho}(k, q)}{2\vec{K} \cdot \vec{Q} + Q^2} \right), \end{aligned} \quad (\text{E.23})$$

We discuss these contributions one by one.

The A2a contribution

Using Eq. (5.56) and the δ -function appearing in the integrand, we can write

$$\begin{aligned}\Gamma_{\lambda\rho}^{(A2a)}(p) &= -2e^2 \lim_{\vec{Q} \rightarrow 0} \int \frac{d^4k}{(2\pi)^3} \delta(k^2 - m^2) \eta_f(k) \\ &\quad \times \frac{k_\lambda k_\rho (\not{k} - 2m)}{m^2 - k \cdot p} \left(\frac{1}{2\vec{K} \cdot \vec{Q} - Q^2} - \frac{1}{2\vec{K} \cdot \vec{Q} + Q^2} \right).\end{aligned}\quad (\text{E.24})$$

As argued before Eq. (E.8), we can put $\vec{Q} = 0$ in the spinors, and use Eq. (4.58). Performing the k_0 -integration, we obtain

$$(2v^\lambda v^\rho - \eta^{\lambda\rho}) \Gamma_{\lambda\rho}^{(A2a)}(p) = \frac{2e^2}{m} \lim_{\vec{Q} \rightarrow 0} \int \frac{d^3K}{(2\pi)^3} F(\vec{K}) \left(\frac{1}{2\vec{K} \cdot \vec{Q} - Q^2} - \frac{1}{2\vec{K} \cdot \vec{Q} + Q^2} \right), \quad (\text{E.25})$$

where the expression on the left is understood to equal the one on the right only between the spinors, and

$$F(\vec{K}) = \frac{2E_K^2 - m^2}{2E_K} \left[\left(1 + \frac{m^2}{m^2 - k \cdot p} \right) f_e + \left(1 + \frac{m^2}{m^2 + k \cdot p} \right) f_{\bar{e}} \right], \quad (\text{E.26})$$

with $k_0 = E_K$. Since the integrand contains $\vec{K} \cdot \vec{P}$, and we must set $\vec{P} = 0$ only after taking the limit $Q \rightarrow 0$, the angular integrations cannot be performed using Eq. (E.9). So we shift the integration variable to $\vec{K} \pm \frac{1}{2}\vec{Q}$ in the terms having $2\vec{K} \cdot \vec{Q} \mp Q^2$ in the denominator. This gives

$$(2v^\lambda v^\rho - \eta^{\lambda\rho}) \Gamma_{\lambda\rho}^{(A2a)}(p) = \frac{2e^2}{m} \lim_{\vec{Q} \rightarrow 0} \int \frac{d^3K}{(2\pi)^3} \frac{\vec{Q} \cdot \vec{\nabla}_K F}{2\vec{K} \cdot \vec{Q}}. \quad (\text{E.27})$$

Clearly the magnitude of $\vec{Q}$ now cancels out. The derivative with respect to $\vec{K}$ can be taken easily, using

$$\begin{aligned}\vec{\nabla}_K E_K &= \frac{\vec{K}}{E_K}, \\ \vec{\nabla}_K \left(\frac{1}{m^2 \pm k \cdot p} \right) &= \frac{\mp 1}{(m^2 \pm k \cdot p)^2} \left(E_P \frac{\vec{K}}{E_K} - \vec{P} \right).\end{aligned}\quad (\text{E.28})$$

The term proportional to $\vec{P}$ from the last derivative does not contribute because it multiplies a factor whose integrand is odd in $\vec{K}$ at $\vec{P} = 0$. Putting $\vec{P} = 0$ in the other terms,

we obtain the contribution to the gravitational mass:

$$m'_{(A2a)} = \frac{2e^2}{m} \int \frac{d^3K}{(2\pi)^3 2E_K} \times \left\{ \frac{2E_K^2 - m^2}{2E_K} \left(\frac{E_K - 2m}{E_K - m} \frac{\partial f_e}{\partial E_K} + \frac{E_K + 2m}{E_K + m} \frac{\partial f_{\bar{e}}}{\partial E_K} \right) \right. \\ \left. + \frac{2E_K^2 - m^2}{2E_K} \left(\frac{m}{(E_K - m)^2} f_e - \frac{m}{(E_K + m)^2} f_{\bar{e}} \right) \right. \\ \left. + \frac{2E_K^2 + m^2}{2E_K^2} \left(\frac{E_K - 2m}{E_K - m} f_e + \frac{E_K + 2m}{E_K + m} f_{\bar{e}} \right) \right\}. \quad (\text{E.29})$$

The A2b contribution

The integral in the (A2b) term is independent of Q . So, in a straight forward way, we obtain

$$m'_{(A2b)} = \frac{e^2}{m^2} \int \frac{d^3K}{(2\pi)^3 2E_K} (2E_K^2 - m^2) \left[\frac{E_K - 2m}{(E_K - m)^2} f_e(E_K) - \frac{E_K + 2m}{(E_K + m)^2} f_{\bar{e}}(E_K) \right]. \quad (\text{E.30})$$

The A2c contribution

For the (A2c) contribution, first we use the expression for $\Lambda_{\lambda\rho}$ from Eq. (5.37) to find

$$\begin{aligned} \Lambda'_{\lambda\rho}(k, q) &= \eta_{\lambda\rho}(q^2 - 2k \cdot q)(\not{k} - 2m) + (k_\lambda q_\rho + k_\rho q_\lambda)(\not{k} - 4m) + k_\lambda \not{k} \gamma_\rho \not{q} + k_\rho \not{k} \gamma_\lambda \not{q}, \\ \Lambda''_{\lambda\rho}(k, q) &= \eta_{\lambda\rho}(q^2 + 2k \cdot q)(\not{k} - 2m) - (k_\lambda q_\rho + k_\rho q_\lambda)(\not{k} - 4m) - k_\lambda \not{q} \gamma_\rho \not{k} - k_\rho \not{q} \gamma_\lambda \not{k}, \end{aligned} \quad (\text{E.31})$$

dropping irrelevant $O(q^2)$ -terms and using $k^2 = m^2$. In the $\eta_{\lambda\rho}$ terms, the integrand becomes independent of q . Thus, these terms give a regular contribution. Let us denote it by (A2r):

$$m'_{(A2r)} = \frac{2e^2}{m} \int \frac{d^3K}{(2\pi)^3 2E_K} \left[\frac{E_K - 2m}{E_K - m} f_e(E_K) + \frac{E_K + 2m}{E_K + m} f_{\bar{e}}(E_K) \right]. \quad (\text{E.32})$$

The terms which appear next will be called (A2s). For these, we use the fact that

$$(2v^\lambda v^\rho - \eta^{\lambda\rho})(k_\lambda q_\rho + k_\rho q_\lambda) = -2k \cdot q = 2\vec{K} \cdot \vec{Q}, \quad (\text{E.33})$$

using $q \cdot v = q_0 = 0$. The $Q \rightarrow 0$ limit can then be taken easily, and we obtain

$$m'_{(A2s)} = -\frac{e^2}{m} \int \frac{d^3K}{(2\pi)^3 2E_K} \left[\frac{E_K - 4m}{E_K - m} f_e(E_K) + \frac{E_K + 4m}{E_K + m} f_{\bar{e}}(E_K) \right]. \quad (\text{E.34})$$

Finally, we come to the terms with three γ -matrices, which we denote by (A2t). For these, first we note that

$$(2v^\lambda v^\rho - \eta^{\lambda\rho})(k_\lambda \not{k} \gamma_\rho \not{q} + k_\rho \not{k} \gamma_\lambda \not{q}) = 4k \cdot v \not{k} \not{q} - 2m^2 \not{q}, \quad (\text{E.35})$$

and a similar expression with the other term. Since the $\not{q}$ term vanishes between the spinors, we can write

$$\begin{aligned} m'_{(A2t)} &= 2e^2 \lim_{P \rightarrow 0} \lim_{Q \rightarrow 0} \int \frac{d^4 k}{(2\pi)^3} \delta(k^2 - m^2) \eta_f(k) \\ &\quad \times \frac{k_0}{m^2 - k \cdot p} \frac{1}{2\vec{K} \cdot \vec{Q}} \bar{u}(\vec{P}') (\not{k} \not{q} + \not{q} \not{k}) u(\vec{P}), \end{aligned} \quad (\text{E.36})$$

omitting the Q^2 terms in the denominator since they will not contribute for $Q \rightarrow 0$. Using the identity

$$\gamma_\kappa \gamma_\mu \gamma_\nu = \eta_{\kappa\mu} \gamma_\nu + \eta_{\mu\nu} \gamma_\kappa - \eta_{\kappa\nu} \gamma_\mu - i\varepsilon_{\kappa\mu\nu\alpha} \gamma^\alpha \gamma_5, \quad (\text{E.37})$$

we obtain

$$\not{k} \not{q} + \not{q} \not{k} = 2\vec{K} \cdot \vec{Q} \not{q} \quad (\text{E.38})$$

between the spinors, since $q \cdot v = 0$ and $\not{q}$ terms vanish. Putting this back into Eq. (E.36) and using Eqs. (4.58) and (4.59), we obtain

$$m'_{(A2t)} = -\frac{2e^2}{m} \int \frac{d^3 K}{(2\pi)^3 2E_K} \left[\frac{E_K}{E_K - m} f_e(E_K) + \frac{E_K}{E_K + m} f_{\bar{e}}(E_K) \right]. \quad (\text{E.39})$$

The sum of Eqs. (E.29), (E.30), (E.32), (E.34) and (E.39) gives the total contribution of the A2 term, given in Eq. (5.68) in the text.

E.3 The X contribution

The part of the integral from Eq. (5.82) that we consider here is given by

$$I^{(f)}(Q) = \int \frac{d^3 K}{(2\pi)^3} F(E_K) \left(\frac{1}{2\vec{K} \cdot \vec{Q} - Q^2} - \frac{1}{2\vec{K} \cdot \vec{Q} + Q^2} \right), \quad (\text{E.40})$$

where

$$F(E) \equiv (f_f(E) - f_{\bar{f}}(E))(2E^2 - m^2). \quad (\text{E.41})$$

Shifting the variables, the integral can be written as

$$I^{(f)}(Q) = \int \frac{d^3 K}{(2\pi)^3} \frac{F(E_{\vec{K}+\frac{1}{2}\vec{Q}}) - F(E_{\vec{K}-\frac{1}{2}\vec{Q}})}{2\vec{K} \cdot \vec{Q}}. \quad (\text{E.42})$$

We have to expand the numerator to $O(Q^3)$ in order to obtain the integral to $O(Q^2)$.

Writing ∂_i to denote a partial derivative with respect to K^i ,

$$F(E_{\vec{K}\pm\frac{1}{2}\vec{Q}}) = F(E) \pm \frac{1}{2}Q^i \partial_i F + \frac{1}{2} \left(\frac{1}{4}Q^i Q^j \right) \partial_i \partial_j F \pm \frac{1}{3!} \left(\frac{1}{8}Q^i Q^j Q^l \right) \partial_i \partial_j \partial_l F. \quad (\text{E.43})$$

The derivatives we need to use are:

$$\begin{aligned} \partial_i F &= K^i \left(\frac{1}{E} \frac{\partial}{\partial E} \right) F, \\ \partial_i \partial_j \partial_l F &= (\delta^{ij} K^l + \delta^{il} K^j + \delta^{jl} K^i) \left(\frac{1}{E} \frac{\partial}{\partial E} \right)^2 F + K^i K^j K^l \left(\frac{1}{E} \frac{\partial}{\partial E} \right)^3 F. \end{aligned} \quad (\text{E.44})$$

Using

$$K^i K^j \rightarrow \frac{1}{3} K^2 \delta^{ij} \quad (\text{E.45})$$

within the integrand, we have

$$I^{(f)}(Q) = \int \frac{d^3 K}{(2\pi)^3} \left[\frac{1}{2} \left(\frac{1}{E} \frac{\partial}{\partial E} \right) F + \frac{1}{3!} \frac{Q^2}{8} \left\{ 3 \left(\frac{1}{E} \frac{\partial}{\partial E} \right)^2 F + \frac{1}{3} K^2 \left(\frac{1}{E} \frac{\partial}{\partial E} \right)^3 F \right\} \right] \quad (\text{E.46})$$

Therefore, the quantity that we must substitute in Eq. (5.79) is

$$I^{(f)}(Q) - I^{(f)}(Q \rightarrow 0) = \frac{1}{3!} \frac{Q^2}{8} \int \frac{d^3 K}{(2\pi)^3} \left\{ 3 \left(\frac{1}{E} \frac{\partial}{\partial E} \right)^2 F + \frac{K^2}{3} \left(\frac{1}{E} \frac{\partial}{\partial E} \right)^3 F \right\} \quad (\text{E.47})$$

We now use the identity

$$\int_0^\infty dK K^n \left(\frac{1}{E} \frac{\partial}{\partial E} \right)^\nu F = -(n-1) \int_0^\infty dK K^{n-2} \left(\frac{1}{E} \frac{\partial}{\partial E} \right)^{\nu-1} F, \quad (\text{E.48})$$

which holds for $n \geq 2$, so that the surface term vanishes. It is obtained by using Eq. (5.122) and performing a partial integration. Using it repeatedly, we can rewrite Eq. (E.47) as

$$I^{(f)}(Q) - I^{(f)}(Q \rightarrow 0) = -\frac{Q^2}{48\pi^2} \int_0^\infty dK \left(\frac{1}{E} \frac{\partial}{\partial E} \right) F. \quad (\text{E.49})$$

Putting this back into Eq. (5.82), we obtain the total X contribution given in Eq. (5.83).

E.4 Alternative evaluation of the B1 and A2 contributions

If we straightaway put $q = 0$ in Eqs. (5.43) and (5.40), they are seen to involve singular expressions of the form $\mathcal{P}r(1/x)\delta(x)$ with $x = k^2$ and $x = k^2 - m^2$ respectively. Using Eqs. (5.60), we can obtain the following regularized form for such expressions [69]:

$$\begin{aligned}
\mathcal{P}r\left(\frac{1}{x}\right)\delta(x) &= \lim_{\epsilon \rightarrow 0} \frac{\epsilon x}{\pi(x^2 + \epsilon^2)^2} \\
&= \lim_{\epsilon \rightarrow 0} \lim_{\lambda \rightarrow 0} \frac{\epsilon(x - \lambda^2)}{\pi[(x - \lambda^2)^2 + \epsilon^2]^2} \\
&= \lim_{\epsilon \rightarrow 0} \lim_{\lambda \rightarrow 0} \frac{1}{2} \frac{\epsilon}{\pi} \frac{d}{d\lambda^2} \frac{1}{(x - \lambda^2)^2 + \epsilon^2} \\
&= \frac{1}{2} \lim_{\lambda \rightarrow 0} \frac{d}{d\lambda^2} \delta(x - \lambda^2). \tag{E.50}
\end{aligned}$$

We shall find that the use of Eq. (E.50) leads to expressions for $m'_{(B1)}$ and $m'_{(A2)}$ which are identical to the ones deduced earlier in this Appendix. Let us also compare Eq. (E.50) with Eq. (5.61). It is clear that Eq. (E.50) for $x = k^2$ is identical with Eq. (5.61) for $m \rightarrow 0$. For the case of $m \neq 0$, Eq. (5.61) is quite convenient to use if the rest of the expression in which it occurs does not involve m^2 (so that d/dm^2 can be performed at the end, as in Ref. [54]). But in our problem, the rest of the expression involves not only m^2 , but m and m^3 as well (see Eqs. (E.51) and (E.52) below). Consequently, we have found it convenient to use Eq. (E.50) with $x = k^2 - m^2$.

It may be noted that it is the form (E.50) which was essentially employed by DHR to regularize $\mathcal{P}r(1/k^2)\delta(k^2)$ in $\Gamma_{\lambda\rho}^{(B1)}(p, p)$ [70], who, however, made a detour into the imaginary-time formulation for this purpose. Here we first give the $m'_{(A2)}$ calculation, and then, for the sake of completeness, summarize the $m'_{(B1)}$ calculation.

Putting $q = 0$ in Eq. (5.40), we get

$$\begin{aligned}
\Gamma_{\lambda\rho}^{(A2)}(p, p) &= 2e^2 \int \frac{d^4k}{(2\pi)^3} \Lambda_{\lambda\rho}(k, k) \mathcal{P}r\left(\frac{1}{(k-p)^2}\right) \mathcal{P}r\left(\frac{1}{k^2 - m^2}\right) \delta(k^2 - m^2) \eta_f(k) \\
&= e^2 \int \frac{d^3K}{(2\pi)^3} \lim_{\lambda \rightarrow 0} \frac{d}{d\lambda^2} \int dk_0 \Lambda_{\lambda\rho}(k, k) \frac{1}{2m^2 + \lambda^2 - 2p \cdot k} \\
&\quad \times \delta(k^2 - m^2 - \lambda^2) \eta_f(k), \tag{E.51}
\end{aligned}$$

using Eq. (E.50) and also setting $k^2 = m^2 + \lambda^2$ as dictated by the delta function.

Making use of the delta function and Eqs. (4.58) and (4.59), we obtain, from Eq. (5.56),

$$(2v^\lambda v^\rho - \eta^{\lambda\rho}) \bar{u}(\vec{P}) \Lambda_{\lambda\rho}(k, k) u(\vec{P}) = -8k_0^3 + 20mk_0^2 - 8m^3 + 2k_0(2m^2 + \lambda^2 - 2mk_0) \tag{E.52}$$

at $v^\mu = (1, \vec{0})$ and $p^\mu = (m, \vec{0})$.

Use of Eq. (E.52), and of $p^\mu = (m, \vec{0})$, in Eq. (E.51) then leads to

$$m'_{(A2)} = e^2(-8I_1 + 20mI_2 - 8m^3I_3 + 2I_4), \quad (\text{E.53})$$

where

$$I_1 = \int [d^4k]_\lambda \frac{k_0^3}{2m^2 + \lambda^2 - 2mk_0}, \quad (\text{E.54})$$

$$I_2 = \int [d^4k]_\lambda \frac{k_0^2}{2m^2 + \lambda^2 - 2mk_0}, \quad (\text{E.55})$$

$$I_3 = \int [d^4k]_\lambda \frac{1}{2m^2 + \lambda^2 - 2mk_0}, \quad (\text{E.56})$$

$$I_4 = \int [d^4k]_\lambda k_0. \quad (\text{E.57})$$

We have used the notation

$$\begin{aligned} \int [d^4k]_\lambda \text{function}(k_0, \lambda) &\equiv \int \frac{d^3K}{(2\pi)^3} \lim_{\lambda \rightarrow 0} \frac{d}{d\lambda^2} \int dk_0 \delta(k_0^2 - E_{K\lambda}^2) \eta_f(k) \\ &\times \text{function}(k_0, \lambda), \end{aligned} \quad (\text{E.58})$$

with

$$E_{K\lambda} \equiv (K^2 + m^2 + \lambda^2)^{1/2}. \quad (\text{E.59})$$

Performing the k_0 integration first, and then removing the regulator λ in Eqs. (E.54) – (E.57) (it is useful to make note of

$$\lim_{\lambda \rightarrow 0} \frac{d}{d\lambda^2} f_{e,\bar{e}}(E_{K\lambda}) = \frac{1}{2E_K} \frac{\partial}{\partial E_K} f_{e,\bar{e}}(E_K), \quad (\text{E.60})$$

for this purpose), we arrive at

$$\begin{aligned} I_1 &= -\frac{1}{8m^2} \int \frac{d^3K}{(2\pi)^3} \left[\frac{E_K + 2m}{E_K - m} f_e(E_K) - \frac{E_K - 2m}{E_K + m} f_{\bar{e}}(E_K) \right. \\ &\quad \left. + mE_K \left(\frac{1}{E_K - m} \frac{\partial f_e}{\partial E_K} + \frac{1}{E_K + m} \frac{\partial f_{\bar{e}}}{\partial E_K} \right) \right], \end{aligned} \quad (\text{E.61})$$

$$\begin{aligned} I_2 &= -\frac{1}{8m^2} \int \frac{d^3K}{(2\pi)^3} \left[\frac{1}{E_K} \left(\frac{E_K + m}{E_K - m} f_e(E_K) + \frac{E_K - m}{E_K + m} f_{\bar{e}}(E_K) \right) \right. \\ &\quad \left. + m \left(\frac{1}{E_K - m} \frac{\partial f_e}{\partial E_K} - \frac{1}{E_K + m} \frac{\partial f_{\bar{e}}}{\partial E_K} \right) \right], \end{aligned} \quad (\text{E.62})$$

$$I_3 = -\frac{1}{8m^2} \int \frac{d^3 K}{(2\pi)^3} \left[\frac{1}{E_K^3} (f_e(E_K) + f_{\bar{e}}(E_K)) + \frac{m}{E_K^2} \left(\frac{1}{E_K - m} \frac{\partial f_e}{\partial E_K} - \frac{1}{E_K + m} \frac{\partial f_{\bar{e}}}{\partial E_K} \right) \right], \quad (\text{E.63})$$

$$I_4 = \frac{1}{4} \int \frac{d^3 K}{(2\pi)^3} \frac{1}{E_K} \frac{\partial}{\partial E_K} (f_e - f_{\bar{e}}). \quad (\text{E.64})$$

Substitution in Eq. (E.53) finally gives $m'_{(A2)}$, which is found to be the same as the result given by Eq. (5.68).

For $\Gamma_{\lambda\rho}^{(B1)}(p, p)$, given by Eq. (5.43), similar steps are to be carried out. Thus the analogues of Eqs. (E.51), (E.52) and (E.53) are

$$\begin{aligned} \Gamma_{\lambda\rho}^{(B1)}(p, p) &= e^2 \int \frac{d^3 K}{(2\pi)^3} \lim_{\lambda \rightarrow 0} \frac{d}{d\lambda^2} \int dk_0 \gamma^\nu (\not{p} - \not{k} + m) \gamma^\mu C_{\mu\nu\lambda\rho}(k, k) \\ &\quad \times \frac{1}{\lambda^2 - 2p \cdot k} \delta(k^2 - \lambda^2) \eta_\gamma(k), \end{aligned} \quad (\text{E.65})$$

$$(2v_\lambda v_\rho - \eta_{\lambda\rho}) \bar{u}(\vec{P}) \gamma^\nu (\not{p} - \not{k} + m) \gamma^\mu u(\vec{P}) C_{\mu\nu\lambda\rho}(k, k) = 2\lambda^2(k_0 - 2m) + 4mk_0^2 - 8k_0^3, \quad (\text{E.66})$$

$$m'_{(B1)} = e^2(2J_1 + 4mJ_2 - 8J_3). \quad (\text{E.67})$$

Here

$$J_1 = \int [d^4 k]_\lambda \frac{\lambda^2(k_0 - 2m)}{\lambda^2 - 2mk_0}, \quad (\text{E.68})$$

$$J_2 = \int [d^4 k]_\lambda \frac{k_0^2}{\lambda^2 - 2mk_0}, \quad (\text{E.69})$$

$$J_3 = \int [d^4 k]_\lambda \frac{k_0^3}{\lambda^2 - 2mk_0}. \quad (\text{E.70})$$

The notation $[d^4 k]_\lambda$ is now defined by Eq. (E.58) with $\eta_f \rightarrow \eta_\gamma$, and the notation $E_{K\lambda}$ by Eq. (E.59) with $m = 0$.

The evaluation of the above three integrals is straightforward. Each one reduces to the standard integral

$$\int_0^\infty dK K f_\gamma(K) = \frac{\pi^2 T^2}{6}, \quad (\text{E.71})$$

while J_3 also involves

$$\int_0^\infty dK K^2 \frac{e^{\beta K}}{(e^{\beta K} - 1)^2} = -\frac{\partial}{\partial \beta} \int_0^\infty dK K f_\gamma(K) \quad (\text{E.72})$$

We then have $J_1 = -T^2/24m$, $J_2 = -T^2/48m^2$ and $J_3 = T^2/48m$, so that the value of $m'_{(B1)}$ is the same as the one stated in Eq. (5.63).

Finally, we comment that the regularization presented above is not useful for evaluating the X -contribution, as we need the $O(Q^2)$ contribution from the fermion loop in that case, and not the the value at $q = 0$.

APPENDIX F

Thermal contribution to the equation for the effective mass at high temperature

In this appendix we arrive at Eq. (6.71), which expresses the high T contribution to the equation for the effective mass, by explicit calculation of the expressions given in Eqs. (6.62), (6.63), (6.65), (6.66) and (6.67).

We begin with $M'_{\text{FG}}{}^2$, given by Eq. (6.62). After the p_0 integration is carried out, the integrand is a function of P (the magnitude of the three-momentum) alone. Then integrating over the angles trivially, we are left with a one-dimensional integral. On doing some rearrangement, and making use of the identity

$$f_B(P) - f_F(P) = 2f_B(2P) \quad (\text{F.1})$$

(here and elsewhere in this appendix, $f_F(P)$ and $f_B(P)$ denote (6.11) and (6.15) with $|p_0|$ replaced by P), we obtain

$$M'_{\text{FG}}{}^2 = \frac{e^2}{2\pi^2} \int_0^\infty dP P [f_B(P) + f_F(P)] - \frac{e^2 M^2}{4\pi^2} \int_0^\infty dP P f_B(P) \text{Re} \frac{1}{P^2 - M^2 - i\epsilon}. \quad (\text{F.2})$$

Now the first part of the R.H.S. of Eq. (F.2) can be evaluated exactly by using

$$\int_0^\infty dP P f_B(P) = \frac{\pi^2 T^2}{6}, \quad (\text{F.3})$$

$$\int_0^\infty dP P f_F(P) = \frac{\pi^2 T^2}{12}. \quad (\text{F.4})$$

The second part of the R.H.S. of Eq. (F.2) involves the integral

$$I \equiv \int_0^\infty dP P \frac{1}{e^{P/T} - 1} \mathcal{P}r \frac{1}{P^2 - M^2} \quad (\text{F.5})$$

$$= \int_0^\infty dx x \frac{1}{e^x - 1} \mathcal{P}r \frac{1}{x^2 - (M/T)^2}. \quad (\text{F.6})$$

Here $\mathcal{P}r$ denotes the principal value. The functional dependence of I on T/M was determined on a computer for large T/M , yielding

$$I \approx -\frac{1}{2} \ln \frac{T}{M}. \quad (\text{F.7})$$

Note that this logarithmic behaviour is in accord with the ultraviolet behaviour of (F.5) without the distribution function. Thus at high T

$$M'_{\text{FG}}{}^2 = e^2 \left(\frac{T^2}{8} + \frac{M^2}{8\pi^2} \ln \frac{T}{M} \right) \quad (\text{F.8})$$

neglecting terms independent of T .

Next we turn to $M'_{\xi,F}{}^2$, given by Eq. (6.63). Integrating over p_0 and the angles, we are led to

$$\begin{aligned} M'_{\xi,F}{}^2 &= \frac{\xi e^2 M^2}{8\pi^2} \int_0^\infty dP P f_F(P) \operatorname{Re} \frac{1}{P^2 - \frac{M^2}{4} - \frac{i\epsilon}{2}} \\ &\quad + \frac{\xi e^2 M^4}{32\pi^2} \int_0^\infty dP P f_F(P) \operatorname{Re} \frac{1}{(P^2 - \frac{M^2}{4} - \frac{i\epsilon}{2})^2}. \end{aligned} \quad (\text{F.9})$$

Moving on to $M'_{\xi,B}{}^2$, integration over the angles in Eqs. (6.65), (6.66) and (6.67) lead to

$$M'_{\xi,B(\text{I})}{}^2 = -\frac{\xi e^2 M^2}{8\pi^2} \int_0^\infty dP P f_B(P) \operatorname{Re} \frac{1}{P^2 - \frac{M^2}{4} - \frac{i\epsilon}{2}}, \quad (\text{F.10})$$

$$M'_{\xi,B(\text{II})}{}^2 = -\frac{\xi e^2 M^4}{32\pi^2} \int_0^\infty dP P f_B(P) \operatorname{Re} \frac{1}{(P^2 - \frac{M^2}{4} - \frac{i\epsilon}{2})^2}, \quad (\text{F.11})$$

$$M'_{\xi,B(\text{III})}{}^2 = -\frac{\xi e^2 M^2 \beta}{8\pi^2} \int_0^\infty dP P^2 \frac{e^{\beta P}}{(e^{\beta P} - 1)^2} \operatorname{Re} \frac{1}{P^2 - \frac{M^2}{4} - \frac{i\epsilon}{2}}. \quad (\text{F.12})$$

Using the identity (F.1) we then arrive at

$$\begin{aligned} M'_{\xi,F}{}^2 + M'_{\xi,B(\text{I})}{}^2 + M'_{\xi,B(\text{II})}{}^2 &= -\frac{\xi e^2 M^2}{4\pi^2} \int_0^\infty dP P f_B(P) \operatorname{Re} \frac{1}{P^2 - M^2 - 2i\epsilon} \\ &\quad - \frac{\xi e^2 M^4}{4\pi^2} \int_0^\infty dP P f_B(P) \operatorname{Re} \frac{1}{(P^2 - M^2 - 2i\epsilon)^2}. \end{aligned} \quad (\text{F.13})$$

While the first part again involves (F.5), the second part, being ill-defined, is regularized in a way similar to (6.30):

$$\int_0^\infty dP P f_B(P) \operatorname{Re} \frac{1}{(P^2 - M^2 - 2i\epsilon)^2} = \lim_{\lambda \rightarrow M} \frac{\partial}{\partial \lambda^2} \int_0^\infty dP P f_B(P) \operatorname{Re} \frac{1}{P^2 - \lambda^2 - 2i\epsilon}. \quad (\text{F.14})$$

The integral on the R.H.S. of Eq. (F.14) is similar to I in Eq. (F.5) and equals $-\frac{1}{2} \ln(T/\lambda)$ for high T . So the second part of (F.13) equals $-\xi e^2 M^2/16\pi^2$ which can be neglected as it is independent of T . Thus we are left with the first part of (F.13), so that

$$M'_{\xi,F}{}^2 + M'_{\xi,B(I)}{}^2 + M'_{\xi,B(II)}{}^2 = \frac{\xi e^2 M^2}{8\pi^2} \ln \frac{T}{M}. \quad (\text{F.15})$$

In $M'_{\xi,B(III)}{}^2$, we encounter the integral

$$J \equiv \frac{1}{T} \int_0^\infty dP P^2 \frac{e^{P/T}}{(e^{P/T} - 1)^2} \mathcal{P}r \frac{1}{P^2 - M_1^2} \quad (\text{F.16})$$

where $M_1 = M/2$. Converting J into a function of T/M_1 alone (as we did in the case of I in Eq. (F.6)), the functional dependence was found out for large T/M_1 using a computer, which gave $J \approx -\frac{1}{2}$. This constancy again agrees with the expectation from the argument involving the ultraviolet behaviour. Alternatively one can note that

$$J = T \frac{\partial}{\partial T} \int_0^\infty dP P \frac{1}{e^{P/T} - 1} \mathcal{P}r \frac{1}{P^2 - M_1^2}. \quad (\text{F.17})$$

Then using our knowledge of I (see Eqs. (F.5) and F.7)), we obtain $J \approx -\frac{1}{2}$ at large T . Thus $M'_{\xi,B(III)}{}^2$ equals $\xi e^2 M^2/16\pi^2$ and can be neglected.

Therefore in the high- T limit, M'^2 receives contribution from (F.8) and (F.15), thereby leading us to Eq. (6.71).

BIBLIOGRAPHY

- [1] J. I. Kapusta, *Finite Temperature Field Theory* (Cambridge University Press, Cambridge, 1989).
- [2] M. Le Bellac, *Thermal Field Theory* (Cambridge University Press, Cambridge, 1996).
- [3] A. Das, *Finite temperature Field Theory* (World Scientific, Singapore, 1997).
- [4] L. Dolan and R. Jackiw, Phys. Rev. **D9**, 3320 (1974); S. Weinberg, Phys. Rev. **D9**, 3357 (1974); A. Linde, Rep. Prog. Phys. **42**, 389 (1979).
- [5] H. A. Weldon, Phys. Rev. **D26**, 2789 (1982).
- [6] H. A. Weldon, Phys. Rev. **D26**, 1394 (1982).
- [7] R. F. Sawyer, Phys. Rev. **D46**, 1180 (1992); J. C. D’Olivo, J. F. Nieves and P. B. Pal, Phys. Lett. **B365**, 178 (1996).
- [8] D. Notzold and G. Raffelt, Nucl. Phys. **B307**, 924 (1988); P. B. Pal and T. N. Pham, Phys. Rev. **D40**, 259 (1989); J. F. Nieves, Phys. Rev. **D40**, 866 (1989).
- [9] J. C. D’Olivo, J. F. Nieves and M. Torres, Phys. Rev. **D46**, 1172 (1992).
- [10] J. C. D’Olivo, J. F. Nieves and P. B. Pal, Phys. Rev. **D40**, 3679 (1989).
- [11] J. F. Nieves and P. B. Pal, Phys. Rev. **D58**, 096005 (1998).
- [12] J. C. D’Olivo, J. F. Nieves and P. B. Pal, Phys. Rev. Lett. **64**, 1088 (1990).
- [13] V. N. Oraevsky and V. B. Semikoz, Sov. J. Nucl. Phys. **42**, 446 (1985); Physica **A142**, 135 (1987); J. F. Nieves and P. B. Pal, Phys. Rev. **D49**, 1398 (1994).
- [14] J. F. Nieves and P. B. Pal, Phys. Rev. **D39**, 652 (1989); *ibid.* **40**, 2148(E) (1989); S. Mohanty, J. F. Nieves and P. B. Pal, Phys. Rev. **D58**, 093007 (1998).
- [15] K. Olive, Phys. Rep. **190**, 307 (1990).
- [16] L. Wolfenstein, Phys. Rev. **D17**, 2369 (1978); S. P. Mikheyev and A. Yu. Smirnov, Sov. J. Nucl. Phys. **42**, 913 (1985); for a recent review see, for example, G. G. Raffelt, Nucl. Phys. Proc. Suppl. **70**, 169 (1999).

- [17] E. Braaten, Phys. Rev. Lett. **66**, 1655 (1991).
- [18] E. Braaten and D. Segel, Phys. Rev. **D48**, 1478 (1993).
- [19] A. Kusenko and G. Segre, Phys. Rev. Lett. **77**, 4872 (1996); Phys. Lett. **B396**, 197 (1997).
- [20] J. F. Donoghue, B. R. Holstein and R. W. Robinett, Phys. Rev. **D30**, 2561 (1984).
- [21] J. F. Donoghue, B. R. Holstein, and R. W. Robinett, Phys. Rev. **D34**, 1208 (1986).
- [22] I. Mitra, J. F. Nieves and P. B. Pal, Phys. Rev **D64**, 085004 (2001).
- [23] I. Mitra, Phys. Rev. **D62**, 045023 (2000).
- [24] R. Kobes, G. Kunstatter and A. Rebhan, Phys. Rev. Lett. **64**, 2992 (1990).
- [25] E. Braaten and R. Pisarski, Nucl. Phys. **B337**, 569 (1990).
- [26] J. Frenkel and J. C. Taylor, Nucl. Phys. **B334**, 199 (1990).
- [27] E. Braaten and R. Pisarski, Phys. Rev. **D42**, 2156 (1990).
- [28] N. P. Landsman and Ch. G. van Weert, Phys. Rep. **145**, 141 (1987).
- [29] T. Altherr, Int. J. Mod. Phys. **A8**, 5605 (1993).
- [30] J. F. Nieves, Phys. Rev. **D42**, 4123 (1990).
- [31] See also M. Le Bellac (Ref. [2]), Appendix B.
- [32] For the derivations, in the canonical approach, see J. F. Nieves (Ref. [30]). Useful background material is to be found in C. Itzykson and J. -B. Zuber, *Quantum Field Theory* (McGraw-Hill, New York, 1980), Sec. 6-1-1.
- [33] For a simple example, see M. Le Bellac (Ref. [2]), p. 55.
- [34] S. Weinberg, *Gravitation and Cosmology: Principles and Applications of the General Theory of Relativity* (John Wiley, New York, 1972), Chapter 10.
- [35] R. Adler, M. Bazin and M. Schiffer, *Introduction to General Relativity* (2nd ed., McGraw-Hill, Tokyo, 1975), Chapter 9.
- [36] See S. Weinberg (Ref. [34]), p. 165.
- [37] F. A. Berends and R. Gastmans, Nucl. Phys. **B88**, 99 (1975). Note that $g_{\mu\nu} = \eta_{\mu\nu} + \kappa h_{\mu\nu}$ and $\kappa = \sqrt{32\pi G}$ in this reference, and also in Refs. [43] and [44].
- [38] M. E. Peskin and D. V. Schroeder, *An Introduction to Quantum Field Theory* (Addison-Wesley, Reading, Massachusetts, 1995), p. 126.

- [39] S. Weinberg (Ref. [34]), Sec. 12.5.
- [40] S. Weinberg, *The Quantum Theory of Fields, Volume III, Supersymmetry* (Cambridge University Press, Cambridge, 2000), Chapter 31.
- [41] M. Green, J. Schwartz and E. Witten, *Superstring Theory* (Cambridge University Press, Cambridge, 1987), Secs. 4.3.4 and 12.1.
- [42] J. F. Nieves and P. B. Pal, hep-ph/9812268.
- [43] H. T. Cho and K. L. Ng, Phys. Rev. **D47**, 1692 (1993).
- [44] S. Y. Choi, J. S. Shim and H. S. Song, Phys. Rev. **D51**, 2751 (1995).
- [45] J. C. D’Olivo and J. F. Nieves, Phys. Rev. **D52**, 2987 (1995).
- [46] H. A. Weldon, Phys. Rev. **D40**, 2410 (1989).
- [47] H. A. Weldon, Physica **A158**, 169 (1989).
- [48] E. Braaten, Astrophys. J. **392**, 70 (1992).
- [49] See F. Mandl and G. Shaw, *Quantum Field Theory* (John Wiley, New York, 1984), p. 303, and reference therein.
- [50] See J. J. Sakurai, *Advanced Quantum Mechanics* (Addison-Wesley, Reading, Massachusetts, 1967), p. 285. For the case of photon in a medium, see the Appendices of Ref. [18]; see also M. Le Bellac (Ref. [2]), p. 219.
- [51] See, for example, S. Weinberg (Ref. [34]), Sec. 3.4.
- [52] For the analogous case of electric charge in a medium, see Appendix C of Ref. [46], and Appendix E of H. A. Weldon, Phys. Rev. **D61**, 036003 (2000).
- [53] See, for example, H. A. Weldon, Phys. Rev. **D26**, 1394 (1982) and **D47**, 594 (1993); P. S. Gribosky and B. R. Holstein, Z. Phys. **C47**, 205 (1990); H. Matsumoto, Y. Nakano and H. Umezawa, Phys. Rev. **D31**, 1495 (1985); Y. Fujimoto and H. Yamada, Z. Phys. **C37**, 265 (1988); T. S. Evans, Z. Phys. **C41**, 333 (1988); P. F. Bedaque and A. Das, Phys. Rev. **D45**, 2906 (1992); P. Arnold, S. Vokos, P. Bedaque and A. Das, Phys. Rev. **D47**, 4698 (1993); J. F. Nieves and P. B. Pal, Phys. Rev. **D51**, 5300 (1995).
- [54] P. F. Bedaque and A. Das (Ref. [53]).
- [55] E. M. Lifshitz and L. P. Pitaevskii, *Physical Kinetics* (Pergamon Press, Oxford, 1981), p. 115.
- [56] E. M. Lifshitz and L. P. Pitaevskii (Ref. [55]), p. 162.

- [57] G. C. Raffelt, *Stars as Laboratories for Fundamental Physics* (Chicago University Press, Chicago, 1996), p. 597.
- [58] It may be noted that Refs. [5] and [23] use upper-case letter for the electron four-momentum, and define a and b with an extra minus sign.
- [59] See Chapter 4 of this work, and also Sec. II of Ref. [23].
- [60] For derivation of the expression (6.14) for the gauge dependent part of the photon propagator, see K. L. Kowalski, Z. Phys. **C36**, 665 (1987). This expression has been used by P. Elmfors, D. Persson and B-S. Skagerstam, Nucl. Phys. **B464**, 153 (1996) to prove gauge independence of the strong-field limit of the one-loop electron dispersion relation in a medium with a magnetic field.
- [61] One way of making $\partial f/\partial\lambda^2$ at $\lambda = 0$ finite in the present case is to use in it a regularization similar to the one given in (6.30), and keep the regulator (small but) non-zero.
- [62] We reserve the symbol M for the effective electron mass including the subleading T -dependence.
- [63] Similar idea has been used for neutrino dispersion relation in Ref. [9].
- [64] See footnote 4 of Ref. [5].
- [65] A. Erdas, C. W. Kim and J. A. Lee, Phys. Rev. **D48**, 3901 (1993).
- [66] See Appendix of H. A. Weldon (Ref. [6]).
- [67] F. T. Brandt and J. Frenkel, Phys. Rev. **D55**, 7808 (1997).
- [68] See, for example, C. Itzykson and J. -B. Zuber (Ref. [32]), Sec. 3-1-5.
- [69] Alternatively, take the real part of the relation (3.104) of M. Le Bellac (Ref. [2]), and use Eq. (6.31) of the present work in it.
- [70] The finite-temperature part in Eq. (A13) of the Appendix of Ref. [20] is the same as our Eq. (E.65).

LIST OF PUBLICATIONS

1. I. Mitra, J. F. Nieves and P. B. Pal, “Gravitational couplings of charged leptons in a medium,” *Phys. Rev* **D64**, 085004 (2001) [hep-ph/0104248].
2. I. Mitra, “Gauge independence of limiting cases of one-loop electron dispersion relation in high-temperature QED,” *Phys. Rev.* **D62**, 045023 (2000) [hep-ph/9905362].
3. I. Mitra, A. DasGupta, and B. Dutta-Roy, “Regularization and renormalization in scattering from Dirac delta potentials,” *Am. J. Phys.* **66**, 1101 (1998): Not included in this thesis.